

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
10 October 2002 (10.10.2002)

PCT

(10) International Publication Number
WO 02/078693 A2

(51) **International Patent Classification⁷:** **A61K 31/4045**,
C07D 209/16, A61P 25/18, 25/28, C07D 403/12, 401/12,
A61K 31/4406, 31/4409, 31/506, 31/4402, C07C 217/54,
323/01, 317/14, C07D 213/64, 277/34, 239/34, 233/24,
333/20, 307/91, 209/14

(21) **International Application Number:** PCT/US02/05115

(22) **International Filing Date:** 15 March 2002 (15.03.2002)

(25) **Filing Language:** English

(26) **Publication Language:** English

(30) **Priority Data:**
60/279,928 29 March 2001 (29.03.2001) US
60/329,449 15 October 2001 (15.10.2001) US

(71) **Applicant (for all designated States except US):** **ELI LILLY AND COMPANY** [US/US]; Lilly Corporate Center, Indianapolis, IN 46285 (US).

(72) **Inventors; and**

(75) **Inventors/Applicants (for US only):** **CHEN, Zhaogen** [CN/US]; 9972 Parkshore Drive, Noblesville, IN 46060 (US). **COHEN, Michael, Philip** [US/US]; 8141 Bowline Court, Indianapolis, IN 46236 (US). **FISHER, Matthew, Joseph** [US/US]; 10650 Quail Ridge Court, Mooresville, IN 46158 (US). **GIETHLEN, Bruno** [FR/FR]; Prestwick Chemical, Inc., Bat. Tycho Brahe, Rue Tobias Stimmer, F-67400 Illkirch-Graffenstaden (FR). **GILLIG, James, Ronald** [US/US]; 3634 Toronto Court, Indianapolis, IN 46268 (US). **MCCOWAN, Jefferson, Ray** [US/US]; 2653 Crescent Hill Lane, Indianapolis, IN 46228 (US). **MILLER, Shawn, Christopher** [US/US]; 7359 Hardin Oak Drive, Noblesville, IN 46060 (US). **SCHAUS, John, Mehnert** [US/US]; 135 Raintree Drive, Zionsville, IN 46077 (US).

(74) **Agents:** **DAWALT, Elizabeth, A.** et al.; Eli Lilly and Company, Lilly Corporate Center, Indianapolis, IN 46285 (US).

(81) **Designated States (national):** AE, AG, AL, AM, AT, AT (utility model), AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, CZ (utility model), DE, DE (utility model), DK, DK (utility model), DM, DZ, EC, EE,

EE (utility model), ES, FI, FI (utility model), GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SK (utility model), SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.


(84) **Designated States (regional):** ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

- as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii)) for the following designations AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZM, ZW, ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG)
- as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii)) for the following designations AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZM, ZW, ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG)

[Continued on next page]

(54) **Title:** N-(2-ARYLETHYL)BENZYLAMINES AS ANTAGONISTS OF THE 5-HT₆ RECEPTOR



(57) **Abstract:** The present invention provides compounds of formula (I), which are antagonists of the 5-HT₆ receptor.

**Published:**

— without international search report and to be republished
upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

-1-

N-(2-ARYLETHYL)BENZYLAMINES AS
ANTAGONISTS OF THE 5-HT₆ RECEPTOR

5 The present invention relates to the field of pharmaceutical and organic chemistry and is concerned with compounds which are antagonists of the 5-HT₆ receptor.

10 The 5-HT₆ receptor is a member of the G-protein coupled receptor superfamily of serotonin receptors, and, like the 5-HT₄ and 5-HT₇ receptors, it is positively coupled to adenylate cyclase.¹ The rat 5-HT₆ receptor was first cloned in 1993^{2,3} and the cloning of the human homologue, to which it shares a 89% sequence identity, was reported in 1996.⁴ The localization of 5-HT₆ receptors in rat brain has been studied using mRNA quantification by Northern analysis and RT-PCR, immunohistochemistry, and autoradiography.^{2, 3, 5, 6, 7, 8} These methods have consistently found high levels of the receptor in olfactory tubercle, hippocampus, striatum, nucleus accumbens, and cortical
15 regions. 5-HT₆ receptors are either absent or present in very low levels in peripheral tissues.^{2, 3}

To date, there are no known high affinity, selective agonists at the 5-HT₆ receptor. Serotonin itself has only moderate affinity for the 5-HT₆ receptor (K_i = 65 nM) and the most selective agonist reported to date, N,N-dimethyl-2-ethyl-5-methoxytryptamine, has
20 K_i = 81 nM and only 3.5-fold selectivity versus the 5-HT_{2A} receptor.⁹

Much of the recent interest in the 5-HT₆ receptor is due to the observation that several psychotropic agents are high affinity antagonists at the human 5-HT₆ receptor.^{4, 10} These compounds include amitriptyline (K_i = 65 nM) and the atypical antipsychotics clozapine (K_i = 9.5 nM), olanzapine (K_i = 10 nM), and quetiapine (33 nM). None of these
25 compounds, however, are selective. The first selective 5-HT₆ receptor antagonists reported are Ro 04-6790 and Ro 63-0563. Their usefulness is limited by their moderate affinity (K_i = 50nM and 12 nM, respectively) and poor pharmacokinetics.¹¹ A series of 5-HT₆ receptor antagonists, culminating in SB-271,046, has been reported.¹² This compound has high affinity (K_i = 1.2 nM) and selectivity (>200-fold versus >55
30 receptors, enzymes and ion channels) and is 80% bioavailable. A selective radioligand [¹²⁵I]-SB-258,585 has been used for radioligand binding and autoradiographic studies.^{13, 14} These compounds are useful tools for preclinical studies on the 5-HT₆ receptor.

The rationale for the use of selective 5-HT₆ receptor antagonists to treat cognitive dysfunction is based on three lines of reasoning: the ability of selective 5-HT₆ receptor
35 antagonists to modulate cholinergic and glutamatergic neuronal function, clinical studies of the atypical antipsychotics clozapine and olanzapine on cognitive function, the activity of selective 5-HT₆ receptor antagonists in animal models of cognitive function.

-2-

Selective 5-HT₆ receptor antagonists modulate cholinergic and glutamatergic neuronal function. Cholinergic and glutamatergic neuronal systems play important roles in cognitive function. Cholinergic neuronal pathways are known to be important to memory formation and consolidation. Centrally acting anticholinergic agents impair cognitive function in animal and clinical studies and loss of cholinergic neurons is one of the hallmarks of Alzheimer's disease. Conversely, stimulation of cholinergic function has been known to improve cognitive performance and the only two agents currently approved for the treatment of cognitive deficit in Alzheimer's disease, tacrine and donepezil, are both acetylcholinesterase inhibitors. The glutamatergic system in the prefrontal cortex is also known to be involved in cognitive function.^{26, 27}

Blocking 5-HT₆ receptor function has been shown to elicit procholinergic effects *in vivo*. Administration (icv) to rats of antisense oligonucleotides targeting the 5-HT₆ receptor sequence induced yawning and stretching behavior that was blocked by the cholinergic antagonist atropine.¹⁵ The selective 5-HT₆ receptor antagonist Ro 04-6790 induced stretching behavior in a dose-dependent manner. This behavior was blocked by the centrally acting anticholinergic agents scopolamine and atropine but not by methylscopolamine at doses known to be peripherally selective.¹⁶ Ro 04-6790 was also shown to block the rotational behavior induced by scopolamine administration to rats with unilateral nigrostriatal 6-OH-DA lesions. It did not block rotational behavior induced by L-DOPA or amphetamine.¹⁷ Ro 04-6790 reversed scopolamine induced performance deficits in the object recognition test, a model of cognitive function. Another selective 5-HT₆ receptor antagonist, SB-271046, potentiated the yawning behavior induced by the cholinesterase inhibitor physostigmine.¹⁸ These studies suggest that 5-HT₆ receptor blockade facilitates cholinergic transmission. In *in vivo* microdialysis studies, SB-271,046 (10 mg/kg, sc) increases glutamate release in the prefrontal cortex through a neuronal mechanism.²⁵

Clinical studies of the atypical antipsychotics clozapine and olanzapine on cognitive function. The atypical antipsychotics clozapine and olanzapine are both high affinity, albeit nonselective, 5-HT₆ receptor antagonists.⁴ On the other hand, risperidone and the typical antipsychotic haloperidol do not have significant affinity for the 5-HT₆ receptor. Clinical differences seen with these sets of drugs may be attributable to 5-HT₆ receptor blockade. Goldberg et al. reported no beneficial cognitive effect of clozapine treatment in a small (N = 15) trial in treatment resistant schizophrenics.¹⁹ In contrast, Meltzer et al.²⁰ in a larger study of treatment-resistant schizophrenics (N = 36), observed improvements in several domains of neuropsychological function at six weeks and six months following initiation of clozapine treatment. In non-treatment resistant schizophrenics, clozapine was more effective than placebo in improving cognitive

-3-

function by several measures.²¹ This effect was seen at six months and persisted throughout the 12 month study. The effect of olanzapine, risperidone, and haloperidol on cognitive function has been compared in a multicenter, double blind study in schizophrenics.²² The olanzapine group showed a statistically significant improvement in cognitive function over either haloperidol or risperidone treatment. This effect was apparent after 6 weeks treatment and continued throughout the 54 weeks of the study. Animal studies suggest that these effects could be mediated through the release of acetylcholine in the prefrontal cortex.²³

Activity of selective 5-HT₆ receptor antagonists in animal models of cognitive function. With the recent development of the selective 5-HT₆ receptor antagonists Ro-04,6790 and SB-271,046, there have been several reports on the activity of these compounds in models of cognitive function. The selective 5-HT₆ receptor antagonist SB-271,046 improved performance in the Morris water maze.²⁴ These results are consistent with the finding that chronic icv administration of antisense oligonucleotides directed toward the 5-HT₆ receptor sequence led to improvements in some measures of performance in the Morris water maze.¹⁶ SB-271,046 treatment also led to improvements in the spatial alternation operant behavior test in aged rats.²⁴

The compounds of the present invention are selective, high affinity antagonists of 5-HT₆, and thus, provide a valuable treatment for 5-HT₆ receptor mediated disorders.

Background References


1. Branchek, T. A., et al. (2000). *Annu Rev Pharmacol Toxicol* 40: 319-34.
2. Monsma, F. J., Jr., et al. (1993). *Mol Pharmacol* 43(3): 320-7.
3. Ruat, M., et al. (1993). *Biochem Biophys Res Commun* 193(1): 268-76.
4. Kohen, R., et al. (1996). *J Neurochem* 66(1): 47-56.
5. Ward, R. P., et al. (1996). *J Comp Neurol* 370(3): 405-14.
6. Ward, R. P., et al. (1995). *Neuroscience* 64(4): 1105-11.
7. Gerard, C., et al. (1997). *Brain Res* 746(1-2): 207-19.
8. Gerard, C., et al. (1996). *Synapse* 23(3): 164-73.
9. Glennon, R. A., et al. (2000). *J Med Chem* 43(5): 1011-8.
10. Roth, B. L., et al. (1994). *J Pharmacol Exp Ther* 268(3): 1403-10.
11. Sleight, A. J., et al. (1998). *Br J Pharmacol* 124(3): 556-62.
12. Routledge, C., et al. (2000). *Br. J. Pharmacol.* 130(7): 1606.
13. Hirst, W. D., et al. (1999). *Br. J. Pharmacol. Suppl.*((in press)).
14. Hirst, W. D., et al. (2000). *Br. J. Pharmacol.* 130: 1597.
15. Bourson, A., et al. (1995). *J Pharmacol Exp Ther* 274(1): 173-80.
16. Bentley, J. C., et al. (1999). *Br J Pharmacol* 126(7): 1537-42.

-4-

17. Bourson, A., et al. (1998). Br J Pharmacol 125(7): 1562-6.
18. Routledge, C., et al. (1999). Br. J. Pharmacol. 127(Suppl.): 21P.
19. Goldberg, T. E., et al. (1993). Br J Psychiatry 162: 43-8.
20. Hagger, C., et al. (1993). Biol Psychiatry 34(10): 702-12.
- 5 21. Lee, M. A., et al. (1994). J Clin Psychiatry 55 Suppl B: 82-7.
22. Purdon, S. E., et al. (2000). Arch Gen Psychiatry 57(3): 249-58.
23. Parada, M. A., et al. (1997). J Pharmacol Exp Ther 281(1): 582-8.
24. Rogers, D. C., et al. (1999). Br J Pharmacol 127(suppl.): 22P.
25. Dawson, L. A., et al. (2000). Br J Pharmacol 130(1): 23-6.
- 10 26. Dudkin, K. N., et al. (1996). Neurosci Behav Physiol 26(6): 545-51.
27. Koechlin, E., et al. (1999). Nature 399(6732): 148-51.

The present invention provides compounds of

15 formula I:



formula I


wherein

X is selected from the group consisting of -O-, -NH-, -S-, -SO₂-, -CH₂-, -CH(F)-, -CH(OH)-, and -C(O)-;

- 20 R₁ is selected from the group consisting of optionally substituted phenyl, optionally substituted naphthyl, optionally substituted 5 to 6 membered monocyclic aromatic heterocycle having one heteroatom selected from the group consisting of nitrogen, oxygen, and sulfur and which 5 to 6 membered monocyclic aromatic heterocycle is optionally benzofused;
- 25 R₂ is selected from the group consisting of hydrogen and C₁-C₃ alkyl;
- R₃ is selected from the group consisting of hydrogen, fluoro, and methyl;
- R₄ is selected from the group consisting of hydrogen, allyl, C₂-C₄ alkyl, fluorinated C₂-C₄ alkyl, optionally substituted phenyl, optionally substituted phenylsulfonyl, optionally substituted benzyl, and optionally substituted 5 to 6 membered monocyclic aromatic
- 30 heterocycle having one or two heteroatoms selected from the group consisting of nitrogen, oxygen, and sulfur, provided that R₄ is not optionally substituted phenylsulfonyl when X is -SO₂-, -CH₂-, -CH(F)-, -CH(OH)-, or -C(O)-; and pharmaceutically acceptable salts thereof.

The present invention also provides compounds of formula II:

-5-



formula II

wherein

Y is selected from the group consisting of O, NH, and NR₉, wherein R₉ is selected from the group consisting of C₁-C₄ alkyl, and optionally substituted phenyl;

5 R₅ and R₆ are hydrogen or taken together with the atoms to which they are attached form a benzo ring, provided that R₅ and R₆ are hydrogen when Y is NR₉;

R₇ is selected from the group consisting of optionally substituted phenyl, optionally substituted naphthyl, optionally substituted 5 to 6 membered monocyclic aromatic heterocycle having one heteroatom selected from the group consisting of nitrogen,
10 oxygen, and sulfur and which 5 to 6 membered monocyclic aromatic heterocycle is optionally benzofused;

R₈ is selected from the group consisting of hydrogen and C₁-C₃ alkyl; and pharmaceutically acceptable salts thereof.

15 The present invention also provides for novel pharmaceutical compositions, comprising: a compound of the formula I or II and a pharmaceutically acceptable diluent.

Because the compounds of formula I and II are antagonists of 5-HT₆ receptor, the compounds of formula I and II are useful for the treatment of a variety of disorders. Thus, in another embodiment the present invention provides methods of treating disorders
20 associated with 5-HT₆, comprising: administering to a patient in need thereof an effective amount of a compound of formula I or II. That is, the present invention provides for the use of a compound of formula I or II and pharmaceutical compositions thereof for the treatment disorders associated with 5-HT₆. More specifically, the present invention provides a method of treating disorders selected from the group consisting of cognitive
25 disorders, age-related cognitive disorder, mild cognitive impairment, mood disorders (including depression, mania, bipolar disorders), psychosis (in particular schizophrenia), anxiety (particularly including generalized anxiety disorder, panic disorder, and obsessive compulsive disorder), idiopathic and drug-induced Parkinson's disease, epilepsy, convulsions, migraine (including migraine headache), substance withdrawal (including, substances such as opiates, nicotine, tobacco products, alcohol, benzodiazepines, cocaine,
30 sedatives, hypnotics, etc.), sleep disorders (including narcolepsy), attention deficit/hyperactivity disorder, conduct disorder, learning disorders, dementia (including Alzheimer's disease and AIDS-induced dementia), Huntington's Chorea, cognitive deficits

-6-


subsequent to cardiac bypass surgery and grafting, stroke, cerebral ischemia, spinal cord trauma, head trauma, perinatal hypoxia, cardiac arrest, and hypoglycemic neuronal damage, vascular dementia, multi-infarct dementia, amyotrophic lateral sclerosis, and multiple sclerosis, comprising: administering to a patient in need thereof an effective amount of a compound of formula I or an effective amount of a compound of formula II.

In addition, the present invention also provides processes for preparing the compounds of formula I and II and intermediate thereof.

As used herein, the following terms have the meanings indicated:


The term "C₁-C₃ alkyl" refers to a straight or branched alkyl chain having from one to three carbon atoms, and includes methyl, ethyl, propyl, and iso-propyl.

The term "optionally substituted phenyl" refers to a radical of the formula




wherein R_a is from 1 to 3 groups independently selected from the group consisting of hydrogen, hydroxy, C₁-C₄ alkyl, C₁-C₄ alkoxy, halogen, benzyloxy, carboxy, C₁-C₄ alkoxycarbonyl, amido, N-(C₁-C₄ alkyl)amido, sulfonylamido, cyano, trifluoromethyl, trifluoromethoxy, nitro, and phenyl optionally substituted with C₁-C₄ alkyl, C₁-C₄ alkoxy, halogen, cyano, and trifluoromethyl.

The term "optionally substituted naphthyl" refers to a radical of the formula



wherein R_c is from 1 to 2 groups independently selected from the group consisting of hydrogen, C₁-C₄ alkyl, C₁-C₄ alkoxy, halogen, cyano, trifluoromethyl, and nitro.

The term "optionally substituted 5 to 6 membered monocyclic aromatic heterocycle having one heteroatom selected from the group consisting of nitrogen, oxygen, and sulfur and which 5 to 6 membered monocyclic aromatic heterocycle is optionally benzofused" refers to radicals of the formula




wherein Q₁ is selected from the group consisting of -O-, -S-, and -NR_g- wherein R_g is selected from the group consisting of hydrogen and C₁-C₄ alkyl; and Q₂ is -N=, R_d, each R_e, and R_f are each independently selected from the group consisting of hydrogen, C₁-C₄ alkyl, C₁-C₄ alkoxy, halogen, cyano, and trifluoromethyl, or R_d and R_e (or one of R_e) are taken together with the atoms to which they are attached to form an benzo ring which

-7-

benzo ring is optionally substituted with 1 to 4 substituents independently selected from the group consisting of hydrogen, hydroxy, C₁-C₄ alkyl, C₁-C₄ alkoxy, trifluoromethoxy, 2,2,2-trifluoroethoxy, trifluoromethyl, halogen, carboxy, C₁-C₄ alkoxycarbonyl, amido, N-(C₁-C₄ alkyl)amido, amino, (C₁-C₄ alkyl)amino, acylamino wherein the acyl group is
 5 selected from the group consisting of C₁-C₄ alkyl and phenyl; cyano, nitro, sulfonylamido, phenyl optionally substituted with C₁-C₄ alkyl, C₁-C₄ alkoxy, halogen, cyano, and trifluoromethyl; phenoxy, benzyloxy, -NHS(O)₂R_h, wherein R_h is selected from the group consisting of C₁-C₄ alkyl and phenyl; and -S(O)_pR_i, wherein p is 0, 1, or 2 and R_i is
 10 C₁-C₄ alkyl, C₁-C₄ alkoxy, halogen, cyano, and trifluoromethyl; and R_f is selected from the group consisting of hydrogen, C₁-C₄ alkyl, C₁-C₄ alkoxy, trifluoromethyl, and halogen. The term specifically includes furyl, thienyl, pyrrolyl, pyridyl, benzofuryl, benzothienyl, indolyl and quinoliny; each optionally substituted as described above.

The term "fluorinated C₂-C₄ alkyl" refers to a straight or branched alkyl chain
 15 having from two to four carbon atoms substituted with one or more fluorine atoms. The term includes 2-fluoroethyl, 2,2-difluoroethyl, 2,2,2-trifluoroethyl, 3-fluoropropyl, 3,3-difluoropropyl, 3,3,3-trifluoropropyl, 2,2,3,3,3-pentafluoropropyl, 2,2,3,3-tetrafluoropropyl, 4,4,4-trifluorobutyl, 3,3,4,4,4-pentafluorobutyl, and the like.


The term "optionally substituted phenylsulfonyl" refers to a radical of the formula



20

wherein R_j is from 1 to 3 groups independently selected from the group consisting of hydrogen, C₁-C₄ alkyl, C₁-C₄ alkoxy, halogen, cyano, trifluoromethyl, nitro, and phenyl.


The term "optionally substituted benzyl" refers to a radical of the formula



25

wherein R_k is from 1 to 3 groups independently selected from the group consisting of hydrogen, C₁-C₄ alkyl, C₁-C₄ alkoxy, cyano, nitro, trifluoromethyl, and halogen.

The term "optionally substituted 5 to 6 membered monocyclic aromatic heterocycle having one or two heteroatoms selected from the group consisting of nitrogen,
 30 oxygen, and sulfur" refers to radicals of the formula



-8-

wherein Q_3 is selected from the group consisting of -O-, -S-, and -NR_g- wherein R_g is selected from the group consisting of hydrogen and C₁-C₄ alkyl; and Q₄ and Q₅ are -CR_m, wherein each R_m is independently selected from the group consisting of hydrogen, C₁-C₄ alkyl, halogen, and trifluoromethyl or one or both of Q₄ and Q₅ is -N=; and wherein one or two of Q₆ are -N=, while the others are -CR_n; wherein each R_n is independently selected from the group consisting of hydrogen, C₁-C₄ alkyl, C₁-C₄ alkoxy, halogen, cyano, nitro, and trifluoromethyl. The term specifically includes furyl, thienyl, thiazolyl, pyrazolyl, imidazolyl, triazolyl, oxazolyl, thiazolyl, isoxazolyl, thioisoxazolyl, pyridyl, pyrimidyl, pyridazinyl, and pyrazidiny; each optionally substituted as described above.

10 The term "C₁-C₄ alkyl" refers to a straight or branched alkyl chain having from one to four carbon atoms, and includes methyl, ethyl, propyl, iso-propyl, butyl, sec-butyl, iso-butyl, and t-butyl.

The term "C₂-C₄ alkyl" refers to a straight or branched alkyl chain having from two to four carbon atoms, and includes ethyl, propyl, iso-propyl, butyl, sec-butyl, iso-butyl, and t-butyl.

15

The term "C₁-C₄ alkoxy" refers to a straight or branched alkyl chain having from one to four carbon atoms attached to an oxygen atom, and includes methoxy, ethoxy, propoxy, iso-propoxy, butoxy, iso-butoxy, sec-butoxy, and t-butoxy.

The term "halogen" refers to a chloro, fluoro, bromo or iodo atom.

20 The term "pharmaceutically-acceptable addition salt" refers to an acid addition salt.

The compound of formula I or II and the intermediates described herein form pharmaceutically acceptable acid addition salts with a wide variety of organic and inorganic acids and include the physiologically acceptable salts which are often used in pharmaceutical chemistry. Such salts are also part of this invention. A pharmaceutically-acceptable addition salt is formed from a pharmaceutically-acceptable acid as is well known in the art. Such salts include the pharmaceutically acceptable salts listed in Journal of Pharmaceutical Science, 66, 2-19 (1977) which are known to the skilled artisan. Typical inorganic acids used to form such salts include hydrochloric, hydrobromic, hydriodic, nitric, sulfuric, phosphoric, hypophosphoric, metaphosphoric, pyrophosphoric, and the like. Salts derived from organic acids, such as aliphatic mono and dicarboxylic acids, phenyl substituted alkanoic acids, hydroxyalkanoic and hydroxyalkandioic acids, aromatic acids, aliphatic and aromatic sulfonic acids, may also be used. Such pharmaceutically acceptable salts thus include chloride, bromide, iodide, nitrate, acetate, phenylacetate, trifluoroacetate, acrylate, ascorbate, benzoate, chlorobenzoate, dinitrobenzoate, hydroxybenzoate, methoxybenzoate, methylbenzoate, o-acetoxybenzoate, isobutyrate, phenylbutyrate, α-hydroxybutyrate, butyne-1,4-

25

30

35

dicarboxylate, hexyne-1,4-dicarboxylate, caprate, caprylate, cinnamate, citrate, formate, fumarate, glycollate, heptanoate, hippurate, lactate, malate, maleate, hydroxymaleate, malonate, mandelate, mesylate, nicotinate, isonicotinate, oxalate, phthalate, teraphthalate, propiolate, propionate, phenylpropionate, salicylate, sebacate, succinate, suberate, 5 benzenesulfonate, p-bromobenzenesulfonate, chlorobenzenesulfonate, ethylsulfonate, 2-hydroxyethylsulfonate, methylsulfonate, naphthalene-1-sulfonate, naphthalene-2-sulfonate, naphthalene-1,5-sulfonate, p-toluenesulfonate, xylenesulfonate, tartrate, and the like.

As with any group of pharmaceutically active compounds, some groups are 10 preferred in their end use application. Preferred embodiments of the present invention are given for the compounds of formula I below:

Compounds in which wherein X is selected from the group consisting of -O-, -NH-, and -S- are preferred, with compounds in which X is -O- being more preferred.

Compounds in which R₁ is optionally substituted phenyl or optionally substituted 15 5 to 6 membered monocyclic aromatic heterocycle having one heteroatom selected from the group consisting of nitrogen, oxygen, and sulfur and which 5 to 6 membered monocyclic aromatic heterocycle is optionally benzofused are preferred.

When R₁ is optionally substituted phenyl preferred substituents are 1 to 3 groups 20 independently selected from the group consisting of hydrogen, C₁-C₄ alkyl, halogen, benzyloxy, carboxy, C₁-C₄ alkoxycarbonyl, amido, N-(C₁-C₄ alkyl)amido, sulfonylamido, cyano, trifluoromethyl, trifluoromethoxy, nitro, and phenyl optionally substituted with C₁-C₄ alkyl, C₁-C₄ alkoxy, halogen, cyano, and trifluoromethyl.


When R₁ is optionally substituted phenyl more preferred substituents are 1 to 3 25 groups independently selected from the group consisting of hydrogen, C₁-C₄ alkyl, halogen, cyano, and trifluoromethyl.

Compounds in which R₃ is hydrogen or fluorine are preferred.

Compound in which R₁ is optionally substituted 5 to 6 membered monocyclic aromatic heterocycle having one heteroatom selected from the group consisting of 30 nitrogen, oxygen, and sulfur and which 5 to 6 membered monocyclic aromatic heterocycle is optionally benzofused, the compounds which are benzo fused are preferred, with indolyl being preferred, and indol-3-yl being even more preferred.

When R₁ is indol-3-yl, preferred groups are depicted as the radical below:

-10-



- a) R_o is selected from the group consisting of hydrogen and C_1 - C_4 alkyl, with hydrogen being more preferred;
- b) R_p is selected from the group consisting of hydrogen and C_1 - C_4 alkyl, with hydrogen being more preferred;
- 5 c) R_q is selected from the group consisting of hydrogen, C_1 - C_4 alkyl, C_1 - C_4 alkoxy, and halogen, with hydrogen being more preferred;
- d) $R_{q'}$ is selected from the group consisting of hydrogen, C_1 - C_4 alkyl, C_1 - C_4 alkoxy, trifluoromethoxy, 2,2,2-trifluoroethoxy, trifluoromethyl halogen, and $-S(O)_pR_i$ wherein p is 2 and R_i is phenyl optionally substituted with C_1 - C_4 alkyl, C_1 - C_4 alkoxy, trifluoromethyl, with halogen being more preferred;
- 10 e) $R_{q''}$ is selected from the group consisting of hydrogen, C_1 - C_4 alkyl, C_1 - C_4 alkoxy, halogen, nitro, cyano, trifluoromethyl, and $-S(O)_pR_i$, wherein p 2 and R_i is phenyl optionally substituted with C_1 - C_4 alkyl, with halogen being more preferred; and
- 15 f) $R_{q'''}$ is selected from the group consisting of hydrogen, C_1 - C_4 alkyl, C_1 - C_4 alkoxy, halogen, trifluoromethyl, cyano, and nitro, with hydrogen and halogen being preferred.

Compounds in which R_4 is selected from the group consisting of C_2 - C_4 alkyl, fluorinated C_2 - C_4 alkyl and optionally substituted phenyl are preferred.

- When R_4 is C_2 - C_4 alkyl, particularly preferred groups include propyl, isopropyl, and butyl.
- 20

When R_4 is fluorinated C_2 - C_4 alkyl, preferred groups include 2,2-difluoroethyl, 2,2,2-trifluoroethyl, 3-fluoropropyl, 3,3-difluoropropyl, 3,3,3-trifluoropropyl, 2,2,3,3,3-pentafluoropropyl, and 2,2,3,3-tetrafluoropropyl.

- When R_4 is optionally substituted phenyl preferred groups include 1 to 3 groups independently selected from the group consisting of hydrogen, C_1 - C_4 alkyl, C_1 - C_4 alkoxy, halogen, cyano, and trifluoromethyl.
- 25

Preferred embodiments of the present invention are given for the compounds of formula II below:


- Compounds in which R_7 is optionally substituted phenyl or optionally substituted 5 to 6 membered monocyclic aromatic heterocycle having one heteroatom selected from the group consisting of nitrogen, oxygen, and sulfur and which 5 to 6 membered monocyclic aromatic heterocycle is optionally benzofused are preferred.
- 30

-11-

When R₇ is optionally substituted phenyl preferred substituents are 1 to 3 groups independently selected from the group consisting of hydrogen, C₁-C₄ alkyl, C₁-C₄ alkoxy, halogen, cyano, trifluoromethoxy, and trifluoromethyl.

5 Compounds in which R₇ is optionally substituted 5 to 6 membered monocyclic aromatic heterocycle having one heteroatom selected from the group consisting of nitrogen, oxygen, and sulfur and which 5 to 6 membered monocyclic aromatic heterocycle is optionally benzofused, the compounds which are benzo fused are preferred, with indolyl being preferred, and indol-3-yl being even more preferred, with the indol-3-yl depicted above for formula I being more preferred.


10 Preferred compounds of formula II having the points of attachment depicted below:



While only compounds of formula I are depicted, the compounds of formula I and
15 II are prepared as described in Schemes A and B below. In the Schemes below all substituents, unless otherwise indicated, are as previously defined, and all starting materials and reagents are well known and appreciated in the art and readily available or prepared by methods described herein. In the Schemes below, it is understood that
20 protecting groups can be used, where appropriate to allow for elaboration of a portion of the compounds of formula I or II. The selection, use, and removal of suitable protecting groups is well known and appreciated in the art (Protecting Groups in Organic Synthesis, Theodora Greene (Wiley-Interscience)).

-12-

Scheme A



Scheme A depicts alternative methods for the preparation of compounds of formula I by reductive amination.

In one alternative of Scheme A, step a, an appropriate compound of formula (1) is contacted with an appropriate compound of formula (2) in a reductive amination reaction to give a compound of formula I. An appropriate compound of formula (1) is one in which R_1 and R_2 are as desired in the final product of formula I or give rise to groups desired in the final product of formula I. An appropriate compound of formula (2) is one in which X , R_3 , and R_4 are as desired in the final product of formula I, or give rise to groups desired in the final product of formula I.

In another alternative of Scheme A, step a, an appropriate compound of formula (3) is contacted with an appropriate compound of formula (4) in a reductive amination reaction to give a compound of formula I. An appropriate compound of formula (3) is one in which R_1 and R_2 are as desired in the final product of formula I or give rise to groups desired in the final product of formula I. An appropriate compound of formula (4) is one in which X , R_3 , and R_4 are as desired in the final product of formula I, or give rise to groups desired in the final product of formula I.

The reductive amination depicted in Scheme A, step a, can be carried out under a variety of conditions, such as by hydrogenation using a suitable catalyst or using a suitable reducing agent.


For example, an appropriate amine of formula (1) is contacted with an appropriate aldehyde of formula (2) (or alternately an appropriate amine of formula (4) and an appropriate aldehyde of formula (3)) and a suitable reducing agent to give a compound of formula I. The reaction is carried out in a suitable solvent, such as methanol, ethanol, tetrahydrofuran, or mixtures of methanol or ethanol and tetrahydrofuran, dichloromethane, and 1,2-dichloroethane. The reaction may be carried out in the presence of a drying agent, such as sodium sulfate, cupric sulfate, or molecular sieves. The reaction

-13-

is carried out in the presence of from about 1 to 20 molar equivalents of a suitable reducing agent, such as, sodium borohydride, sodium cyanoborohydride, and sodium triacetoxyborohydride. It may be advantageous to allow Schiff base formation to proceed before addition of the suitable reducing agent. When sodium cyanoborohydride is used it may be advantageous to monitor and adjust the pH during the course of the reaction as is known in the art. The reaction is generally carried out at temperatures of from 0°C to the refluxing temperature of the solvent. Generally, the reactions require 1 to 72 hours. The product can be isolated and purified by techniques well known in the art, such as filtration, extraction, evaporation, trituration, chromatography, and recrystallization.

Scheme A, optional step b, not shown, an acid addition salt of a compound of formula I is formed using a pharmaceutically-acceptable acid. The formation of acid addition salts is well known and appreciated in the art.

Scheme B



Scheme B depicts alternative methods for the preparation of compounds of formula I by formation and reduction of an amide.

In one alternative, Scheme B, step a, depicts contacting an appropriate compound of formula (1) with an appropriate compound of formula (5) in an amide forming reaction to give a compound of formula (6). An appropriate compound of formula (1) is as described in Scheme A. An appropriate compound of formula (5) is one in which A is an activating group, taking the form of an acid halide, activated ester, activated amide, or

-14-

anhydride, and X, R₃, and R₄ are as desired in the final product of formula I, or give rise to groups desired in the final product of formula I.

In another alternative, Scheme B, step a, depicts contacting an appropriate compound of formula (7) with an appropriate compound of formula (4) in a amide forming reaction to give a compound of formula (8). An appropriate compound of formula (7) is one in which A is an activating group as described above and R₁ is as desired in the final product of formula I. An appropriate compound of formula (4) is as described in Scheme A. Appropriate compounds of formula (4) and (7) are generally available from commercial sources and can also be prepared by methods described herein and by methods described in the art.

The amide formation reaction depicted in Scheme B, step a, is readily accomplished by a number of methods readily available to the skilled person, including those which are conventionally conducted for peptide synthesis. Such methods can be carried out on the acid, acid halide, activated esters, activated amides, and anhydrides.

For example, well known coupling reagents such as carbodiimides with or without the use of well known additives such as N-hydroxysuccinimide, 1-hydroxybenzotriazole, etc. can be used to facilitate amide formation. Such coupling reactions are typically use about 1 to 1.5 molar ratios of acid, amine, and coupling reagent and are conventionally conducted in an inert aprotic solvent such as pyridine, dimethylformamide, dichloromethane, chloroform, 1,2-dichloroethane, acetonitrile, tetrahydrofuran and the like. It may be advantageous to use a suitable base, such as triethylamine or N,N-diisopropylethylamine, in such coupling reactions. The reaction is preferably conducted at from about 0°C to about 60°C until reaction completion which typically occurs within 1 to about 48 hours. Upon reaction completion, the product can be isolated and purified by techniques well known in the art, such as filtration, extraction, evaporation, trituration, chromatography, and recrystallization.

Alternatively, for example, an acid halide can be employed in the reaction. It may be advantageous to use a suitable base to scavenge the acid generated during the reaction, suitable bases include, by way of example, triethylamine, N,N-diisopropylethylamine, N-methylmorpholine, pyridine, and the like. Typically, about 1 to 1.5 molar ratios of the acid halide and amine are used. The reaction can be carried out in a variety of inert aprotic solvents such as pyridine, dichloromethane, chloroform, 1,2-dichloroethane, tetrahydrofuran, and the like. The reaction is preferably conducted at from about 0°C to about 60°C until reaction completion which typically occurs within 1 to about 12 hours. Upon reaction completion, the product can be isolated and purified by techniques well known in the art, such as filtration, extraction, evaporation, trituration, chromatography, and recrystallization.

-15-

Alternatively, for example, an acid halide can be employed in the reaction under Schotten-Baumann conditions. Typically, under such conditions 1 to 10 molar equivalents of amine are used. Such couplings generally use a suitable base to scavenge the acid generated during the reaction, such as sodium hydroxide, potassium hydroxide, sodium carbonate, potassium carbonate, sodium bicarbonate, potassium bicarbonate, and the like. The reaction can be carried out in a variety of mixed solvent systems such as dichloromethane, chloroform, ethyl acetate, tetrahydrofuran and the like; and water. The reaction is preferably conducted at from about 0°C to about 80°C until reaction completion which typically occurs within 1 to about 6 hours. Upon reaction completion, the product can be isolated and purified by techniques well known in the art, such as filtration, extraction, evaporation, trituration, chromatography, and recrystallization.

Alternatively, for example, an anhydride (either symmetrical or mixed) can be employed in the reaction. Such anhydrides are formed by numerous methods known in art. Typically, about 1 to 1.5 molar equivalents of the anhydride and amine are used. It may be advantageous to use a suitable base to scavenge the acid generated during the reaction. Suitable bases include, by way of example, triethylamine, N,N-diisopropylethylamine, N-methylmorpholine, pyridine, sodium carbonate, potassium carbonate, sodium bicarbonate, and the like. The reaction can be carried out in a variety of solvents. The reaction is preferably conducted at from about 0°C to about 60°C until reaction completion which typically occurs within 1 to about 12 hours. Upon completion, the product can be isolated and purified by techniques well known in the art, such as filtration, extraction, evaporation, trituration, chromatography, and recrystallization.

Scheme B, steps b, depicts reduction of a compound of formula (6) or (8) to give a compound of formula I.


For example, a compound of formula (6) or (8) is contacted with a suitable reducing agent to give a compound of formula I. Suitable reducing agents are those which are capable of reducing an amide to an amine and include, borane reagents, such as borane dimethyl sulfide complex, hydride transfer reagents, such as aluminum hydride and lithium aluminum hydride, and the like. The reaction is carried out in a solvent, such as tetrahydrofuran or diethyl ether, typically using 1 to 10 equivalents of reducing agent. The reaction is generally conducted at from about 0°C the refluxing temperature of the selected solvent and typically occurs within 1 to about 48 hours. The product can be isolated and purified by techniques well known in the art, such as quenching, filtration, extraction, evaporation, trituration, chromatography, and recrystallization.

Scheme B, as an optional step, not shown, an acid addition salt of a compound of formula I is formed using a pharmaceutically-acceptable acid. The formation of acid addition salts is well known and appreciated in the art.

-16-

In Schemes A and B, as an optional step, not shown, as will be appreciated by the skilled person, a compound of formula I in which R_2 is hydrogen can be alkylated to give a compound in which R_2 is not hydrogen. Methods for alkylating such secondary amines are well known in the art and discussed in Scheme C, step c, below.

- 5 In Schemes A and B, as will be appreciated by the skilled person, compounds of formula II are also prepared by the methods described in Schemes A and B using compounds of the formula (9) and (10), below:




- 10 An appropriate compound of formula (9) is one in which Y, R_5 and R_6 are as desired in the final product of formula II and an appropriate compound of formula (10) is one in which A is an activating group, as described above, and Y, R_5 and R_6 are as desired in the final product of formula II.

- Starting material for Schemes A and B are prepared in the Schemes below. In the Schemes below all substituents, unless otherwise indicated, are as previously defined, and
15 all starting material and reagents are well known and appreciated in the art.

Scheme C describes methods for preparing compounds of formula (1).

Scheme C



- Scheme C, step a, depicts the reaction of an appropriate aldehyde of formula (24) and nitromethane to give the compound of formula (25). An appropriate aldehyde of
20 formula (24) is one in which R_1 is as desired in the final product of formula I. The reaction of the anion of nitromethane with aldehydes to give nitro olefins is well known and appreciated in the art. Modern Synthetic Reactions, H.O. House (2nd ed. The Benjamin/Cummings Publishing Company 1972).

- For example, an appropriate aldehyde of formula (24) is condensed with
25 nitromethane to give the compound of formula (25). Typically the reaction is carried out

-17-

in the presence of an excess of nitromethane. The reaction is performed in a suitable solvent, such as tetrahydrofuran, nitromethane, and dimethyl sulfoxide. The reaction is performed using from about 1.1 to about 3 molar equivalents of a suitable base, such as sodium bis(trimethylsilyl)amide, potassium t-butoxide, sodium hydride, sodium acetate, triethylamine, N,N-diisopropylethylamine, ammonium salts, such as ammonium acetate. The reaction is carried out at temperatures of from about -20°C to the reflux temperature of the selected solvent and generally require from 6 hours to 48 hours. The product of the coupling reaction can be isolated and purified using techniques well known in the art, including extraction, evaporation, chromatography and recrystallization.

10 Scheme C, step b, depicts the reduction of a compound of formula (25) to give a compound of formula (1) in which R₂ is hydrogen.

For example, an appropriate compound of formula (25) is hydrogenated over a suitable catalyst, such as Raney® nickel or a palladium catalyst. When Raney nickel is used, the reaction is carried out in a suitable solvent, such as ethanol, methanol, water, and mixtures thereof. It may be advantageous to carry out the hydrogenation under acidic conditions, for example, using hydrochloric or sulfuric acid. When a palladium catalyst is used palladium-on-carbon is preferred and the reaction is carried out in a suitable solvent, such as ethanol, methanol, tetrahydrofuran, water, and mixtures thereof. It may be advantageous to carry out the hydrogenation under acidic conditions, for example, using hydrochloric, trifluoroacetic acid, or sulfuric acid. The reaction is generally carried out at temperatures of from ambient temperature to 70°C. The reaction is carried out with hydrogen at pressures of from 15 psi to 120 psi in an apparatus designed for carrying out reactions under pressure, such as a Parr® hydrogenation apparatus. The product can be isolated by carefully removing the catalyst by filtration and evaporation. The product can be purified by extraction, evaporation, trituration, chromatography, and recrystallization.

Alternately, for example, an appropriate compound of formula (25) is contacted with a suitable reducing agent. Suitable reducing agents include hydride transfer reagents, such as aluminum hydride and lithium aluminum hydride, and the like. The reaction is carried out in a solvent, such as tetrahydrofuran or diethyl ether, typically using 1 to 10 equivalents of reducing agent. The reaction is generally conducted at from about 0°C the refluxing temperature of the selected solvent and typically occurs within 1 to about 48 hours. The product can be isolated and purified by techniques well known in the art, such as quenching, filtration, extraction, evaporation, trituration, chromatography, and recrystallization.

35 Additionally, an appropriate compound of formula (25) can be reduced in two steps to a compound of formula (1). For example, the vinyl group of a compound of formula (25) can be reduced using reagents such as sodium borohydride. The reaction is

typically carried out using an excess of borohydride in a solvent, such as methanol, ethanol, isopropanol, water, and the like. The intermediate 2-nitroethyl compound can be isolated and purified by techniques well known in the art, such as quenching, filtration, extraction, evaporation, trituration, chromatography, and recrystallization. The intermediate 2-nitroethyl compound can then be reduced using a variety of methods, such as the hydrogenation and hydride transfer reagents as discussed above. Also, the intermediate 2-nitroethyl compound can be reduced using metals such as zinc to give the desired amine of formula (1) in which R_2 is hydrogen.

5
10 Scheme C, step c, depicts the optional alkylation of a compound of formula (1) in which R_2 is hydrogen to give a compound of formula (1) in which R_2 is not hydrogen.

For example, a compound of formula (1) in which R_2 is hydrogen is contacted with a suitable alkylating agent. A suitable alkylating agent is one which transfers a group R_2 as is desired in the final product of formula I. Suitable alkylating agents include C_1 - C_3 alkyl halides. The reaction is carried out in a suitable solvent, such as dioxane, tetrahydrofuran, tetrahydrofuran/water mixtures, or acetonitrile. The reaction is carried out in the presence of from 1.0 to 6.0 molar equivalents of a suitable base, such as sodium carbonate, sodium bicarbonate, potassium carbonate, potassium bicarbonate, triethylamine, or N,N-diisopropylethylamine. The reaction is generally carried out at temperatures of from -78°C . to the refluxing temperature of the solvent. Generally, the reactions require 1 to 72 hours. The product can be isolated and purified by techniques well known in the art, such as extraction, evaporation, trituration, chromatography, and recrystallization.

Alternately, for example, a compound of formula (1) in which R_2 is hydrogen undergoes a reductive amination with an aldehyde or ketone which gives a compound of formula (1) in which R_2 is not hydrogen. Suitable aldehydes include formaldehyde, acetaldehyde, propionaldehyde, and acetone. The reaction is carried out as described in Scheme A, step a.

25
30 In another alternate, for example, a compound of formula (1) in which R_2 is hydrogen undergoes amide or carbamate formation followed by reduction to give a compound of formula (1) in which R_2 is not hydrogen. Suitable aldehydes include formaldehyde, acetaldehyde, propionaldehyde, and acetone. The reaction is carried out as described in Scheme A, step a.

Scheme C, steps d and e, depict an alternative approach to preparing the compounds of formula (1) via formation of an amide using an appropriate compound of formula (7) and an appropriate amine of formula (26) to give an amide of formula (27), followed by reduction to give a compound of formula (1). An appropriate compound of formula (7) is as described in Scheme B. An appropriate amine of formula (26) is one


-19-

which gives R_2 as desired in final compound of formula I. The skilled person will recognize that many of the amides of formula (27) are commercially available and available in the art.

5 The amide formation and reduction in Scheme C are carried out as described in the Scheme B.

Scheme D describes methods for preparing compounds of formula (1) in which R_1 is optionally substituted indol-3-yl.

Scheme D



10 Scheme D, step a, depicts the two-step reaction of an appropriate indole of formula (28) with oxalyl chloride followed by an appropriate amine of formula (26), R_2NH_2 to give a compound of formula (29). An appropriate indole of formula (28) is one in which Z' represents optional substituents on the indole 2- and 4- to 7-positions as desired in the final product of formula I. An appropriate amine of formula (26) is as described in Scheme C, above.

15 For example, an appropriate indole of formula (28) is contacted with about 1 to 2 molar equivalents of oxalyl chloride to give an intermediate keto-acid chloride. The reaction is carried out in a suitable solvent, such as diethyl ether or tetrahydrofuran. The reaction is generally carried out at temperatures of from 0°C to 40°C and generally require from 6 hours to 48 hours. The intermediate keto-acid chloride product can be isolated and
 20 purified using techniques well known in the art, including extraction, evaporation, chromatography and recrystallization. Generally, the intermediate keto-acid chloride product is used directly after isolation. The intermediate keto-acid chloride product is contacted with an appropriate amine, R_2NH_2 , as defined above and using the procedures described above.

-20-

Scheme D, step b, depicts the reduction of a compound of formula (29) to give a compound of formula (1) in which R₁ is optionally substituted indol-3-yl.

For example, a compound of formula (29) is reduced using a suitable reducing reagent such as, lithium aluminum hydride to give a compound of formula (1) which R₁ is optionally substituted indol-3-yl. The reaction is carried out in a solvent, such as tetrahydrofuran or diethyl ether, typically using 1 to 12 molar equivalents of reducing agent. The reaction is generally conducted at from about 0°C the refluxing temperature of the selected solvent and typically occurs within 12 to about 48 hours. The product can be isolated and purified by techniques well known in the art, such as quenching, filtration, extraction, evaporation, trituration, chromatography, and recrystallization.

In Scheme D, step c, an appropriate indole of formula (28) is formylated to give a compound of formula (30). An appropriate indole of formula (28) is as described in step a, above.

For example, an appropriate indole of formula (28) is reacted with a suitable formyl transfer reagent, such as the Vilsmeier reagent formed from dimethylformamide. Generally, about 1 molar equivalent of formyl transfer reagent is used. The reaction is performed in a suitable solvent, such as benzene, dimethylformamide, tetrahydrofuran, or diethyl ether. The reaction is carried out at temperature of from about -70°C to about 20°C and generally require from 1 hours to 6 hours. The product of the reaction can be isolated and purified using techniques well known in the art. These techniques include extraction, evaporation, chromatography and recrystallization.

In Scheme D, step d, an appropriate indole of formula (28) is contacted with (CH₃)₂N-CH=CH-NO₂ to give a compound of formula (30). An appropriate indole of formula (28) is as described in step a, above.


For example, an appropriate indole of formula (28) is reacted with 1-dimethylamino-2-nitroethylene. Generally, about 1 equimolar amounts of reagents. The reaction is performed in a suitable solvent, such as trifluoroacetic acid or dichloromethane containing about 2 to 15 equivalents of trifluoroacetic acid. The reaction is carried out at temperature of from about -70°C to about 20°C and generally require from 1 hours to 24 hours. The product of the reaction can be isolated and purified using techniques well known in the art. These techniques include extraction, evaporation, chromatography and recrystallization.

Scheme D, steps e and f, depict an the reaction of an aldehyde of formula (30) to give a nitro olefin of formula (31) and the reduction of the nitro olefin to give a compound of formula (1) in which R₁ is optionally substituted indol-3-yl. These steps can be carried out using the methods described in Scheme C.

As will be appreciated by the skilled person, in steps not shown, the indole nitrogen of a compound of formula (1) can be substituted, as desired, using suitable amine protecting groups to give compounds in which R_1 is 1-substituted indol-3-yl. Also as will be appreciated by the skilled person, in steps described in Scheme C, R_2 groups which are not hydrogen can be introduced by various methods.

Scheme E describes methods for preparing compounds of formula (2) in which X is -O- or -S-.

Scheme E



Scheme E, step a, depicts the formation of an acetal of an appropriate compound of formula (11) to give a compound of formula (12). An appropriate compound of formula (11) is one in which X and R_3 are as desired in the final compound of formula I. Such acetal formation reactions are readily accomplished by methods well known in the art. (Protecting Groups in Organic Synthesis, Theodora Greene (Wiley-Interscience)).

For example, a compound formula (11) is contacted under acid catalysis with an appropriate alcohol, HOR. An appropriate alcohol is one which gives an acetal with is stable to the reaction in step b and can be removed in step c to give a compound of formula (2). Appropriate alcohols include methanol, ethanol, propanol, 1,3-propane diol, ethylene glycol, and the like.

In Scheme E, step b, an appropriate compound of formula (11), (12), or (14) is reacted with an R_4 group transfer reagent, as desired, to give a compound of formula (2), (13), or (15); respectively. Appropriate compounds of formula (11), (12), and (14) are

-22-

ones in which X and R₃ are as desired in the final product of formula I. A variety of reagents that transfers an R₄ as desired in the final product are available and suitable for the reaction depicted in Scheme E. Such reagents include halopyridines, halopyridine N-oxides, allyl halides, C₂-C₄ alkanols, C₂-C₄ alkyl halides and sulfonates, fluorinated C₂-C₄ alkanols, fluorinated C₂-C₄ alkyl halides and sulfonates, optionally substituted phenyl
5 having at least one fluoro or chloro atom, optionally substituted phenylsulfonyl halides or anhydrides, and optionally substituted benzyl halides.

For example, where the appropriate R₄ group transfer reagent is a halide, sulfonate, or anhydride, an appropriate compound of formula (11), (12), or (14) is coupled
10 under basic conditions to give a compound of formula (2), (13), or (15); respectively. The reaction is performed in a suitable solvent, such as acetonitrile, dimethylformamide, dimethylacetamide, tetrahydrofuran, pyridine, and dimethyl sulfoxide. The reaction is carried out in the presence of from about 1 to about 3 molar equivalents of a suitable base, such as potassium hydride, sodium hydroxide, sodium hydride, sodium carbonate,
15 potassium carbonate, cesium carbonate, N,N-diisopropylethylamine, triethylamine, and the like. The reaction is carried out at temperature of from about -30°C to about 100°C and generally require from 6 hours to 48 hours. The product of the reaction can be isolated and purified using techniques well known in the art. These techniques include extraction, evaporation, chromatography and recrystallization.

20 Of course, when a halopyridine N-oxide is used the N-oxide is removed by reduction to give the R₄ as desired in the final product of formula I. Such reductions are readily accomplished by the skilled person, and include catalytic reduction over palladium catalysts using hydrogen or ammonium formate in a suitable solvent such as methanol, ethanol, water, and mixtures thereof.

25 Alternately, for example, where the appropriate R₄ group transfer reagent is an alkanol, the coupling can be carried out under Mitsunobu conditions which are well known in the art. The reaction is carried out in a suitable solvent, such as tetrahydrofuran and diethyl ether using a phosphine, such as triphenylphosphine or a resin bound phosphine and a dialkyl azodicarboxylate, such as diethyl azodicarboxylate, diisopropyl
30 azodicarboxylate or di-*t*-butyl azodicarboxylate. The reaction is generally carried out at temperatures of from ambient temperatures to 60°C. The reaction generally requires from 1 hour to 12 hours. The product can be isolated by techniques well known in the art, such as extraction and evaporation. The product can then be purified by techniques well known in the art, such as distillation, chromatography, or recrystallization.

35 Scheme E, step c, depicts the deprotection of an acetal of formula (13) to give a compound of formula (2). Such deprotections are readily accomplished by methods well

-23-

known in the art. (Protecting Groups in Organic Synthesis, Theodora Greene (Wiley-Interscience)).

For example, a compound formula (13) is contacted under acid under aqueous conditions to give a compound of formula (2).


5 In Scheme E, step d, a bromo compound of formula (15) is formylated to give a compound of formula (2).

For example, a compound of formula (15) is metalated by treatment with a metalation reagent such as butyl lithium. The reaction is performed in a suitable solvent, such as hexane, benzene, toluene, tetrahydrofuran or diethyl ether. The reaction is
10 typically carried out in the presence of from about 1 to about 1.5 molar equivalents of a metalating reagent. The metalation reaction is carried out at temperature of from about -70°C to about 20°C and generally require from 1 hours to 6 hours. The metalated species is then treated with a formyl transfer reagent, such as dimethylformamide or an alkyl chloroformate to give a compound of formula (2) or a alkoxycarbonyl compound which
15 can be elaborated to an aldehyde as described herein. The product of the reaction can be isolated and purified using techniques well known in the art. These techniques include extraction, evaporation, chromatography and recrystallization.

Scheme F describes methods for preparing compounds of formula (2) from the versatile intermediate, compound (17), which readily prepared by acetal formation as
20 described above.

-24-

Scheme F



Scheme F, step a, depicts an aromatic displacement reaction of an appropriate compound of formula (17) and an appropriate alcohol (R_4OH) or an appropriate thiol (R_4SH) to give a compound of formula (13) in which X is -O- or -S- are defined above in Scheme E. An appropriate compound of formula (17) is one in which R_3 is as desired in the final product of formula I. In an appropriate alcohol (R_4OH) or an appropriate thiol (R_4SH), R_4 is as desired in the final product of formula I, and includes $\text{C}_2\text{-C}_4$ alkyl alcohols and thiols, fluorinated $\text{C}_2\text{-C}_4$ alkyl alcohols and thiols, optionally substituted phenols and thiophenols, optionally substituted benzyl alcohols and thiols.

For example, an appropriate compound of formula (17) and an appropriate alcohol (R_4OH) or an appropriate thiol (R_4SH) are coupled give a compound of formula (13). The reaction is performed in a suitable solvent, such as dimethylformamide,

-25-

dimethylacetamide, and dimethyl sulfoxide. The reaction is performed using from about 1.1 to about 3 molar equivalents of an appropriate alcohol or thiol. The reaction is carried out in the presence of from about 1 to about 6 molar equivalents of a suitable base, such as potassium hydride, sodium hydroxide, potassium carbonate, sodium carbonate, or sodium hydride. The coupling is performed using a suitable catalyst, such as copper salts. The reaction generally requires from 6 hours to 48 hours. The product of the coupling reaction can be isolated and purified using techniques well known in the art. These techniques include extraction, evaporation, chromatography and recrystallization.

Scheme F, steps b-e, depict a number of reactions of an appropriate compound of formula (17), after metalation as described in Scheme E, step d, to give compounds of formula (18)-(21), respectively. In these steps an appropriate compound of formula (17) is one in which R_3 is as desired in the final product of formula I and is not adversely affected by the metalation conditions of the reaction. Generally, these reactions are performed in the solvent used and at the temperature used to form the metalated species. The products of these reactions can be isolated and purified using techniques well known in the art, include quenching, extraction, evaporation, trituration, chromatography, and recrystallization.

For example, in Scheme F, step b, a metalated compound of formula (17) is contacted with an appropriate disulfide $(R_4S)_2$, to give a compound of formula (18). An appropriate disulfide is one that gives R_4 as desired in the final product of formula I and gives rise to compounds in which X is -S-. Appropriate disulfides include C_1 - C_4 alkyl disulfides, optionally substituted phenyl disulfides, and optionally substituted benzyl disulfides. The reaction is performed using from about 1 to about 2 molar equivalents of an appropriate disulfide. The reaction is typically carried out in the same solvent used for the metallation and at temperatures of about -78°C to about 50°C . The reaction generally require from 12 hours to 48 hours.

For example, in Scheme F, step c, a metalated compound of formula (17) is contacted with an appropriate sulfonyl fluoride (R_4SO_2F) to give a compound of formula (19). An appropriate sulfonyl fluoride is one that transfers R_4 as desired in the final product of formula I and gives rise to compounds in which X is $-SO_2-$. Appropriate sulfonyl fluorides include an optionally substituted phenyl sulfonyl fluoride. The reaction is performed using from about 1 to about 3 molar equivalents of an appropriate sulfonyl fluoride. The reaction is typically carried out in the same solvent used for the metallation and at temperatures of about -78°C to about 0°C . The reaction generally require from 2 hours to 12 hours.

For example, in Scheme F, step d, a metalated compound of formula (17) is contacted with an appropriate acid chloride $(R_4C(O)Cl)$ to give a compound of formula

-26-

(20). An appropriate acid chloride is one that transfers R_4 as desired in the final product of formula I and gives rise to compounds in which X is $-C(O)-$. Appropriate acid chlorides include C_2-C_4 alkyl acid chlorides, fluorinated C_2-C_4 alkyl acid chlorides, optionally substituted phenyl acid chlorides, optionally substituted benzyl acid chlorides, and optionally substituted 5 to 6 membered monocyclic aromatic heterocycle acid chlorides. The reaction is performed using from about 0.8 to about 1.2 molar equivalents of an appropriate acid chloride. The reaction is typically carried out in the same solvent used for the metallation and at temperatures of about -78°C to about 50°C . The reaction generally require from 1 hours to 12 hours.

For example, in Scheme F, step e, a metalated compound of formula (17) is contacted with an appropriate aldehyde ($R_4C(O)H$) to give a compound of formula (21). An appropriate aldehyde is one that transfers R_4 as desired in the final product of formula I and gives rise to compounds in which X is $-CH(OH)-$. Appropriate aldehydes include C_2-C_4 alkyl aldehyde, fluorinated C_2-C_4 alkyl aldehyde, optionally substituted phenyl aldehyde, optionally substituted benzyl aldehyde, and optionally substituted 5 to 6 membered monocyclic aromatic heterocycle aldehyde. The reaction is performed using from about 1 to about 3 molar equivalents of an appropriate aldehyde. The reaction is typically carried out in the same solvent used for the metallation and at temperatures of about -50°C to about 50°C . The reaction generally requires from 4 hours to 24 hours.


As will be appreciate by the skilled person, compounds of formula (18)-(21) can undergo a number of other transformations which are depicted in Scheme F, steps f-i, to give, ultimately, compounds of formula I having various groups at X. These transformations are trivial and well within the ability of the skilled person. These transformations include oxidation of sulfides (step f) which can be accomplished by peroxide, peracids, and other reagents known in the art; reduction of a benzyl alcohol (step g) which can be accomplished by a variety of reagents, such as triethylsilane/trifluoroacetic acid; halogenation of a benzyl alcohol to give fluoro (step h) using reagents such as DAST and fluorinating reagents; reduction of a ketone (step i) using various hydride transfer reagents or oxidation of a benzylic alcohol (step i) which can be accomplished by manganese dioxide or Swern conditions.

In Scheme F, step j, compounds of the formula (13) and (18)-(23) are deprotected to give an aldehyde of formula (2) as described in Scheme E, step c.

Scheme G describes methods for preparing compounds of formula (5).

-27-

Scheme G



Scheme G, step a, a bromo compound of formula (15) is carboxylated to give a compound of formula (5) in which A is -OH.

For example, a compound of formula (15) is metalated as described in Scheme E, step d, and the metalated species is then treated with carbon dioxide to give a compound of formula (5) in which A is -OH. The product of the reaction can be isolated and purified using techniques well known in the art. These techniques include extraction, evaporation, chromatography and recrystallization.

Scheme G, step b, a bromo compound of formula (15) is alkoxyformylated using an appropriate chloroformate or carbonate to give a compound of formula (32). An appropriate chloroformate or carbonate is one that transfers an RO(O)C- group in which R is methyl, ethyl, or benzyl.

For example, a compound of formula (15) is metalated as described in Scheme E, step d, and the metalated species is then treated with about 1 to 3 molar equivalents of an appropriate chloroformate or carbonate. The reaction is typically carried out in the same solvent used for the metallation and at temperatures of about -78°C to about 50°C. The reaction typically requires from 1 to 24 hours. The product of the reaction can be isolated and purified using techniques well known in the art. These techniques include extraction, evaporation, chromatography and recrystallization.

In Scheme G, step c, an appropriate compound of formula (33) is reacted with an R₄ group transfer reagent, as desired, to give a compound of formula (32). An appropriate compound of formula (33) is one in which X and R₃ are as desired in the final product of formula I. Reagents that transfers an R₄ are as described in Scheme E.

-28-

For example, where the appropriate R₄ group transfer reagent is a halide or anhydride, an appropriate compound of formula (34) is coupled under basic conditions with to give a compound of formula (33). The reaction is performed in a suitable solvent, such as dimethylformamide, tetrahydrofuran, or pyridine. The reaction is typically carried out in the presence of from about 1 to about 3 molar equivalents of a suitable base, such as sodium carbonate, potassium carbonate, cesium carbonate, N,N-diisopropylethylamine, triethylamine, and the like. The reaction is carried out at temperature of from about -30°C to about 100°C and generally require from 6 hours to 48 hours. The product of the reaction can be isolated and purified using techniques well known in the art. These techniques include extraction, evaporation, chromatography and recrystallization.

Alternately, for example, where the appropriate R₄ group transfer reagent is an alkanol, the coupling can be carried out under Mitsunobu conditions which are well known in the art and described in Scheme E.

Scheme G, step d, an ester of formula (32) is deprotected to give a compound of formula (5) in which A is -OH. Such deprotections are readily accomplished by methods well known in the art. (Protecting Groups in Organic Synthesis, Theodora Greene (Wiley-Interscience)).


Scheme G, step e, a compound of formula (5) in which A is -OH is converted to a compound of formula (5) in which A is an activating group, such as acid halide, activated ester, activated amide, or anhydride. The formation of such activated intermediates is well known and appreciated in the art.

For example, an acid halide can be prepared by a variety of reagent such as oxalyl chloride, oxalyl bromide, thionyl chloride, thionyl bromide, phosphorous oxychloride, phosphorous trichloride, and phosphorous pentachloride; a mixed anhydride of substituted phosphoric acid, such as dialkylphosphoric acid, diphenylphosphoric acid, halophosphoric acid; of aliphatic carboxylic acid, such as formic acid, acetic acid, propionic acid, butyric acid, isobutyric acid, pivalic acid, 2-ethylbutyric acid; an activated ester, such as phenol ester, p-nitrophenol ester, N-hydroxysuccinimide ester, N-hydroxyphthalimide ester, 1-hydroxybenztriazole ester; or activated amide, such as imidazole, dimethylpyrazole, triazole; are prepared by method which are well known and appreciated in the art. Such activated intermediates may be prepared and used directly or are prepared and isolated before use in the schemes above.

Scheme H describes methods for preparing compounds of formula (4).

-29-

Scheme H



Scheme H, step a, a bromo compound of formula (15) is converted to a nitrile of formula (35).

For example, a compound of formula (15) is treated with copper (I) cyanide to give a compound of formula (35). The reaction is performed in a suitable solvent, such as dimethylformamide. The reaction is typically carried out in the presence of from about 1 to about 3 molar equivalents of copper (I) cyanide. The reaction is carried out at temperature of from about ambient temperature to about 100°C and generally require from 6 hours to 48 hours. The product of the reaction can be isolated and purified using techniques well known in the art. These techniques include extraction, evaporation, chromatography and recrystallization.

Scheme H, step b, a nitrile compound of formula (35) reduced to give a compound formula (4) in which R_2 is hydrogen.

For example, a nitrile compound of formula (35) is contacted with sodium borohydride in the presence of cobalt chloride. The reaction is carried out in a suitable solvent, such as methanol, or ethanol. The reaction is generally carried out at temperatures of from 0°C. to 50°C. Generally, the reactions require 1 to 72 hours. The product can be isolated and purified by techniques well known in the art, such as extraction with aqueous acid, evaporation, trituration, chromatography, and recrystallization.

-30-

Alternately, for example, a nitrile compound of formula (35) is hydrogenated over a suitable catalyst, such as Raney® nickel. The reaction is carried out in a suitable solvent, when Raney® nickel is used as the catalyst, suitable solvents will generally contain ammonia, such as ethanol/ammonium hydroxide. The reaction is generally carried out at temperatures of from ambient temperature to 50°C. The reaction is carried out at pressures of from 15 psi (103 kPa) to 120 psi (827 kPa) in an apparatus designed for carrying out reactions under pressure, such as a Parr hydrogenation apparatus. The product can be isolated by carefully removing the catalyst by filtration and evaporation. The product can be purified by extraction, evaporation, trituration, chromatography, and recrystallization.

Scheme H, step c, a nitrile compound of formula (35) is converted to an amide of formula (36).

For example, a compound of formula (35) is treated with acid or base under hydrolysis conditions to give a compound of formula (36). The reaction is performed in a suitable solvent, such as ethanol, isopropanol, dimethylsulfoxide, each containing water. The hydrolysis of an aromatic nitrile to an amide is well known and appreciated in the art. The product of the reaction can be isolated and purified using techniques well known in the art. These techniques include extraction, evaporation, chromatography and recrystallization.

Scheme H, step d, depicts formation of an amide of formula (37) by reacting a compound of formula (5) and an appropriate amine of formula H_2NR_2 in an amide forming reaction. An appropriate amine of formula H_2NR_2 is one which gives R_2 as desired in the final product of formula I. Suitable methods of forming amides are well known in the art and are described in Scheme B, above.

Scheme H, step e, a amide compound of formula (36) or (37) is reduced to a compound of formula (4). Such reductions of amides are readily carried out as described in Scheme B, above, and as known in the art.

Scheme H, step f, a compound of formula (2) and an appropriate amine of formula H_2NR_2 undergo reductive amination to give a compound of formula (4). Such reductive aminations are readily carried out as described in Scheme B, above, and by other methods known in the art.

As will be appreciated by the skilled person, the compounds of formula II are readily prepared by methods analogous to those described above.

The present invention is further illustrated by the following examples and preparations. These examples and preparations are illustrative only and are not intended to limit the invention in any way.

-31-

The terms used in the examples and preparations have their normal meanings unless otherwise designated. For example, "°C" refers to degrees Celsius; "N" refers to normal or normality; "M" refers to molar or molarity; "mmol" refers to millimole or millimoles; "g" refers to gram or grams; "mL" refers milliliter or milliliters; "mp" refers to melting point; "brine" refers to a saturated aqueous sodium chloride solution; etc. In the ¹H NMR, all chemical shifts are given in δ, unless otherwise indicated.

Example 1

2-(3-(4-Fluorophenoxy)phenyl)-(1,3)dioxolane

Combine 4-fluorophenol (3.0 g, 227.8 mmol), 2-(3-bromophenyl)-1,3-dioxolane (5.0 ml, 33.3 mmol), potassium carbonate (anhydrous, 8.0 g, 55.6 mmol), and dry pyridine (50 ml). Heat to 90°C and add copper oxide (5.5 g, 69.5 mmol). Heat at reflux. After 24 hours, cool to room temperature, dilute with dichloromethane, and filter. Concentrate the filtrate in vacuum to give a yellow oil. Chromatograph the oil on silica gel eluting with 95:5 hexane:EtOAc to give the title compound.

By the method of Example 1 the following compounds were prepared: a) 2-(3-(2-Fluorophenoxy)phenyl)-(1,3)dioxolane;

b) 2-(3-(3-Fluorophenoxy)phenyl)-(1,3)dioxolane;

c) 2-(3-(Naphth-2-yloxy)phenyl)-(1,3)dioxolane;

d) 2-(3-(Naphth-1-yloxy)phenyl)-(1,3)dioxolane;

e) 2-(3-(Pyrid-3-yloxy)phenyl)-(1,3)dioxolane;

f) 3-(Pyridin-3-yloxy)benzaldehyde;

g) 3-(Pyrimidin-5-yloxy)benzaldehyde; and

h) 3-(Pyridin-4-yloxy)benzaldehyde.

Example 2

3-[1,3]Dioxolan-2-ylphenyl)phenylamine

Combine 2-(3-bromophenyl)-1,3-dioxolane (0.7 ml, 4.3 mmol), aniline (0.4 ml, 4.7 mmol), sodium t-butoxide (0.6 g, 6.0 mmol), BINAP (10.0 mg, 0.03 mol), Pd₂(dba)₃ (30.0 mg, 0.01 mmol) and toluene (20 ml). Heat at 80 °C. After 18 hours, cool to room temperature, dilute with ether (40 ml), filter, and concentrate to give a residue. Chromatograph the residue on silica gel eluting with 9:1 hexane:EtOAc to give the title compound.

By the method of Example 2 the following compounds were prepared: a) Benzyl-(3-[1,3]dioxolan-2-ylphenyl)-amine;

b) (3-[1,3]Dioxolan-2-ylphenyl)-pyridin-3-ylamine;

c) (3-[1,3]Dioxolan-2-ylphenyl)-pyridin-4-ylamine; and

-32-

d) (3-[1,3]Dioxolan-2-ylphenyl)-pyridin-2-ylamine.

Example 3

2-(3-Phenylsulfanylphenyl)-[1,3]-dioxolane

5 Combine 2-(3-bromophenyl)-1,3-dioxolane (3.0 ml, 20.0 mmol) and tetrahydrofuran (100 ml). Cool to about -78°C . Add dropwise a solution of n-butyllithium, 1.6 M solution in hexane (13.4 ml, 21.0 mmol). After 10 minutes, add phenyl disulfide (4.3 g, 20.0 mmol) as a solution in tetrahydrofuran (50 ml). After 1 hour, warm to room temperature over 1 hour then quenched with water (150 ml) and extract
10 with ether. Extract the combined organic layers sequentially with distilled water and brine and then dry (Na_2SO_4), filter, and concentrate to give a residue. Chromatograph the residue eluting with 9:1 hexane:EtOAc gives the title compound.

By the method of Example 3 the following compounds were prepared: a) 2-(3-p-Tolylsulfanylphenyl)-[1,3]-dioxolane.

15

Example 4

2-(3-Benzenesulfonylphenyl)-[1,3]-dioxolane

Combine 2-(3-phenylsulfanylphenyl)-[1,3]-dioxolane (1.0 g, 3.6 mmol) and dichloromethane (15 ml). Cool to about -78°C . Add a slurry of m-chloroperbenzoic acid (2.3 g, 7.2 mmol) in dichloromethane (10 ml). After 30 minutes, warm to room
20 temperature then add a solution 1N of sodium thiosulfate (20 ml). After 15 minutes, add a solution of saturated sodium bicarbonate. Separate the organic layer and extract the aqueous layer with dichloromethane. combine the organic layers and wash sequentially with distilled water and brine and then dry (Na_2SO_4), filter, and concentrate to give a
25 residue. Chromatograph the residue eluting with EtOAc to give the title compound.

By the method of Example 4 the following compounds were prepared: a) 2-(3-(Toluene-4-sulfonyl)-phenyl)-[1,3]-dioxolane.

30

Example 5

(3-[1,3]Dioxolan-2-ylphenyl)phenylmethanol

Combine 2-(3-bromophenyl)-1,3-dioxolane (10.0 ml, 66.0 mmol) and tetrahydrofuran (100 ml) and cool to about -78°C . Add dropwise a solution of n-butyllithium, 1.6 M solution in hexane (44.0 ml, 66.0 mmol). After 10 min, add a solution
35 of benzaldehyde (7.6 ml, 66.0 mmol) in tetrahydrofuran (50 ml) via cannula. After 1 hour, warm to room temperature. After 18 hours, quench into water and extract with dichloromethane. Combine the organic layers and wash sequentially with distilled water

-33-

and brine and then dry (Na_2SO_4), filter, and concentrate to give a residue. Chromatograph the residue eluting with 7:3 hexane:EtOAc to give the title compound.

Example 6

5 (3-[1,3]Dioxolan-2-ylphenyl)phenylmethanone

Combine (3-[1,3]dioxolan-2-ylphenyl)-phenyl-methanol (5.0 g, 18.5 mmol) and 18-crown-6 (160 mg, 0.6 mmol) in dichloromethane. Add potassium permanganate (8.8 g, 55.5 mmol). Heat to about 40°C. After 4 hours, cool to room temperature add water and sodium hydrogensulfite (6.0 g). Basify with a solution of 1N sodium hydroxide (about 10 60 ml) and extract with dichloromethane. Combine the organic layers and wash sequentially with distilled water and brine and then dry (Na_2SO_4), filter, and concentrate to give a residue. Chromatograph the residue eluting with EtOAc to give the title compound.

15 Example 7

3-Benzylbenzaldehyde

Combine (3-[1,3]dioxolan-2-ylphenyl)-phenyl-methanol (2.3 g, 8.7 mmol) and sodium iodide (5.3 g, 35.0 mmol) in acetonitrile (25 ml). Add dichloromethylsilane (2.1 ml, 17.4 mmol) via syringe. After 10 min, dilute with EtOAc and wash with water, 20 saturated sodium bicarbonate, 10% sodium thiosulfate, and then brine. Dry the organic layers (Na_2SO_4), filter, and concentrate to give a residue. Chromatograph the residue eluting with 9:1 hexane:EtOAc to give the title compound.

Example 8

25 2-(3-(α -Fluorobenzyl)phenyl)-[1,3]-dioxolane

Combine (3-[1,3]-dioxolan-2-ylphenyl)phenylmethanol (2.3 g, 8.9 mmol) and dichloromethane (50 ml). Cool to about -78°C. Add dropwise a solution of (diethylamino)sulfur trifluoride (1.7 ml, 12.9 mmol). After 10 minutes, quench with water and extract with dichloromethane. Combine the organic layers and wash sequentially with 30 distilled water and brine and then dry (Na_2SO_4), filter, and concentrate to give a residue. Chromatograph the residue eluting with 7:3 hexane:EtOAc to give the title compound.

Example 9

3-Nitrodibenzofuran

35 Combine dibenzofurane (20.0 g, 0.11 mol) and acetic acid (80 ml). Heat to 65°C. Add 98% nitric acid (20.0 g, 11.8 mol). After 3 hours, cool to room temperature to give a solid. Collect the solid by filtration, rinse with water, and dry to give the title compound.

-34-

Example 10N-Dibenzofuran-3-ylamine

Combine 3-nitrodibenzofuran (22.0 g, 0.1 mol) and Raney nickel (2.75g), and
5 ethanol (365 ml) and hydrogenate at room temperature and 40 psi (276 kPa). After 18
hours, filter and concentrate the filtrate to residue. Chromatograph the residue eluting
with 9:1 hexane:EtOAc to give the title compound.

Example 1110 3-Bromodibenzofuran

Combine N-dibenzofuran-3-ylamine (2.0 g, 10.8 mmol), water (20 ml), and conc.
hydrobromic acid (6 ml). Cool to 0 °C. Add a solution of sodium nitrite (0.7 g, 10.8
mmol) in water (16 ml). After 15 minutes add the mixture above to a mixture of copper
bromide (1.7 g, 12.3 mmol) in water (9.2 ml) and hydrobromic acid (4 ml). Warm to
15 ambient temperature. After 18 hours, add water and extract with dichloromethane.
Combine the organic layers and wash sequentially with distilled water and brine and then
dry (Na₂SO₄), filter, and concentrate to give a residue. Chromatograph the residue eluting
with 8:2 hexane:EtOAc to give the title compound.

20 Example 12Dibenzofuran-3-carbaldehyde

Combine 3-bromodibenzofuran (0.5 g, 2.0 mmol) and tetrahydrofuran (30 ml).
Cool to about -78°C. Add a solution of *t*-butyllithium, 1.6 M solution in hexane (2.2 ml,
3.0 mmol), then warm to about 0°C for 10 min. Cool to about -78°C and add
25 dimethylformamide (0.5 ml, 5.9 mmol). Warm to room temperature, quench with water,
and extract with dichloromethane. Combine the organic layers and wash sequentially with
distilled water and brine and then dry (Na₂SO₄), filter, and concentrate to give a residue.
Chromatograph the residue eluting with 8:2 hexane:EtOAc to give the title compound.

30 Example 133-(Thiazol-2-yloxy)benzaldehyde

In a sealed tube, combine 2-bromo-thiazole (2.0 ml, 22.2 mmol), 3-hydroxy-
benzaldehyde (1.8 g, 15.0 mmol) and potassium carbonate (2.1 g, 15.0 mmol) in
dimethylformamide (20 ml). Heat to 100 °C. After 48 hours, cool, pour into water (150
35 ml), and extract with ether. Combine the organic layers and wash sequentially with
distilled water and brine and then dry (Na₂SO₄), filter, and concentrate to give a residue.
Chromatograph the residue eluting with 9:1 hexane:EtOAc to give the title compound.

Example 146-Bromo-1H-indole

Combine 4-bromo-2-nitrotoluene (5.0 g, 23.1 mmol), dimethylformamide (50 ml),
5 DMF-dimethylacetal (9.0 ml, 69.4 mmol), and pyrrolidine (2.0 ml, 23.1 mmol). Heat to
110 °C. After 3 hours, cool to room temperature, dilute with ether, and wash with water.
Combine the organic layers, and concentrate to give a residue. Combine the residue and
80% aq. acetic acid (120 ml) and heat at 75°C. Add zinc dust (13.1 g, 200.5 mmol)
portionwise. Heat to 85°C. After 2 hours, cool and filter. Dilute the filtrate with ether,
10 wash with water dry (Na₂SO₄), and concentrate to give a residue. Chromatograph the
residue eluting with 9:1 hexane:EtOAc to give the title compound.

By the method of Example 14 the following compounds were prepared: 4-Bromo-
1H-indole.

Example 151H-Indole-6-carbaldehyde

Combine hexane washed potassium hydride (1.3 g, 10.7 mmol) and ether (20 ml).
Cool to about 0 °C and add a solution of 6-bromo-1H-indole (2.1 g, 10.7 mmol) in ether
(5 ml). After 15 min, cool to about -78°C and add a solution of t-butyllithium, 1.4 M
20 solution in hexane (14.0 ml, 10.7 mmol). After 10 min, add dimethylformamide (1.7 ml,
20.0 mmol) in ether (5ml). Slowly warm to room temperature and then pour into a ice
cold solution of 1M phosphoric acid and extract with EtOAc. Combine the organic layers
and wash sequentially with distilled water and brine and then dry (Na₂SO₄), filter, and
concentrated to give a residue. Chromatograph the residue eluting with 9:1 hexane:EtOAc
25 to give the title compound.

By the method of Example 15 the following compounds were prepared: 1H-
Indole-4-carbaldehyde.

Example 161-Phenyl-1H-indole-6-carbaldehyde

Combine in a sealed tube 1H-indole-6-carbaldehyde 0.9 g, 6.2 mmol), copper(I)
trifluoromethanesulfate-complex (0.2 g, 0.3 mmol), phenanthroline (1.3 g, 6.2 mmol),
transdibenzylidenacetone (0.1 g, 0.3 mmol), cesium carbonate (2.6 g, 7.9 mmol) and
iodobenzene (1.6 ml, 14.3 mmol) in xylene (40 ml). Heat at about 110°C. After 24
35 hours, cool to room temperature, dilute with dichloromethane and saturated ammonium
chloride. Separate the layer and Extract the aqueous layer with dichloromethane.
Combine the organic layers and wash sequentially with distilled water and brine and then

-36-

dry (Na_2SO_4), filter, and concentrated to give a residue. Chromatograph the residue eluting with 8:2 hexane:EtOAc to give the title compound.

By the method of Example 16 the following compounds were prepared: 1-Phenyl-1H-indole-4-carbaldehyde.

5

Example 17

3-Phenylsulfanylbenzaldehyde

Combine 2-(3-phenylsulfanylphenyl)-[1,3]-dioxolane (0.3 g, 1.1 mmol) and acetonitrile (8.0 ml) add a solution of hydrochloric acid (1N, 2.0 ml). After 18 hours, concentrate in vacuum to remove most of the acetonitrile, dilute with water and extract with ether. Combine the organic extracts and wash once with saturated sodium bicarbonate, then with brine. Dry (Na_2SO_4) the organics, filter, and concentrated to give the title compound.

By the method of Example 17 the following compounds were prepared: a) 3-Benzenesulfonylbenzaldehyde;

- b) 3-p-Tolylsulfanylbenzaldehyde;
- c) 3-(p-Tosyl)benzaldehyde;
- d) 3-Benzylaminobenzaldehyde;
- e) 3-Phenylaminobenzaldehyde;
- f) 3-Benzoylbenzaldehyde;
- g) 3-(α -Fluorobenzyl)benzaldehyde;
- h) 3-(4-Fluorophenoxy)benzaldehyde;
- i) 3-(2-Fluorophenoxy)benzaldehyde;
- j) 3-(3-Fluorophenoxy)benzaldehyde;
- k) 3-(Naphth-2-yloxy)benzaldehyde;
- l) 3-(Naphth-1-yloxy)benzaldehyde;
- m) 3-(Pyridin-3-ylamino)benzaldehyde;
- n) 3-(Pyridin-4-ylamino)benzaldehyde;
- o) 3-(Pyridin-2-ylamino)benzaldehyde; and
- p) 3-(Pyridin-2-yloxy)benzaldehyde.

Example 18

2-Naphth-2-ylethylamine

Combine naphth-2-ylacetonitrile (1.0 g, 6.0 mmol) and nickel (II) chloride hexahydrate (0.7g, 3.0 mmol) and tetrahydrofuran (30 ml). Add dropwise borane-tetrahydrofuran complex, 1M solution in tetrahydrofuran (24.0 ml, 24.0 mmol). After 1

-37-

hour, evaporate to give a residue. Chromatograph on silica gel eluting with 8:2 EtOAc:MeOH + 2% NH₄OH) to give the title compound.

By the method of Example 18 the following compounds were prepared: 2-Naphth-1-ylethylamine.

5

Example 19

5-Methanesulfonyltryptamine

Combine 2-(3-Chloropropyl)-(1,3)-dioxolane (6.69 g, 44.5 mmol), (4-methanesulfonylphenyl) hydrazine hydrochloride (9.92 g, 44.5 mmol), and Na₂HPO₄ (1.58 g, 11.1 mmol) in 300 ml of methanol/water (1:1). Heat to reflux. After 4.5 hours, cool to ambient temperature, then evaporate to residue. Dissolve the residue in 1 N NaOH and extract with dichloromethane. Combine the organic extracts, wash with brine, dry (Na₂SO₄), filter, then concentrate to residue. Chromatograph the residue on silica gel eluting with dichloromethane/2N NH₃(methanol) (84/16) to give the title compound as a light brown solid: mp 134-138 °C, MS (ACPI): m/e 239.1 (M+1). Analysis for C₁₁H₁₄N₂O₂S: Calcd: C, 55.44; H, 5.92; N, 11.76; found: C, 55.33; H, 5.97; N, 11.48.

15

Example 20

N-t-Butoxycarbonyl-2-(6-chloro-1H-indol-3-yl)ethylamine

Combine di-tert-butyl dicarbonate (1.2 g, 5.34 mmol), 6-chlorotryptamine (866.4 mg, 4.45 mmol) and NaHCO₃ (598.2 mg) in dioxane (50 ml). Stir at ambient temperature. After 15 hours, evaporate to residue, partition the residue between water and dichloromethane. Separate the layer and extract the aqueous layer with dichloromethane. Combine the organic extracts, wash with brine, dry (Na₂SO₄), filter and then evaporated to give the title compound as a light yellow oil.

20

25

Example 21A

N-Methyl-2-(6-chloro-1H-indol-3-yl)ethylamine

Combine N-t-butoxycarbonyl-2-(6-chloro-1H-indol-3-yl)ethylamine (1.3 g, 4.41 mmol) and dry THF (20 ml) and add dropwise to an ice bath cooled suspension of LiAlH₄ (1.0 g, 26.5 mmol) in dry THF (30 ml). Heat to reflux. After 2 hours, cool to ambient temperature and stir. After 15 hours, quench with saturated NaSO₄ (100 ml/mol), stir for 1 hour at ambient temperature, then filter under vacuum. Wash precipitate with THF and evaporated filtrate and washes to residue. Chromatograph the residue on silica gel eluting with dichloromethane/2N NH₃(methanol) (84/16) to give the title compound: MS (ACPI): m/e 209.0 (M+1).

30

35

-38-

Example 21B5-(4-Fluorophenyl)-1H-indole

Combine 5-bromoindole (5.00 g, 25.50 mmol) and $\text{Pd}(\text{Ph}_3\text{P})_4$ (1.47 g, 1.28 mmol) in toluene (510 ml). After 30 minutes, add a solution of 4-fluorobenzeneboronic acid (5.35 g, 38.26 mmol) in ethanol (153 ml) then add saturated NaHCO_3 (255 ml). Heat to reflux. After 4 hours, cool to ambient temperature, pour into saturated NaCl (250 ml), and separate the organic layer. Extract the aqueous layer with ethyl acetate. Combine the organic extracts, wash with brine, dried, then evaporate to residue. Chromatograph the residue on silica gel eluting with ethyl acetate/hexanes (10/90) to give the title compound: mp 84-89 °C. MS (ACPI): m/e 212.0 (M+1). Analysis for $\text{C}_{14}\text{H}_{10}\text{FN}$: Calcd: C, 79.60; H, 4.77; N, 6.63; found: C, 79.33; H, 4.92; N, 6.64.

By the method of Example 21 the following compounds were prepared: a) 5-Phenyl-1H-indole: mp 71-74 °C. MS (ACPI): m/e 194.0 (M+1). Analysis for $\text{C}_{14}\text{H}_{11}\text{N}$: Calcd: C, 87.01; H, 5.74; N, 7.25; found: C, 86.67; H, 5.82; N, 7.31.

b) 4-Phenylphenethylamine hydrochloride: (Exception-Chromatograph the residue on silica gel using dichloromethane/2N NH_3 (methanol) (90/10) to give the final product. The HCl salt was prepared in ethyl acetate: MS (ACPI): m/e 198.1 (M+1). Analysis for $\text{C}_{14}\text{H}_{16}\text{ClN}$: Calcd: C, 71.94; H, 6.90; N, 5.99; found: C, 71.66; H, 6.90; N, 5.94.

Example 227-Cyano-1H-indole

Combine 7-bromoindole (4.72 g, 24.0 mmol) and copper cyanide (4.30 g, 48.1 mmol) in 1-methyl-2-pyrrolidine (40 mL). Heat to 200 °C. After 2.5 hours, cool to room temperature, add water-ethyl acetate (200 mL, 1/1) to give a solid. Filter through the celite, extract the filtrate with ethyl acetate, combine the organic layers, wash with brine, dry over Na_2SO_4 , filter and evaporate *in vacuo* to give a residue. Chromatograph the residue on silica gel eluting with hexanes : ethyl acetate (10 :1) to give (1.87 g) of the title compound as a yellow solid: ^1H NMR (300 MHz, $\text{DMSO}-d_6$) 6.64-6.66 (m, 1H), 7.17 (t, 1H, $J = 7.6$ Hz), 7.51-7.53 (m, 1H), 7.60-7.62 (m, 1H), 7.94 (d, 1H, $J = 8.0$ Hz), 12.03 (br, 1H).

Example 233-Formyl-5-cyano-1H-indole

Add phosphorous oxychloride (11.76 g, 76.67 mmol) dropwise to DMF (24.3 ml) while maintaining the temperature at less than 10°C. Warm to ambient temperature and stir for 15 minutes at ambient temperature. Add dropwise 5-cyanoindole (10.00 g, 70.34 mmol) as a solution in DMF (30 ml) while keeping the temperature below 35°. After 1

-39-

hour, pour the reaction mixture into ice/water (300 ml) and then add 5N NaOH (54 ml) with stirring. Add slowly an additional amount of 5N NaOH (19.7 ml) and then heated to 90° for 1 minute and then cooled to ambient temperature to give a precipitate. Filter the precipitate and washed with water and dry to give the title compound: mp 248-250 °C.

5 MS (ACPI): m/e 171.0 (M+1). Analysis for C₁₀H₆N₂O: Calcd: C, 70.58; H, 3.55; N, 16.46; found: C, 70.41; H, 3.53; N, 16.33.

By the method of Example 23 the following compounds were prepared: a) 3-Formyl-5-(4-fluorophenyl)-1H-indole; mp 215-217 °C. MS (ACPI): m/e 239.1 (M+1). Analysis for C₁₅H₁₀FNO: Calcd: C, 75.30; H, 4.21; N, 5.85; found: C, 74.94; H, 4.17; N, 5.84;

10

b) 3-Formyl-5-phenyl-1H-indole; mp >250 °C. MS (ACPI): m/e 222.1 (M+1). Analysis for C₁₅H₁₁NO: Calcd: C, 81.43; H, 5.01; N, 6.33; found: C, 81.04; H, 5.05; N, 6.36;

c) 3-Formyl-6-methyl-1H-indole; mp 178-180 °C. MS (ACPI): m/e 159.9 (M+1). Analysis for C₁₀H₉NO: Calcd: C, 75.45; H, 5.70; N, 8.80; found: C, 75.60; H, 5.78; N, 8.97;

15

d) 3-Formyl-6-cyano-1H-indole; mp 246 °C. MS (ACPI): m/e 171.0 (M+1). Analysis for C₁₀H₆N₂O: Calcd: C, 70.58; H, 3.55; N, 16.46; found: C, 70.51; H, 3.59; N, 16.40; and

20 e) 3-Formyl-6-trifluoromethoxy-1H-indole; mp 189-192 °C. MS (ACPI): m/e 230.0 (M+1). Analysis for C₁₀H₆F₃NO₂: Calcd: C, 52.41; H, 2.64; N, 6.11; found: C, 52.31; H, 2.61; N, 6.09.

f) 3-Formyl-7-cyano-1H-indole; ¹H NMR (300 MHz, DMSO-*d*₆) δ 7.41 (t, 1H, *J* = 7.6 Hz), 7.80-7.82 (m, 1H), 8.42-8.50 (m, 2H), 10.02 (s 1H), 13.06 (br, 1H).

25 g) 3-Formyl-6-bromo-1H-indole; mp 197-200 °C. Analysis for C₉H₆BrNO: Calcd: C, 48.25; H, 2.70; N, 6.25; found: C, 47.87; H, 2.68; N, 6.17.

h) 3-Formyl-7-fluoro-1H-indole; mp 211-214 °C. MS(ACPI): m/e 163.9 (M+1). Analysis for C₉H₆FNO: Calcd: C, 66.26; H, 3.71; N, 8.59; found: C, 66.12; H, 3.67; N, 8.56.

30

Example 25

3-(2-Nitrovinyl)-5-cyano-1H-indole

Combine 3-formyl-5-cyano-1H-indole (10.60 g, 63.32 mmol) and a solution of ammonium acetate (10.60 g) in nitromethane (660 ml). Heat to 90°C. After 2 hours, cool to ambient temperature to give a precipitate. Collect the precipitate by filtration, wash with 1:1

35

-40-

MeOH/water (500 ml), and dry to give the title compound: mp 247-251 °C. MS (ACPI): m/e 214.0 (M+1).

By the method of Example 25 the following compounds were prepared: a) 3-(2-Nitrovinyl)-5-(4-fluorophenyl)-1H-indole; mp 217-220 °C. MS (ACPI): m/e 282.2 (M+1).

5 Analysis for $C_{15}H_{10}FN_2O_2$: Calcd: C, 68.08; H, 3.93; N, 9.92; found: C, 67.73; H, 3.92; N, 9.73;

b) 3-(2-Nitrovinyl)-5-phenyl-1H-indole; MS (ACPI): m/e 265.1 (M+1);

c) 3-(2-Nitrovinyl)-6-methyl-1H-indole; MS (ACPI): m/e 203.1 (M+1);

d) 3-(2-Nitrovinyl)-6-cyano-1H-indole; mp >250 °C. MS (ACPI): m/e 212.0 (M-1). Analysis for $C_{11}H_7N_3O_2$: Calcd: C, 61.97; H, 3.31; N, 19.71; found: C, 62.09; H, 3.34; N, 20.06; and

e) 3-(2-Nitrovinyl)-6-trifluoromethoxy-1H-indole; mp 139-143 °C. MS (ACPI): m/e 273.0 (M+1).

f) 3-(2-Nitrovinyl)-6-phenoxy-1H-indole: 1H NMR (DMSO d_6) 12.1 (s, 1H), 8.38-8.34 (d, 1H), 8.20-8.19 (m, 1H), 8.01-7.97 (m, 2H), 7.39-7.35 (m, 2H), 7.14-7.07 (m, 2H), 7.02-7.00 (m, 2H), 6.95-6.92 (m, 1H).

g) 3-(2-Nitrovinyl)-5-(pyridin-3-yloxy)-1H-indole: ISMS 282 (M+1); 1H NMR (DMSO- d_6) 9.5 (bs, 1H), 8.36-8.32 (m, 2H), 8.26-8.24 (m, 2H), 7.98-7.95 (m, 1H), 7.79-7.78 (m, 1H), 7.55-7.53 (m, 1H), 7.34-7.31 (m, 1H), 7.27-7.24 (m, 1H), 7.02-7.00 (m, 1H).

h) 3-(2-Nitrovinyl)-7-cyano-1H-indole: 1H NMR (300 MHz, $CDCl_3$) δ 7.36 (t, 1H, J = 7.7 Hz), 7.76 (d, 1H, J = 7.2 Hz), 8.09-8.14 (m, 1H), 8.36-8.46 (m, 3H); MS (electrospray), m/e: 212.1 (M-1)

i) 3-(2-Nitrovinyl)-6-bromo-1H-indole; mp 210 °C, dec. Analysis for $C_{10}H_7BrN_2O_2$: Calcd: C, 44.97; H, 2.64; N, 10.49; found: C, 44.62; H, 2.70; N, 10.49.

j) 3-(2-Nitrovinyl)-7-fluoro-1H-indole; mp 176-180 °C. MS (ACPI): m/e 207.1 (M+1). Analysis for $C_{10}H_7FN_2O_2$: Calcd: C, 58.26; H, 3.42; N, 13.59; found: C, 58.01; H, 3.31; N, 13.26.

30 Example 26

3-(2-Nitroethyl)-5-cyano-1H-indole

Add sodium borohydride (25.65 g, 678 mmol) to an ice bath cooled solution of 3-(2-nitrovinyl)-5-cyano-1H-indole (12.68 g, 59.5 mmol) in 1:1 MeOH/DMF (600 ml). After 1.5 hours, add brine (600 ml) then adjust the pH to approximately 7 with 5N HCl.

35 Evaporate under reduced pressure to remove the methanol and then extract with dichloromethane. Combine the organic extracts, wash with brine, dry (Na_2SO_4), then evaporate to residue. Chromatograph the residue on silica gel eluting with

-41-

dichloromethane to give, after evaporation, the title compound as colorless prisms: mp 132-136 °C. MS (ACPI): m/e 215.0 (M+1). Analysis for $C_{11}H_9N_3O_2$: Calcd: C, 61.39; H, 4.22; N, 19.52; found: C, 61.09; H, 4.10; N, 19.16.

By the method of Example 26 the following compounds were prepared: a) 3-(2-nitro-ethyl)-7-cyano-1H-indole; 1H NMR (300 MHz, DMSO- d_6) 3.39 (t, 2H, $J = 6.9$ Hz), 4.87 (t, 2H, $J = 7.0$ Hz), 7.17 (t, 1H, $J = 7.4$ Hz), 7.38 (m, 1H), 7.63 (d, 1H, $J = 7.4$ Hz), 7.99 (d, 1H, $J = 7.9$ Hz), 11.96 (br, 1H). MS (electrospray), m/e: 214.1 (M-1).

Example 27

10 5-Cyanotryptamine

Add zinc powder (16.22 g, 248.1 mmol) in four portions to 2N HCl (300.0 ml). Add dropwise 3-(2-nitroethyl)-5-cyano-1H-indole (2.25 g, 10.5 mmol) as a solution in methanol (300.0 ml). Heat to reflux. After 2.5 hours, cool to ambient temperature and adjust the pH to 11 using 5N NaOH, filter through Celite, wash with water, then extract the filtrate with dichloromethane. Combine the organic extracts, wash with brine, dry (Na₂SO₄), then evaporated to give the title compound: mp 102-105 °C, MS (ACPI): m/e 186.1 (M+1). Analysis for $C_{11}H_{11}N_3$: Calcd: C, 71.33; H, 5.99; N, 22.69; found: C, 71.03; H, 5.91; N, 22.64.

By the method of Example 27 the following compounds were prepared: a) 3-(2-aminoethyl)-7-cyano-1H-indole; 1H NMR (300 MHz, DMSO- d_6) 2.76-2.82 (m, 4H), 7.15 (t, 1H, $J = 7.6$ Hz), 7.31 (s, 1H), 7.58 (d, 1H, $J = 7.4$ Hz), 7.91-7.94 (m, 1H), 11.80 (br, 1H); MS (electrospray), m/e: 186.1 (M+1), 184.1 (m-1).

By the method of Example 27 the following compound was prepared: a) 6-Bromotryptamine; mp 114-116 °C. Analysis for $C_{10}H_{11}BrN_2$: Calcd: C, 50.23; H, 4.64; N, 11.72; found: C, 49.96; H, 4.49; N, 11.47.

Example 28

N-t-Butoxycarbonyl-2-(5-cyano-1H-indol-3-yl)ethylamine

Combine di-tert-butyl dicarbonate and a solution of 5-cyanotryptamine (1.33 g, 7.15mmol) and 2N NaOH (4.2 ml) in THF (60 ml). After 3 hours at ambient temperature, evaporate to residue. Dissolve the residue in water and extract with dichloromethane. Combine the organic extracts, wash with brine, dry (Na₂SO₄), and evaporate to residue. Chromatograph the residue on silica gel eluting with dichloromethane/2N NH₃(methanol) (97/3)) to give the title compound: mp 129-132 °C. MS (ACPI): m/e 286.2 (M+1). Analysis for $C_{16}H_{19}N_3O_2$: Calcd: C, 67.35; H, 6.71; N, 14.73; found: C, 67.16; H, 6.68; N, 14.46.

-42-

By the method of Example 28 the following compounds were prepared: a) N-t-Butoxycarbonyl-2-(6-cyano-1H-indol-3-yl)ethylamine.

Example 29

5 N-t-Butoxycarbonyl-2-(5-amido-1H-indol-3-yl)ethylamine

Combine water (64.0 ml) and NaOH (8.53 g) and cool to about 5°C in an ice bath. Add a solution of N-t-butoxycarbonyl-2-(5-cyano-1H-indol-3-yl)ethylamine (1.85 g, 6.50 mmol) in ethanol (128.0 ml) then added to the chilled solution. Add dropwise 30% peroxide (6.4 ml) while keeping the temperature below 5°C. After 30 minutes, warm to
10 ambient temperature. After 22 hours, decompose the excess peroxide by adding a 20% sodium bisulfite solution (45.0 ml) with stirring. After 30 minutes, evaporate under reduced pressure to remove the ethanol and extract the resulting aqueous solution with ethyl acetate. Combine the organic extracts, wash with brine, dry (Na₂SO₄), then evaporate to residue. Chromatograph the residue on silica gel eluting with
15 dichloromethane/2N NH₃(methanol) (96/4) to give, after evaporation, the title compound: mp 65-68 °C. MS (ACPI): m/e 304.2 (M+1). Analysis for C₁₆H₂₁N₃O₃: Calcd: C, 63.35; H, 6.98; N, 13.85; found: C, 63.26; H, 6.99; N, 13.71.

By the method of Example 28 the following compounds were prepared: a) N-t-Butoxycarbonyl-2-(6-amido-1H-indol-3-yl)ethylamine: MS (ACPI): m/e 302.3 (M-1).
20

Example 30

5-Amidotryptamine

Dissolve N-t-butoxycarbonyl-2-(5-amido-1H-indol-3-yl)ethylamine (1.83 g, 6.04 mmol) in dioxane (25.0 ml). Add dropwise a 4M HCl in dioxane (7.5 ml). After 18
25 hours, collect the solid by filtration and wash with diethyl ether to give, after drying, the title compound as the hydrochloride: mp 192-195 °C. MS (ACPI): m/e 202.0 (M+1).

By the method of Example 30 the following compounds were prepared: a) 6-Amidotryptamine: mp 169-173 °C. MS (ACPI): m/e 204.1 (M+1).

Example 32

5-(4-Fluorophenyl)tryptamine

Combine LiAlH₄ (2.66 g, 70.17 mmol) and dry THF (70.0 ml) and cool the suspension in an ice bath. Add dropwise a solution of 5-(4-fluorophenyl)-3-(2-nitrovinyl)-1H-indole (3.30 g, 11.69 mmol) in dry THF (30.0 ml). Heat to reflux. After 1
35 hour, cool to ambient temperature and stir. After 15 hours, quench with saturated Na₂SO₄ (100 ml/mol) and stir at ambient temperature. After 1 hour, filter, rinse the precipitate with THF, and evaporate the filtrate to residue. Chromatograph the residue on silica gel

-43-

eluting with dichloromethane/2N NH₃(methanol) (80/20) to give the title compound.

Prepare the HCl salt in diethyl ether: mp >250 °C. MS (ACPI): m/e 255.0 (M+1).

Analysis for C₁₆H₁₆ClFN₂: Calcd: C, 66.09; H, 5.55; N, 9.63; found: C, 65.78; H, 5.48; N, 9.58.

- 5 By the method of Example 32 the following compounds were prepared: a) 5-Phenyltryptamine; mp 244-246 °C. MS (ACPI): m/e 237.1 (M+1). Analysis for C₁₆H₁₇ClN₂: Calcd: C, 70.45; H, 6.28; N, 10.27; Found: C, 70.75; H, 6.33; N, 10.27. (isolated as the hydrochloride);

- b) 6-Methyltryptamine; mp 139-141 °C. MS (ACPI): m/e 175.0 (M+1). Analysis
10 for C₁₁H₁₄N₂: Calcd: C, 75.82; H, 8.10; N, 16.08; Found: C, 76.05; H, 8.26; N, 16.12.

c) 6-Trifluoromethoxytryptamine; MS (ACPI): m/e 245.0 (M+1). Analysis for C₁₁H₁₁F₃N₂O: Calcd: C, 54.10; H, 4.54; N, 11.47; Found: C, 53.92; H, 4.50; N, 11.06.

d) 7-Fluorotryptamine; MS (ACPI): m/e 179.0 (M+1). Analysis for C₁₀H₁₁FN₂: Calcd: C, 67.40; H, 6.22; N, 15.72; Found: C, 67.06; H, 6.11; N, 15.48.

15

Example 33

6-Ethoxycarbonyl-1H-indole

- Combine 6-carboxy-1H-indole and ethanol (110 ml) and cool to 5°C. Add dropwise concentrated H₂SO₄ (96%, 11.08 ml) while keeping the temperature below
20 10°C. Heat to reflux. After 4 hours, cool and pour onto ice/water, adjust the pH to about pH 9 and extract with ethyl acetate. Combine the organic extracts, wash with brine, dry (Na₂SO₄), then concentrate to residue. Chromatograph the residue on silica gel eluting with chloroform to give, after evaporation, the title compound: mp 72-75 °C. MS (ACPI): m/e 189.9 (M+1).

25

Example 34

3-(2-Nitrovinyl)-6-ethoxycarbonyl-1H-indole

- Combine 1-dimethylamino-2-nitroethylene (1.93 g, 16.58 mmol) and TFA (10.0 ml) and stir until dissolved. Add 6-ethoxycarbonyl-1H-indole (3.14 g, 16.58 mmol) and
30 stir at ambient temperature. After 1 hour, pour the reaction mixture into ice/water, extract with ethyl acetate, then evaporate to residue. Stir the residue in warm ethanol, cool to ambient temperature, then filter and dry to give the title compound as a dark yellow powder: mp 241 °C. MS (ACPI): m/e 261.1 (M+1). Analysis for C₁₃H₁₂N₂O₄: Calcd: C, 60.00; H, 4.65; N, 10.76; found: C, 59.99; H, 4.63; N, 10.59.

35

-44-

Example 353-(2-Nitroethyl)-6-ethoxycarbonyl-1H-indole

Combine 3-(2-nitrovinyl)-6-ethoxycarbonyl-1H-indole (4.0 g, 15.37 mmol) and NaBH₄ (726.7 mg, 19.21 mmol) in 100 ml of THF/Methanol (9:1) and stir at ambient temperature. After 1.5 hours, concentrate to residue. Partition the residue between brine and ethyl acetate, wash with brine, combine the organic layers, dry (Na₂SO₄), then evaporate to give the title compound as a yellow powder: mp 124-127 °C. MS (ACPI): m/e 263.0 (M+1). Analysis for C₁₃H₁₄N₂O₄: Calcd: C, 59.54; H, 5.38; N, 10.68; found: C, 59.40; H, 5.36; N, 10.53.

By the method of Example 35 the following compounds were prepared: a) 3-(2-Nitroethyl)-6-cyano-1H-indole: m/e 214.1 (M-1). Analysis for C₁₁H₉N₃O₂: Calcd: C, 61.39; H, 4.22; N, 19.52; found: C, 61.05; H, 4.09; N, 19.19.

b) 3-(2-Nitroethyl)-6-bromo-1H-indole;

c) 3-(2-Nitroethyl)-6-methanesulfonyl-1H-indole; mp 162-164 °C. MS (ACPI): m/e 269.1 (M+1).

d) 3-(2-Nitroethyl)-6-benzenesulfonyl-1H-indole (exception: 75 ml of THF was used).

Example 366-Ethoxycarbonyltryptamine

Combine Pt₂O (440 mg) and a solution of 3-(2-nitroethyl)-6-ethoxycarbonyl-1H-indole (3.55 g, 13.54 mmol) in ethyl acetate (100 ml). Hydrogenate at 60 psi (410 kPa) and ambient temperature. After 4 hours, filter through celite and concentrate the filtrate to residue. Chromatograph the residue on silica gel eluting with dichloromethane/2N NH₃(methanol) (85/15) to give, after evaporation, the title compound as an off-white powder: mp 127-131 °C. MS (ACPI): m/e 233.0 (M+1). Analysis for C₁₃H₁₆N₂O₂: Calcd: C, 67.22; H, 6.94; N, 12.06; found: C, 66.87; H, 6.86; N, 11.86.

By the method of Example 36 the following compounds were prepared: a) 6-Cyanotryptamine: mp 144-147 °C. MS (ACPI): m/e 186.0 (M+1). Analysis for C₁₁H₁₁N₃: Calcd: C, 71.33; H, 5.99; N, 22.69; found: C, 71.10; H, 5.89; N, 22.38.

b) 6-Methanesulfonyltryptamine: mp 149-153 °C. MS (ACPI): m/e 239.1 (M+1). Analysis for C₁₁H₁₄N₂O₂S: Calcd: C, 55.44; H, 5.92; N, 11.76; found: C, 55.12; H, 5.82; N, 11.97.

c) 6-Benzenesulfonyltryptamine: mp 169-172 °C. MS (ACPI): m/e 301.0 (M+1).

-45-

Example 386-Trifluoromethoxy-1H-indole

Combine 1-methyl-4-trifluoromethoxybenzene (5.44 g, 30.87 mmol) and H₂SO₄ (96%, 30.9 ml). Cool to about 0°C. Add dropwise fuming HNO₃ (2.06 g, 32.72 mmol) while maintaining the temperature below 10°C. When the addition is complete, warm to ambient temperature. After 2.5 hours, pour the mixture onto ice/water, extract with dichloromethane. Combine the organic extracts, wash with brine, dry (Na₂SO₄), then concentrate to residue. Chromatograph the residue on silica gel eluting with hexanes/ethyl acetate (75/25) to give, after evaporation, 1-methyl-2-nitro-4-trifluoromethoxybenzene: MS (ACPI): m/e 220.1 (M-1).

Combine 1-methyl-2-nitro-4-trifluoromethoxybenzene (3.73 g, 16.86 mmol), pyrrolidine (1.32 g, 18.55 mmol), N,N-dimethylformamide dimethyl acetal (6.03 g, 50.58 mmol), and dry DMF (35 ml). Heat to about 105°. After 19 hours, remove the DMF under reduced pressure to give an oily residue. Combine the residue and ethyl acetate, wash with brine, dry (Na₂SO₄), then concentrate to give N,N-dimethyl-2-(2-nitro-4-trifluoromethoxyphenyl)vinylamine.

Combine N,N-dimethyl-2-(2-nitro-4-trifluoromethoxyphenyl)vinylamine (4.64 g, 16.8 mmol) and Raney® nickel (900 mg) in ethanol (100 ml). Hydrogenate at 60 psi (410 kPa) and ambient temperature. After 18 hours, filter through celite, concentrate the filtrate to residue, and chromatograph on silica gel eluting with hexanes/ethyl acetate (30/70) to give, after evaporation, the title compound as an off-white powder: mp 59 °C. MS (ACPI): m/e 200.0 (M-1).

Example 392-Phenylphenethylamine

Combine 2-phenylphenylacetonitrile (4.69 g, 24.26 mmol) and diethyl ether (10 ml) and add dropwise to a cooled (-10°) solution of LiAlH₄ (2.76 g, 72.81 mmol) in diethyl ether (100 ml). Warm to ambient temperature. After 2 hours, quench with saturated sodium sulfate (100 ml/mol). Filter to remove the precipitate, dry (Na₂SO₄) the filtrate, filter, and concentrate to residue. Chromatograph the residue on silica gel eluting with dichloromethane/2N NH₃(methanol) (95/5) to give the title compound as a yellow oil. Prepare the HCl salt in diethyl ether: mp 197-199 °C. MS (ACPI): m/e 198.1 (M+1). Analysis for C₁₄H₁₆ClN: Calcd: C, 71.94; H, 6.90; N, 5.99; found: C, 72.15; H, 6.84; N, 6.09.

35

-46-

Example 407-Chloro-1H-indole

By the method of *J. Med. Chem.* 1990, 33, 2777, add dropwise 2-chloroaniline (5.8 g, 45.45 mmol) in anhydrous toluene (80 mL) to a cold 1M solution of BCl₃ (50 mL) in dichloromethane. After addition, allow the reaction to stir at 0°C for 10 minutes. After 10 minutes at 0°C, add chloroacetonitrile (13.72 g, 11.53 mL, 181.8 mmol, 4 eq) and aluminum trichloride (6.67 g, 50 mmol, 1.1 eq) in 5 equal portions over 45 minutes and then heat to reflux (~65°C). After 6 hours, cool to room temperature. After 16 hours, cool the reaction in an ice bath and add 2N HCl (61.4 mL) and then heat the reaction to reflux. After 45 minutes, cool in an ice bath, neutralize the acid with 2N NaOH keeping the temperature of the reaction below 15°C until the pH is about 5. Transfer the reaction to a separatory funnel and remove the organic layer. Extract the aqueous layer with dichloromethane (2x200 mL). Combine the organic layers, dry over MgSO₄, filter, and remove the solvent *in vacuo* to give 1-(2-amino-3-chlorophenyl)-2-chloroethanone which may be used without further purification.

Dissolve 1-(2-amino-3-chlorophenyl)-2-chloroethanone (7.0 g, 34.30 mmol) in 10% aqueous 1,4-dioxane (75 mL). Carefully add NaBH₄ (2.6 g, 68.6 mmol, 2eq.) as a solid. Heat to reflux. After 4 hours, cool to room temperature, dilute with water (300 mL), and extract with dichloromethane (2x200 mL). Combine the organic layers, dry over MgSO₄, filter, and remove the solvent *in vacuo* leaving a light brown oil in the flask. Purify the oil by HPLC (silica gel mobile phase: 100% hexane to 50% EtOAc in hexanes over 50 minutes), to give the title compound as a brown oil: ¹H NMR (300MHz, d₆-DMSO-d₆): 5.16 (m, 1H), 5.39 (d, 1H), 5.70 (bs, 1H), 6.59 (t, 1H), 7.09 (m, 2H); MS (ES⁺): m/z 154, 152 (M+H)⁺.

By the method of Example 40 the following compounds were prepared: a) 5-Bromo-7-ethyl-1H-indole: ¹H NMR (300MHz, d₆-DMSO-d₆): 1.25 (t, 3H), 2.85 (m, 2H), 6.41 (m, 1H), 7.02 (M, 1H), 7.36 (m, 1H), 7.55 (m, 1H), 11.28 (bs, 1H); MS(ES⁺): m/z 224, 226 (M+H)⁺; Analysis for C₁₀H₁₀BrN: Calcd.; C, 53.60; H, 4.50; N, 6.25; found; C, 53.50; H, 4.34; N, 6.22.

Example 426-Trifluoromethyl-1H-indole

Combine 2-bromo-5-trifluoromethylphenylamine (27.06 g, 112.74 mmol) and 200 mL of pyridine. Cool in an ice bath and add ethyl chloroformate (18.35 g, 169.11 mL, 1.5 eq). After addition was complete, allow the reaction to gradually warming to room temperature. After 18 hours, evaporate *in vacuo* to give a residue. Dissolve the residue in Et₂O/water and transfer to a separatory funnel. Separate the layer and extract the aqueous

-47-

layer with Et₂O (2x300 mL), combine the organic layers, dry over MgSO₄, filter, and evaporate *in vacuo* to give N-(2-bromo-5-trifluoromethylphenyl)carbamic acid ethyl ester which may be used without further purification.

Following the procedure described in *J. Org. Chem.* 1997, 62, 6507, combine N-(2-bromo-5-trifluoromethylphenyl)carbamic acid ethyl ester (34.33 g, 110 mmol), triethylamine (300 mL), dichlorobis (triphenylphosphine)palladium(II) (5.4 g, 7.7 mmol), and copper (I) iodide (1.47 g, 7.7 mmol). Evacuate the dark solution and fill with N₂ twice and then quickly add (trimethylsilyl)acetylene (16.21 g, 165 mmol, 23.32 mL) with vigorous stirring. Heat to 80°C. When TLC indicates absence of the starting material, add water and Et₂O and filter through celite and evaporate the filtrate *in vacuo* to give a residue. Dilute residue with water and extract with Et₂O, combine the organic layers and remove the evaporate to give a dark brown oil. Absorb the oil onto silica gel and load onto a short column of silica gel equilibrated with 100% hexanes. Wash with 100% hexanes (2 L) and elute the product with 1% EtOAc in hexanes. Pool fractions containing the product and remove *in vacuo* the solvent to give 5-trifluoromethyl-2-trimethylsilanylethynyl phenylamine as a red/brown oil; MS (IS): m/z 330 (M+H)⁺.

Carefully add NaH (10.83 g, 60% in oil, 270.8 mmol, 4 eq.) to EtOH (200 mL). When cool, add a solution of 5-trifluoromethyl-2-trimethylsilanylethynylphenylamine (22.3 g, 67.7 mmol) in EtOH (400 mL) with vigorous stirring. After 2 hours, heat to reflux. After 4 hours, evaporate *in vacuo* to remove the EtOH and dilute the residue obtained with water and extract with Et₂O. Combine the organic layers and wash with brine, dry over MgSO₄, filter, and evaporate to give a dark oil. Absorb the oil onto silica gel and load onto short column of silica gel. Elute with 20% EtOAc in hexanes. Pool fractions containing the product and remove the solvent leaving a dark brown oil. Further purification of the oil by HPLC (silica column) using a gradient 1% Et₂O in hexanes to 15% Et₂O in hexanes. Pool fractions containing the product and remove the solvent to give the title compound as an orange solid: ¹H NMR (300MHz, d₆-DMSO-d₆): 6.58 (m, 1H), 7.28 (m, 1H), 7.61 (t, 1H), 7.74 (m, 2H), 11.51 (bs, 1H); MS(EI⁺): m/z 185 (M⁺).

By the method of Example 42 the following compounds were prepared: a) 5-Isopropyl-1H-indole: MS(ES⁺): m/z 160 (M+H)⁺; (ES)⁻: m/z 158 (M-H)⁻.

Example 44

6-Fluoro-5-methoxy-1H-indole

Dissolve fuming nitric acid (24 mL) in concentrated H₂SO₄ in a round bottom flask. Add 3,4-difluorobromobenzene (20 g, 104 mmol) dropwise via pipette with vigorous stirring. After addition, stir the reaction at room temperature for 2 hours, pour

-48-

the reaction into ice water and extract with Et₂O (2x250 mL). Collect and combine the organic layers, dry over MgSO₄, filter, and remove the solvent to give 1-bromo-4,5-difluoro-2-nitrobenzene as a light yellow oil.

5 Add 1-bromo-4,5-difluoro-2-nitrobenzene (24 g, 100 mmol) to a solution of sodium methoxide (1.2 eq) in MeOH. After addition, stir the reaction at room temperature for 2.5 hours. Remove the solvent *in vacuo* and dilute the residue with water and extract with Et₂O (2x250 mL). Combine the organic layers, dry over MgSO₄, filter, and the remove solvent *in vacuo* to give 1-bromo-4-fluoro-5-methoxy-2-nitrobenzene as a yellow solid: ¹H NMR (300 MHz, CDCl₃): 3.99 (s, 3H), 7.26 (m, 1H), 7.83 (d, 1H); MS(FD+):
10 m/z 249, 251 (M⁺); Analysis for C₇H₅BrFNO₃: Calcd.: C, 33.63; H, 2.02; N, 5.60; found: C, 33.79; H, 1.98; N, 5.62.

Combine 1-bromo-4-fluoro-5-methoxy-2-nitrobenzene (20.5 g, 82 mmol) and Pt-on-carbon (sulfided) in THF (600 mL) and hydrogenate at 60 psi (414 kPa) over for 4 hours. Filter the reaction through celite to remove the catalyst and remove the solvent to
15 give 2-bromo-5-fluoro-4-methoxyaniline as a brown solid which may taken on without further purification.

Using 2-bromo-5-fluoro-4-methoxyaniline, the method of Example 42 gives N-(2-bromo-5-fluoro-4-methoxyphenyl)carbamic acid ethyl ester as a brown solid: ¹H NMR (300 MHz, CDCl₃): 1.33 (t, 3H), 3.85 (s, 3H), 4.23 (q, 2H), 7.09 (d, 1H), 7.97 (bd, 1H);
20 MS(FD+): m/z 291, 293 (M⁺); N-(5-fluoro-4-methoxy-2-trimethylsilanylethynylphenyl)carbamic acid ethyl ester as a yellow solid: MS(ES+): m/z 310 (M+H)⁺; (ES-): m/z 308 (M-H)⁻; and the title compound as a solid: ¹H NMR (300MHz, CDCl₃): 3.93 (s, 3H), 6.48 (m, 1H), 7.15 (m, 3H), 8.11 (bs, 1H); MS(ES+): m/z 166 (M+H)⁺; (ES-): m/z 164 (M-H)⁻; Analysis for C₉H₈FNO: Calcd.: C, 65.45; H, 4.88; N, 8.48; found: C, 65.17; H, 4.97; N, 8.70.
25

Example 45

5,6-Difluoro-1H-indole

Using the method of Example 42 gives 2-bromo-4,5-difluoroaniline; which gives
30 N-(2-bromo-4,5-difluorophenyl)carbamic acid ethyl ester; which gives N-(4,5-Difluoro-2-trimethylsilanylethynylphenyl)carbamic acid ethyl ester; which gives the title compound as an orange solid: ¹H NMR (300MHz, d₆-DMSO-d₆): 6.43 (m, 1H), 7.38 (m, 2H), 7.50 (m, 1H), 11.25 (bs, 1H); MS(ES-): m/z 152 (M-H)⁻; Analysis for C₈H₅F₂N: Calcd.: C, 62.75; H, 3.29; N, 9.15; found: C, 62.41; H, 3.12; N, 8.98.

35

-49-

Example 465-Trifluoromethoxy-1H-indole

Using the method of Example 42 and 2-bromo-4-(trifluoromethoxy)aniline gives N-(2-bromo-4-trifluoromethoxyphenyl)carbamic acid ethyl ester: ¹H NMR (300MHz, CDCl₃): 1.34 (t, 3H), 4.25 (m, 2H), 7.19 (m, 1H), 7.41 (m, 1H), 8.20 (d, 1H); MS(ES⁻): m/z 326, 328 (M-H)⁻; Analysis for C₁₀H₉BrF₃NO₃: Calcd.; C, 36.6096; H, 2.7650; N, 4.2692; found; C, 36.50; H, 2.67; N, 3.97; which gives N-(4-Trifluoromethoxy-2-trimethylsilanylethynylphenyl) carbamic acid ethyl ester; which gives the title compound as a yellow oil: MS(ES⁻): m/z 200 (M-H)⁻.

10

Example 474-Phenyl-1H-indole

Using the method of Carrera and Sheppard in *Synlett*. 1994, 93-94, 4-bromoindole gives the title compound: ¹H NMR (300MHz, d₆-DMSO-d₆): 6.56 (m, 1H), 7.08 (m, 1H), 7.17 (m, 1H), 7.43 (m, 5H), 7.67 (m, 2H), 11.27 (bs, 1H); MS(ES⁺): m/z 194 (M+H)⁺; (ES⁻): m/z 192 (M-H)⁻.

15

Example 48(2-Nitro-5-trifluoromethylphenyl)-acetonitrile

By the method of Liebigs Ann. Chem. 1988, 203-208, using 4-nitrobenzotrifluoride (15 g, 78.49 mmol) gives the title compound: MS(ES⁻): m/z 229 (M-H)⁻.

20

Example 495-Trifluoromethyl-1H-indole

By the method of Liebigs Ann. Chem. 1988, 203-208 using (2-nitro-5-trifluoromethylphenyl)acetonitrile gives the title compound as a white solid: ¹H NMR (300MHz, d₆-DMSO-d₆): 6.60 (m, 1H), 7.36 (m, 1H), 7.53 (m, 1H), 7.57 (m, 1H), 7.94 (m, 1H), 11.51 (bs, 1H); MS(ES⁻): m/z 184 (M-H)⁻.

25

30

Example 503-Formyl-4-methoxy-1H-indole

Add phosphorus oxychloride (1.1 eq.) with vigorous stirring to DMF (cooled in an ice bath). After the addition is complete, allow to stir in the ice bath for ~10 minutes, then add a solution of 4-methoxy-1H-indole (5 g) in anhydrous DMF with vigorous stirring. Allow to stir at 0°C. After 1 hour, warm to room temperature. After 16 hours, carefully add 4 eq. of 2N NaOH with vigorous stirring. Heat to about 80°C and then cool. Pour the

35

-50-

reaction mixture into cold water with vigorous stirring to give a solid. Collect the solid by filtration and dry in a vacuum oven at room temperature to give the title compound.

Acidify the filtrate and extract with EtOAc. Combine the organic layers and wash with 50% brine. Collect the organic layer, dry (MgSO_4) filter, and remove the solvent to give

5 the title compound as a light purple solid. Total yield of the title compound is 5.44 g: MS (ES⁺): m/z 175 (M+H)⁺, 160 (M-CH₃)⁺; (ES⁻): m/z 174 (M-H)⁻.

- By the method of Example 50 the following compounds were prepared: a) 3-Formyl-6-methoxy-1H-indole, ¹H NMR (300 MHz, d₆-DMSO): 3.79 (s, 3H); 6.85 (dd, 1H); 6.98 (m, 1H); 7.92 (d, 1H); 8.15 (s, 1H); 9.86 (s, 1H); 11.92 (bs, 1H); MS (ES⁺): m/z 176 (M+H)⁺; (ES⁻): m/z 174 (M-H)⁻;
- 10 b) 3-Formyl-7-methoxy-1H-indole;
- c) 3-Formyl-4-chloro-1H-indole;
- d) 3-Formyl-6-chloro-1H-indole, ¹H NMR (300MHz, d₆-DMSO-d₆): 7.24 (dd, 1H), 7.56 (d, 1H), 8.06 (d, 1H), 8.33 (s, 1H), 9.93 (s, 1H), 12.21 (bs, 1H); MS (ES⁺): m/z 182, 180 (M+H)⁺; (ES⁻): m/z 180, 178 (M-H)⁻;
- 15 e) 3-Formyl-7-chloro-1H-indole, ¹H NMR (300MHz, d₆-DMSO): 7.23 (t, 1H), 7.35 (d, 1H), 8.05 (d, 1H), 8.38 (bs, 1H), 9.95 (s, 1H), 12.54 (bs, 1H); MS (ES⁺): m/z 182, 180 (M+H)⁺; (ES⁻): m/z 180, 178 (M-H)⁻;
- f) 3-Formyl-4-fluoro-1H-indole, ¹H NMR (300MHz, d₆-DMSO): 7.01 (m, 1H), 7.24 (m, 1H), 7.36 (d, 1H), 8.30 (s, 1H), 10.03 (d, 1H), 12.48 (bs, 1H); MS (ES⁺): m/z 164 (M+H)⁺; (ES⁻): m/z 162 (M-H)⁻;
- 20 g) 3-Formyl-5-methoxy-6-trifluoromethyl-1H-indole, ¹H NMR (300MHz, d₆-DMSO): 3.91 (s, 3H), 7.77 (dd, 1H), 7.95 (bs, 1H), 8.42 (s, 1H), 9.96 (s, 1H), 12.29 (bs, 1H); MS (ES⁺): m/z 244 (M+H)⁺; (ES⁻): m/z 242 (M-H)⁻;
- 25 h) 3-Formyl-6-chloro-5-methoxy-1H-indole, ¹H NMR (300MHz, d₆-DMSO): 3.88 (s, 3H), 7.58 (s, 1H), 7.71 (s, 1H), 8.26 (s, 1H), 9.91 (s, 1H), 12.08 (bs, 1H); MS (ES⁺): m/z 210, 212 (M+H)⁺; (ES⁻): m/z 208, 210 (M-H)⁻;
- i) 3-Formyl-4-chloro-5-methoxy-1H-indole, ¹H NMR (300MHz, d₆-DMSO): 3.89 (s, 3H), 7.13m (dd, 1H), 7.47 (dd, 1H), 8.23 (s, 1H), 10.5 (s, 1H), 12.39 (bs, 1H); MS (ES⁺): m/z 210, 212 (M+H)⁺; (ES⁻): m/z 208, 210 (M-H)⁻;
- 30 j) 3-Formyl-6-trifluoromethyl-1H-indole, ¹H NMR (300MHz, d₆-DMSO): 7.52 (d, 1H), 8.27 (d, 1H), 8.51 (m, 1H), 9.99 (s, 1H), 12.47 (bs, 1H). MS(ES⁺): m/z 214 (M+H)⁺; (ES⁻): m/z 212 (M-H)⁻;
- k) 3-Formyl-5-methoxy-2-methyl-1H-indole, ¹H (300MHz, d₆-DMSO): 2.65 (s, 3H), 3.76 (s, 3H), 6.78 (dd, 1H), 7.27 (d, 1H), 7.56 (m, 1H), 10.00 (s, 1H), 11.85 (bs, 1H); MS(ES⁺): m/z 190 (M+H)⁺; (ES⁻): m/z 188 (M-H)⁻;
- 35

-51-

- l) 3-Formyl-6-fluoro-5-methoxy-1H-indole, ^1H NMR (300MHz, d6-DMSO): 3.87 (s, 3H), 7.35 (d, 1H), 7.71 (d, 1H), 8.21 (s, 1H), 9.89 (s, 1H), 12.03 (bs, 1H); MS(ES⁺): m/z 194 (M+H)⁺; (ES⁻): m/z 192 (M-H)⁻;
- m) 3-Formyl-5,6-difluoro-1H-indole, ^1H NMR (300MHz, d6-DMSO): 7.56 (m, 1H), 7.92 (m, 1H), 8.36 (s, 1H), 9.92 (s, 1H), 12.25 (bs, 1H); MS(ES⁺): m/z 182 (M+H)⁺ (ES⁻): m/z 180 (M-H)⁻;
- n) 3-Formyl-6-chloro-5-fluoro-1H-indole, ^1H NMR (300MHz, d6-DMSO): 7.72 (d, 1H), 7.91 (d, 1H), 8.40 (s, 1H), 9.93 (s, 1H), 12.29 (bs, 1H); MS(ES⁺): m/z 198 (M+H)⁺; (ES⁻): m/z 196 (M-H)⁻;
- o) 3-Formyl-5-trifluoromethoxy-1H-indole, ^1H NMR (300MHz, d6-DMSO): 7.24 (m, 1H), 7.61 (m, 1H), 7.97 (bs, 1H), 8.42 (d, 1H), 9.95 (s, 1H), 12.35 (bs, 1H); MS(ES⁺): m/z 230 (M+H)⁺; (ES⁻): m/z 228 (M-H)⁻; Analysis for C₁₀H₆F₃NO₂: Calcd.; C, 52.4138; H, 2.6391; N, 6.1122; found; C, 52.70; H, 2.73; N, 6.13;
- p) 3-Formyl-4,6-difluoro-5-methoxy-1H-indole, MS(ES⁺): 212 (M+H)⁺; (ES⁻): 210 (M-H)⁻;
- q) 3-Formyl-4-phenyl-1H-indole, ^1H NMR (300MHz, d6-DMSO): 7.07 (m, 1H), 7.30 (m, 1H), 7.46 (m, 6H), 7.53 (m, 1H), 8.20 (bs, 1H), 9.37 (s, 1H), 12.40 (bs, 1H). MS(ES⁺): m/z 222 (M+H)⁺; (ES⁻): m/z 220 (M-H)⁻;
- r) 3-Formyl-6-phenyl-1H-indole, ^1H NMR (300MHz, d6-DMSO): 7.35 (m, 1H), 7.49 (m, 3H), 7.71 (m, 3H), 8.15 (m, 1H), 8.33 (d, 1H), 9.96 (s, 1H), 12.20 (bs, 1H). MS(EI⁺): m/z 221 (M)⁺;
- s) 3-Formyl-5-isopropyl-1H-indole, ^1H NMR (300MHz, d6-DMSO): 1.24 (d, 6H), 2.99 (m, 1H), 7.15 (m, 1H), 7.41 (m, 1H), 7.94 (m, 1H), 8.22 (m, 1H), 9.90 (s, 1H), 12.02 (bs, 1H); MS(ES⁺): 188 (M+H)⁺; (ES⁻): m/z 186 (M-H)⁻;
- t) 3-Formyl-4,6-difluoro-5-methoxy-1-methyl-1H-indole: ^1H NMR (300MHz, CDCl₃): 3.81 (s, 3H), 4.02 (s, 3H), 6.92 (m, 1H), 7.77 (s, 1H), 10.14 (d, 1H); MS(ES⁺): m/z 226 (M+H)⁺; and
- u) 3-Formyl-4,6-difluoro-1-methyl-1H-indole: ^1H NMR (300MHz, d6-DMSO): 3.87 (s, 3H), 7.10 (m, 1H), 7.41 (m, 1H), 8.32 (s, 1H), 9.93 (d, 1H); MS(ES⁺): 196 (M+H)⁺.

Example 51

3-(2-Nitrovinyl)-4-methoxy-1H-indole

- Combine ammonium acetate (dried from treatment with toluene and removal of the toluene *in vacuo*) as a solid (0.75 eq.), nitromethane (20 eq.), and 4-methoxy-1H-indole-3-carbaldehyde (5.4 g; 30.82 mmol). Heat to about 65°C. After the reaction is judged to be near completion (by TLC), add silica gel and evaporate *in vacuo* to remove

-52-

the nitromethane. Load the silica gel from the reaction mixture on top of short column of silica gel and elute with 25% acetone in hexanes to give, after evaporation, the title compound which may be used in the next step without further purification.

- By the method of Example 51 the following compounds were prepared: a) 3-(2-Nitrovinyl)-6-methoxy-1H-indole;
- 5 b) 3-(2-Nitrovinyl)-7-methoxy-1H-indole, ^1H NMR (300 MHz; d₆-DMSO): 3.95 (s, 3H), 5.02 (m, 1H), 6.86 (d, 1H), 7.17 (t, 1H), 7.50 (d, 1H), 8.38 (d, 1H), 12.40 (bs, 1H); MS (ES⁺): m/z 219 (M+H)⁺; (ES⁻): m/z 217 (M-H)⁻;
- 10 c) 3-(2-Nitrovinyl)-4-chloro-1H-indole, ^1H NMR (300 MHz, d₆-DMSO): 5.08 (m, 1H), 7.24 (m, 2H), 7.51 (dd, 1H), 8.12 (d, 1H), 8.92 (d, 1H), 12.6 (bs, 1H); MS(ES⁻): m/z 221, 223 (M-H)⁻;
- d) 3-(2-Nitrovinyl)-6-chloro-1H-indole, ^1H NMR (300MHz, d₆-DMSO): 5.03 (m, 1H), 7.22 (dd, 1H), 7.58 (d, 1H), 8.03 (m, 2H), 8.38 (d, 1H), 12.23 (bs, 1H); MS (ES⁻): m/z 223, 221 (M-H)⁻;
- 15 e) 3-(2-Nitrovinyl)-7-chloro-1H-indole, ^1H NMR (300MHz, d₆-DMSO): 7.23 (t, 1H), 7.36 (d, 1H), 7.97 (d, 1H), 8.06 (d, 1H), 8.33 (bs, 1H), 8.40 (d, 1H), 12.58 (bs, 1H); MS (ES⁺): m/z 225, 223 (M+H)⁺; (ES⁻): m/z 223, 221 (M-H)⁻;
- f) 3-(2-Nitrovinyl)-4-fluoro-1H-indole,
- g) 3-(2-Nitrovinyl)-5-methoxy-6-trifluoromethyl-1H-indole, MS (ES⁺): m/z 286
- 20 (M⁺); (ES⁻): m/z 285 (M-H)⁻;
- h) 3-(2-Nitrovinyl)-6-chloro-5-methoxy-1H-indole,
- i) 3-(2-Nitrovinyl)-4-chloro-5-methoxy-1H-indole, ^1H NMR (300MHz, d₆-DMSO): 3.88 (s, 3H), 5.03 (m, 2H), 7.13 (d, 1H), 7.46 (d, 1H), 8.08 (d, 1H), 12.42 (bs, 1H); MS(ES⁻): m/z 151, 153 (M-H)⁻;
- 25 j) 3-(2-Nitrovinyl)-6-trifluoromethyl-1H-indole, MS(ES⁺): m/z 257 (M+H)⁺; (ES⁻): m/z 255 (M-H)⁻;
- k) 3-(2-Nitrovinyl)-5-methoxy-2-methyl-1H-indole, ^1H NMR (300MHz, d₆-DMSO): 2.58 (s, 3H), 3.84 (s, 3H), 6.82 (m, 1H), 7.28 (m, 2H), 7.89 (d, 1H), 8.29 (d, 1H), 12.14 (bs, 1H); MS(ES⁺): m/z 233 (M+H)⁺; (ES⁻): m/z 231 (M-H)⁻;
- 30 l) 3-(2-Nitrovinyl)-6-fluoro-5-methoxy-1H-indole;
- m) 3-(2-Nitrovinyl)-5,6-difluoro-1H-indole;
- n) 3-(2-Nitrovinyl)-6-chloro-5-fluoro-1H-indole;
- o) 3-(2-Nitrovinyl)-5-trifluoromethoxy-1H-indole;
- p) 3-(2-Nitrovinyl)-4,6-difluoro-5-methoxy-1H-indole;
- 35 q) 3-(2-Nitrovinyl)-4-phenyl-1H-indole;
- r) 3-(2-Nitrovinyl)-6-phenyl-1H-indole;
- s) 3-(2-Nitrovinyl)-5-isopropyl-1H-indole;

-53-

t) 3-(2-Nitrovinyl)-4,6-difluoro-5-methoxy-1-methyl-1H-indole: ^1H NMR (300MHz, $\text{d}_6\text{-DMSO}$): 3.82 (t, 3H), 3.92 (s, 3H), 7.53 (m, 1H), 7.84 (m, 1H), 8.30 (m, 2H); MS(ES $^+$): m/z 269 (M+H) $^+$; and

u) 3-(2-Nitrovinyl)-4,6-difluoro-1-methyl-1H-indole.

5

Example 52

4-Methoxytryptamine

Combine LiAlH_4 (6.78 g; 178.74 mmol; 6 eq) and anhydrous THF. Dissolve 3-(2-nitrovinyl)-4-methoxy-1H-indole (6.5 g; 29.79 mmol) in anhydrous THF and add dropwise to the LiAlH_4 solution with vigorous stirring. After the addition is complete, heat to reflux. After 1 hour cool to ambient temperature and stir. After 16 hours, quench the excess LiAlH_4 as described in *J. Med. Chem.* 1995, 38, 2050. Filter the gray suspension through celite and rinse the celite with ethyl acetate. Evaporate the filtrate *in vacuo* to residue. Chromatograph the residue on silica gel eluting with 1 L of $\text{CHCl}_3/\text{MeOH}/\text{NH}_4\text{OH}$ (95:10:1) and then 1 L of $\text{CHCl}_3/\text{MeOH}/\text{NH}_4\text{OH}$ (90:10:1) as the mobile phase. Pool fractions containing the product and evaporate to give the title compound as a tan solid: ^1H NMR (300 MHz, $\text{d}_6\text{-DMSO}$): 2.96 (t, 2H); 3.42 (t, 2H); 3.83 (s, 3H); 6.42 (dd, 1H); 6.93 (m, 3H); 10.7 (s, 1H); MS (ES $^+$): m/z 191 (M+H) $^+$; 174 (M-NH $_2$) $^+$; 159 (M-OCH $_3$) $^+$; (ES $^-$): m/z 189 (M-H) $^-$.

By the method of Example 52 the following compounds were prepared: a) 6-Methoxytryptamine, ^1H NMR (300 MHz; $\text{d}_6\text{-DMSO}$): 2.86 (t, 2H); 3.42 (t, 2H); 3.75 (s, 3H); 6.62 (dd, 1H); 6.83 (m, 1H); 6.97 (bs, 1H); 7.37 (m, 1H); 10.55 (s, 1H); MS (ES $^+$): m/z 191 (M+H) $^+$; 174 (M-NH $_2$) $^+$; (ES $^-$): m/z 189 (M-H) $^-$;

b) 7-Methoxytryptamine, ^1H NMR (300 MHz, $\text{d}_6\text{-DMSO}$): 2.88 (t, 2H), 3.42 (t, 2H), 3.89 (s, 3H), 6.61 (d, 1H), 6.89 (t, 1H), 7.02 (m, 1H), 7.10 (d, 1H), 10.85 (bs, 1H); MS (ES $^+$): m/z 191 (M+H) $^+$, 174 (M-NH $_2$) $^+$; (ES $^-$): m/z 189 (M-H) $^-$;

c) 4-Chlorotryptamine, ^1H NMR (300MHz, $\text{d}_6\text{-DMSO}$): 3.11 (t, 2H), 3.44 (t, 2H), 6.99 (m, 2H), 7.22 (m, 1H), 7.30 (d, 1H), 11.19 (bs, 1H); MS(ES $^+$): m/z 178, 180 (M+H) $^+$; (ES $^-$): m/z 193 (M-H) $^-$;

d) 6-Chlorotryptamine, ^1H NMR (300MHz, $\text{d}_6\text{-DMSO}$): 2.89 (t, 2H), 3.42 (t, 2H), 6.96 (dd, 1H), 7.17 (bs, 1H), 7.35 (m, 1H), 7.52 (d, 1H), 10.91 (bs, 1H); MS (ES $^+$): m/z 197, 195 (M+H) $^+$, 180, 178 (M-NH $_2$) $^+$; (ES $^-$): m/z 195, 193 (M-H) $^-$;

e) 7-Chlorotryptamine, ^1H NMR (300MHz, $\text{d}_6\text{-DMSO}$): 2.91 (t, 2H), 3.43 (t, 2H), 6.98 (t, 1H), 7.13 (d, 1H), 7.20 (bs, 1H), 7.51 (d, 1H), 11.15 (bs, 1H); MS (ES $^+$): m/z 197, 195 (M+H) $^+$, 180, 178 (M-NH $_2$) $^+$; (ES $^-$): m/z 195, 193 (M-H) $^-$;

f) 4-Fluorotryptamine,

g) 5-Methoxy-6-trifluoromethyltryptamine,

-54-

- h) 6-Chloro-5-methoxytryptamine, ^1H NMR (300MHz, $\text{d}_6\text{-DMSO}$): 2.89 (t, 2H), 3.42 (t, 2H), 3.84 (s, 3H), 7.12 (bs, 1H), 7.19 (s, 1H), 7.36 (s, 1H), 8.01 (bs, 1H); MS (ES $^+$): m/z 225, 227 (M+H) $^+$, 208, 210 (M-NH $_2$) $^+$; (ES $^-$): m/z 223, 225 (M-H) $^-$;
- i) 4-Chloro-5-methoxytryptamine, ^1H NMR (300MHz, $\text{d}_6\text{-DMSO}$): 3.10 (t, 2H), 3.43 (t, 2H), 3.81 (s, 3H), 6.95 (d, 1H), 7.18 (m, 1H), 7.25 (dd, 1H), 10.93 (bs, 1H); MS(ES $^+$): m/z 208, 210 (M-NH $_2$) $^+$ (ES $^-$): m/z 223, 225 (M-H) $^-$;
- j) 6-Trifluoromethyltryptamine,
- k) 5-Methoxy-2-methyltryptamine, ^1H NMR (300MHz, $\text{d}_6\text{-DMSO}$): 2.28 (s, 3H), 2.80 (t, 2H), 3.31 (bt, 2H), 6.59 (dd, 1H), 6.88 (d, 1H), 7.09 (d, 1H); MS(ES $^+$): m/z 188 (M-NH $_2$) $^+$ (ES $^-$): m/z 203 (M-H) $^-$;
- l) 6-Fluoro-5-methoxytryptamine;
- m) 5,6-Difluorotryptamine;
- n) 6-Chloro-5-fluorotryptamine;
- o) 5-Trifluoromethoxytryptamine;
- p) 4,6-Difluoro-5-methoxytryptamine;
- q) 4-Phenyltryptamine;
- r) 6-Phenyltryptamine;
- s) 5-Isopropyltryptamine;
- t) 4,6-Difluoro-5-methoxy-1-methyltryptamine: ^1H NMR (300MHz, CDCl_3): 3.0 (m, 4H), 3.67 (s, 3H), 3.98 (s, 3H), 6.85 (m, 2H); and
- u) 4,6-Difluoro-5-methoxy-1-methyltryptamine: ^1H NMR(300MHz, $\text{d}_6\text{-DMSO}$): 2.92 (t, 2H), 3.39 (t, 2H), 3.69 (s, 3H), 6.75 (m, 1H), 7.13 (m, 2H); MS(ES $^+$): m/z 211; (M+H) $^+$ 194 (M-NH $_2$) $^+$.

25

Example 534-Methoxytryptamine hydrochloride

- Dissolve 4-methoxytryptamine (1 g, 5.26 mmol) in MeOH and add of NH_4Cl (0.97 eq, 0.27 g, 5.10 mmol). After 30 minutes, evaporate *in vacuo* to remove the MeOH leaving a thick orange oil. Dissolve the oil in MeOH and add dropwise to Et_2O (200 mL) with vigorous stirring giving a gummy white precipitate. Stir with heating to give the title compound as a solid: ^1H NMR ($\text{d}_6\text{-DMSO}$, 300 MHz): 3.06 (bs, 4 H); 3.86 (s, 3H); 6.46 (dd, 1H); 7.06-6.9 (m, 3H); 7.93 (bs, 1H); 10.9 (s, 1H); MS (ES $^+$): m/z 191 (M+H) $^+$; 175 (M-CH $_3$) $^+$; 174 (M-NH $_2$) $^+$; (ES $^-$): 189 (M-H) $^-$; Analysis for $\text{C}_{11}\text{H}_{15}\text{ClN}_2\text{O}$: Calcd.: C, 58.2788; H, 6.6692; N, 12.3566; found C, 58.18; H, 6.73; N, 12.15.

35

-55-

Example 545,6-Difluorotryptamine hydrochloride

Prepare by the method of Example 53 to give the title compound: ^1H NMR (300MHz, $\text{d}_6\text{-DMSO}$): 2.97 (m, 4H), 7.27 (m, 1H), 7.36 (m, 1H), 7.53 (m, 1H), 11.20 (bs, 1H); MS(ES $^+$): m/z 197 (M+H) $^+$, 180 (M-NH $_2$) $^+$; (ES $^-$): m/z 195 (M-H) $^-$.

Example 554-Phenyltryptamine hydrochloride

Add a solution of HCl (4.6 mL of 4 M HCl in 1,4-dioxane) dropwise to a solution of 4-phenyltryptamine (3.33 g, 14.09 mmol) in EtOAc/Et $_2$ O to give a solid. Collect the solid by filtration and dry overnight in a vacuum oven at room temperature to give the title compound as an off white solid: ^1H NMR (300MHz, $\text{d}_6\text{-DMSO}$): 2.54 (m, 4H), 6.82 (m, 1H), 7.14 (t, 1H), 7.27 (m, 1H), 7.41 (m, 5H), 7.68 (bs, 2H), 11.28 (bs, 1H); MS(ES $^+$): m/z 237 (M+H) $^+$, 220 (M-NH $_2$) $^+$; (ES $^-$): m/z 235 (M-H) $^-$; Analysis for C $_{16}$ H $_{17}$ ClN $_2$; Calcd.: C, 70.4517; H, 6.2818; N, 10.2696; found; C, 70.26; H, 6.16; N, 10.20.

Example 565-Chloro-6-fluorotryptamine hydrochloride

Prepare by the method of Example 55 to give the title compound: ^1H NMR (300MHz, $\text{d}_6\text{-DMSO}$): 3.00 (m, 4H), 7.37 (m, 1H), 7.53 (d, 1H), 7.59 (d, 1H), 11.28 (bs, 1H); MS(ES $^+$): m/z 213 (M+H) $^+$, 196, 198 (M-NH $_2$) $^+$; (ES $^-$): m/z 211, 213 (M-H) $^-$.

Example 574-Chlorotryptamine oxalate

Add dropwise oxalic acid (1.32 g, 1.3 eq.) in MeOH to a solution of 4-chlorotryptamine in EtOAc (2.2 g, 11.3 mmol) with vigorous stirring. When addition was complete, add Et $_2$ O to the cloud point and place flask in the freezer to give a solid. Collect the solid by filtration and wash with ether. Dry in a vacuum oven at room temperature to give the title compound as an off-white solid: ^1H NMR (300MHz, $\text{d}_6\text{-DMSO}$): 3.11 (m, 2H), 3.2 (m, 2H), 7.04 (m, 2H), 7.34 (m, 2H), 11.44 (bs, 1H); MS (ES $^+$): m/z 195 (M+H) $^+$, 178 (M-NH $_2$) $^+$; (ES $^-$): m/z 193 (M-H) $^-$; Analysis for C $_{12}$ H $_{13}$ ClN $_2$ O $_4$; Calcd.: C, 50.6263; H, 4.6026; N, 9.8396; found: C, 50.56; H, 4.57; N, 9.66.

Using the method of Example 57 gives the following compounds: a) 6-Phenyltryptamine oxalate: 3.05 (m, 4H), 7.31 (m, 3H), 7.45 (t, 2H), 7.65 (m, 4H), 11.10 (bs, 1H). MS(ES $^+$): m/z 237 (M+H) $^+$, 220 (M-NH $_2$) $^+$; (ES $^-$): m/z 235 (M-H) $^-$;

-56-

b) 4,6-Difluoro-5-methoxytryptamine oxalate: ^1H NMR (300MHz, $\text{d}_6\text{-DMSO}$): 3.04 (m, 4H), 3.85 (s, 3H), 7.10 (m, 1H), 7.22 (m, 1H), 11.29 (bs, 1H); MS(ES⁺): m/z 227 (M+H)⁺; (ES⁻): m/z 225 (M-H)⁻; Analysis for $\text{C}_{13}\text{H}_{14}\text{F}_2\text{N}_2\text{O}_5$: Calcd.; C, 49.3718; H, 4.4620; N, 8.8576; found; C, 49.68; H, 4.57; N, 8.60; and

- 5 c) 5-Isopropyltryptamine oxalate: ^1H NMR (300MHz, $\text{d}_6\text{-DMSO}$): 1.25 (d, 6H), 3.01 (m, 4H), 6.99 (m, 1H), 7.17 (m, 1H), 7.27 (m, 1H), 7.36 (bs, 1H), 10.85 (bs, 1H); MS(ES⁺): m/z 203 (M+H)⁺, 186 (M-NH₂)⁺; (ES⁻): m/z 201 (M-H)⁻.

Example 58

10 5-Trifluoromethoxytryptamine oxalate

Add oxalic acid (1.3 eq.) in acetone to a solution of 5-trifluoromethoxytryptamine in acetone. Warm and add Et₂O to the cloud point and then placed in the freezer overnight to obtain the title compound as a white crystalline solid: ^1H NMR (300MHz, $\text{d}_6\text{-DMSO}$): 3.02 (m, 4H), 7.06 (m, 1H), 7.39 (m, 1H), 7.45 (d, 1H), 7.55 (m, 1H), 11.30 (bs, 1H). MS(ES⁺): m/z 245 (M+H)⁺, 228 (M-NH₂)⁺; (ES⁻): m/z 243 (M-H)⁻; Analysis for $\text{C}_{11}\text{H}_{11}\text{F}_3\text{N}_2\text{O}$: Calcd.; C, 46.7144; H, 3.9203; N, 8.3809; found; C, 46.55; H, 3.62; N, 8.27.

- Using the method of Example 58 gives the following compounds: a) 4,6-Difluoro-5-methoxytryptamine oxalate: ^1H NMR (300MHz, $\text{d}_6\text{-DMSO}$): 3.04 (m, 4H), 3.85 (s, 3H), 7.10 (m, 1H), 7.22 (m, 1H), 11.29 (bs, 1H); MS(ES⁺): m/z 227 (M+H)⁺; (ES⁻): m/z 225 (M-H)⁻; Analysis for $\text{C}_{13}\text{H}_{14}\text{F}_2\text{N}_2\text{O}_5$: Calcd.; C, 49.3718; H, 4.4620; N, 8.8576; found; C, 49.68; H, 4.57; N, 8.60.

Example 60

25 4-Fluorotryptamine oxalate

Add dropwise oxalic acid (1.44 g, 1.2 eq.) in acetonitrile to an acetonitrile solution of 4-fluorotryptamine with vigorous stirring. Warm add MeOH to make a solution. Add Et₂O to the cloud point and cool the solution in the freezer to give a solid. Collect the solid by filtration and dry in a vacuum oven overnight at 45°C to give the title compound as a tan solid: ^1H NMR (300MHz, $\text{d}_6\text{-DMSO}$): 3.07 (m, 4H), 6.73 (m, 1H), 7.04 (m, 1H), 7.22 (m, 2H), 11.30 (bs, 1H); MS (ES⁺): m/z 179 (M+H)⁺; (ES⁻): m/z 177 (M-H)⁻.

Example 61

35 6-Fluoro-5-methoxytryptamine oxalate

Add oxalic acid (3.91 g, 1.2 eq.) in MeOH dropwise to an EtOAc/MeOH solution of 6-fluoro-5-methoxytryptamine with vigorous stirring. Add Et₂O to give a solid and collect the solid by filtration and dry overnight in a vacuum oven at 60°C to give the title

-57-

compound: ^1H NMR (300MHz, d6-DMSO): 3.0 (m, 4H), 3.85 (s, 3H), 7.21 (m, 3H), 10.89 (bs, 1H); MS(ES⁺): m/z 209 (M+H)⁺; Analysis for C₁₃H₁₅FN₂O₅: Calcd.: C, 52.3496; H, 5.0690; N, 9.3919; Found: C, 52.06; H, 4.91; N, 9.20.

5

Example 622-(2-(7-Fluoro-1H-indol-3-yl)ethyl)isoindole-1,3-dione

Combine 2-fluorohydrazine hydrochloride (3.25 g, 20 mmol) and 2-(4,4-diethoxybutyl)-isoindole-1,3-dione (6.99 g, 24 mmol) and dissolve in 4% aqueous H₂SO₄. Heat the reaction to reflux. After 2 hours, cool to ambient temperature. Basify the reaction mixture with 30% aqueous NH₄OH to pH of about 11. Extract with dichloromethane (2x100 mL). Combine the organic phases, dry over MgSO₄, filter, and remove the solvent leaving an orange oil. Absorb the oil onto silica gel and load on top short column of silica gel equilibrated with 15% EtOAc in hexanes. Eluent with 15% EtOAc in hexanes (1500 mL) and then 30% EtOAc in hexanes (2000 mL) to give, after evaporation, the title compound as a yellow solid: ^1H NMR (300MHz, d6-DMSO): 3.03 (t, 2H), 3.85 (t, 2H), 6.91 (m, 2H), 7.25 (m, 1H), 7.36 (d, 1H), 7.83 (m, 4H), 11.32 (bs, 1H); MS (FD): m/z 308 (M⁺).

20

Example 637-Fluorotryptamine oxalate

Dissolve 2-(2-(7-fluoro-1H-indol-3-yl)ethyl)isoindole-1,3-dione in 25 mL of THF. Add ethanolamine (63.4 g, 62.65 mL, 1038 mmol, 100 eq.) with vigorous stirring and heat to 70°C. After 1.5 hours, cool to room temperature. After 18 hours, pour reaction mixture into of water (250 mL) containing of 5N NaOH (3 mL) and extract with Et₂O (2x200 mL). Combine the organic layers and wash with 0.1 N NaOH. Collect the organic layer, dry over MgSO₄, filter, and remove the solvent *in vacuo* to give the title compound as a yellow oil.

Add oxalic acid (0.62 g, 1.2 eq.) in MeOH dropwise to an EtOAc solution of the base (1.02 g, 5.72 mmol) with vigorous stirring. Heat the cloudy suspension to reflux for 30 minutes and then cool to give a solid. Collect the solid by filtration and dry overnight in a vacuum oven at 60°C to give the title compound as an off-white solid: ^1H NMR (300MHz, d6-DMSO): 3.04 (m, 4H), 6.96 (m, 2H), 7.30 (m, 1H), 7.38 (d, 1H), 11.51 (bs, 1H); MS (ES⁺): m/z 179 (M+H)⁺, 162 (M-NH₂)⁺; (ES⁻): m/z 177 (M-H)⁻; Analysis for C₁₂H₁₃FN₂O₄: Calcd.: C 53.7318; H 4.8849; N 10.4431; found: C, 53.50; H, 4.86; N, 10.32.

-58-

Example 646-Trifluoromethyltryptamine oxalate

Add 6-trifluoromethyltryptamine to 1:1 acetone/Et₂O. Add dropwise of oxalic acid (1.2 eq.) in acetone to give a solid. Collect the solid by filtration and dry overnight in the vacuum oven to obtain the title compound: MS(ES⁺): m/z 212 (M-NH₂)⁺; (ES⁻): m/z 227 (M-H)⁻.

Example 654,6-Difluoro-5-methoxy-1H-indole

Dissolve 2,6-difluoro-4-nitrophenol (*J. Heterocyclic. Chem.* 1976, 13, 1253; 10g, 57.11 mmol) in 300 mL of benzene. Add dropwise a solution of 1-methyl-3-p-tolyltriazene (9.37 g, 62.82 mmol, 1.1 eq.) in benzene (150 mL). After TLC indicates absence of starting material, transfer the reaction mixture to a separator funnel and wash with 1 N HCl, then saturated NaHCO₃, and then water. Dry the organic layer over MgSO₄, filter, and remove the solvent to give a residue. Crystallize the residue from MeOH/water to give 1,3-difluoro-2-methoxy-5-nitrobenzene as white needles: ¹H NMR (300MHz, CDCl₃): 4.25 (t, 3H), 7.80 (d, 2H).

Combine 1,3-difluoro-2-methoxy-5-nitrobenzene (10.12 g, 53.51 mmol) and 4-chlorophenoxyacetonitrile (11.21 g, 66.89 mmol, 1.25 eq.) in DMSO (150 mL). Add dropwise to a suspension of solid NaOH (powdered, 10.70 g, 267.55 mmol, 5 eq.) over 5 hours. After 18 hours, pour the reaction mixture into cold, aqueous HCl and extract with Et₂O (2x150 mL). Combine the organic layers, wash with brine, and evaporate to give a residue. Chromatograph the residue on silica gel and eluting with 20% EtOAc in hexanes to give, after evaporation, (2,4-difluoro-3-methoxy-6-nitrophenyl)acetonitrile as a yellow oil: MS(ES⁻): m/z 227 (M-H)⁻.

Using (2,4-difluoro-3-methoxy-6-nitrophenyl)acetonitrile in the cyclization as described in *Israel J. Chem.* 1966, 4, 155-159 gives an oil. Chromatograph the oil on silica gel eluting with 20% EtOAc in hexanes to give, after evaporation, the title compound as a purple solid; ¹H NMR (300MHz, d₆-DMSO): 3.85 (bs, 3H), 6.46 (m, 1H), 7.12 (d, 1H), 7.36 (m, 1H), 11.35 (bs, 1H); MS(ES⁻): m/z 182 (M-H)⁻.


Using the method of Example 65 gives the following compounds: a) 4,6-Difluoro-1H-indole: ¹H NMR (300MHz, CDCl₃): 4.68 (d, 2H), 6.14 (m, 2H), 6.57 (bs, 2H); MS(ES⁺): m/z 205, 207 (M+H)⁺ which gives the title compound.

-59-

Example 664,6-Difluoro-5-methoxy-1-methyl-1H-indole

- Combine 4,6-difluoro-5-methoxy-1H-indole (7.5 g, 40.95 mmol) and cold DMF (100 mL) and treat with NaH (1.8 g, 45.05 mmol, 1.1 eq.) and with vigorously stirring.
- 5 After about 10 minutes, add dropwise iodomethane (11.62 g, 81.90 mmol, 2 eq.). After the addition was complete, allow the reaction to stir at room temperature for several hours until TLC indicates the absence of starting material. Dilute the reaction with water and extract with Et₂O (2x150 mL). Combine the organic layers, dry over MgSO₄, filter, and remove the solvent leaving an oil. Chromatograph the oil on silica gel eluting with 10% EtOAc in hexanes to give, after evaporation, the title compound as a light yellow oil: ¹H NMR (300MHz, CDCl₃): 3.72 (s, 3H), 3.97 (s, 3H), 6.50 (d, 1H), 6.84 (d, 1H), 6.98 (d, 1H); MS(ES⁺): m/z 198 (M+H)⁺; Analysis for C₁₀H₉F₂NO: Calcd.: C, 60.91; H, 4.60; N, 7.10; found: C, 60.93; H, 4.63; N, 7.25.
- 10

- Using the method of Example 66 gives the following compounds: a) 4,6-Difluoro-1-methyl-1H-indole.
- 15

Example 67N-(2-(5-Methoxy-1H-indol-3-yl)ethyl)-3-phenoxybenzylamine


- Combine 3-phenoxybenzaldehyde (5.6 ml, 26.7 mmol), 5-methoxytryptamine (5.0 g, 26.7 mmol) and 3 Å molecular sieves (1.0 g) in methanol (50 ml) and under argon and heat at reflux for 4 hours. Remove the molecular sieves by filtration and then slowly add sodium borohydride (3.0 g, 60.0 mmol) portionwise. Stir at room temperature for 1 hour and concentrate under reduced pressure, dissolve the concentrated reaction mixture in sodium hydroxide 1N (100 ml) and extract with dichloromethane (3 x 50 ml). Combine organic layers and wash sequentially with distilled water (50 ml) and brine (50 ml), dry (Na₂SO₄) the organic layer, and concentrate to give a residue. Chromatograph the residue on silica gel eluting with 9:1 EtOAc:MeOH with 2% NH₄OH gives the title compound.
- 20
- 25

- Formation of oxalate salt: Add a solution of the free base (8.7 g, 23.5 mmol) in EtOAc (50 ml) to a solution of oxalic acid (2.1 g, 23.5 mmol) in EtOAc (5 ml) to give a precipitate. Collect the precipitate and recrystallize from methanol/diethyl ether to give a solid. Collect the solid by filtration, rinse with diethyl ether, and dry in a vacuum oven at 50°C overnight to give the title compound as the oxalate, mp 188-190°C, RMN
- 30

-60-

consistent, Mass: m/z 373.2 (M^+), Anal. Calcd for $C_{26}H_{26}N_2O_6$: C, 67.52; H, 5.67; N, 6.06. Found: C, 67.38; H, 5.46; N, 6.04.

By the method of Example 67 the following compounds were prepared, isolated as the oxalate except where noted:



5

No.	Z'	X	R ₄	Data
68	H	-O-	phenyl	mp 203-205°C, Mass: m/z 343.1 (M^+), Anal. Calcd for $C_{25}H_{24}N_2O_5$: C, 69.43; H, 5.59; N, 6.48 Found: C, 69.25; H, 5.42; N, 6.37
69	H	-S-	phenyl	mp 106-108°C, Mass: m/z 359.2 (M^+), Anal. Calcd for $C_{22}H_{24}N_2O_4S$: C, 66.95; H, 5.39; N, 6.25 Found: C, 66.19; H, 5.49; N, 6.13
70	H	-SO ₂ -	phenyl	mp 203-205°C, Mass: m/z 391.2 (M^+), Anal. Calcd for $C_{25}H_{24}N_2O_6S$: C, 62.49; H, 5.03; N, 5.83 Found: C, 62.05; H, 5.21; N, 5.82
71	5-methoxy	-S-	phenyl	mp 198-200°C, Mass: m/z 389.3 (M^+), Anal. Calcd for $C_{26}H_{26}N_2O_5S$: C, 65.25; H, 5.48; N, 5.85 Found: C, 64.50; H, 5.63; N, 5.73
72	5-methoxy	-SO ₂ -	phenyl	mp 142-144°C, Mass: m/z 421.1 (M^+), Anal. Calcd for $C_{26}H_{26}N_2O_7S$: C, 61.16; H, 5.13; N, 5.49 Found: C, 61.14; H, 5.38; N, 5.25
73	H	-S-	4-methylphenyl	mp 190-192°C, Mass: m/z 373.2 (M^+), Anal. Calcd for $C_{26}H_{26}N_2O_4S$: C, 67.51; H, 5.67; N, 6.06 Found: C, 67.44; H, 5.69; N, 6.13
74	H	-SO ₂ -	4-methylphenyl	mp 212-214°C, Mass: m/z 405.4 (M^+), Anal. Calcd for $C_{26}H_{26}N_2O_6S$: C, 63.14; H, 5.30; N, 5.66 Found: C, 62.59; H, 5.70; N, 5.29
75	5-methoxy	-CH(F)-	phenyl	mp 214-216°C, Mass: m/z 389.3 (M^+), Anal. Calcd for $C_{27}H_{27}FN_2O_5$: C, 67.77; H, 5.69; N, 5.85 Found: C, 67.52; H, 5.77; N, 5.64
76	H	-CH(F)-	phenyl	mp 216-218°C, Mass: m/z 359.2 (M^+), Anal. Calcd for $C_{26}H_{25}FN_2O_4$: C, 69.63; H, 5.62; N, 6.25 Found: C, 69.55; H, 5.36; N, 5.95

-61-

77	5-methoxy	-CH ₂ -	phenyl	mp 199-202°C, Mass: m/z 371.1 (M ⁺), Anal. Calcd for C ₂₇ H ₂₈ N ₂ O ₅ : C, 70.42; H, 6.13; N, 6.08 Found: C, 69.73; H, 6.25; N, 6.05
78	H	-CH ₂ -	phenyl	mp 222-224°C, Mass: m/z 341.2 (M ⁺), Anal. Calcd for C ₂₆ H ₂₆ N ₂ O ₄ : C, 72.54; H, 6.09; N, 6.51 Found: C, 72.23; H, 6.08; N, 6.37
79	5-methoxy	-CH(OH)-	phenyl	mp 146-148, Mass: m/z 387.2, Anal. Calcd for C ₂₇ H ₂₈ N ₂ O ₅ : C, 68.05; H, 5.92; N, 5.88 Found: C, 67.29; H, 6.03; N, 5.51
80	H	-CH(OH)-	phenyl	mp 167-169°C, Mass: m/z 357.3 (M ⁺), Anal. Calcd for C ₂₆ H ₂₆ N ₂ O ₅ : C, 69.94; H, 5.87; N, 6.27. Found: C, 68.11; H, 6.07; N, 6.06
81	5-methoxy	-NH-	phenyl	mp 170-172°C, Mass: m/z 372.3 (M ⁺), Anal. Calcd for C ₂₆ H ₂₇ N ₃ O ₅ : C, 67.67; H, 5.90; N, 9.10 Found: C, 67.24; H, 6.08; N, 8.54
82	H	-NH-	phenyl	mp 196-198°C, Mass: m/z 342.2 (M ⁺), Anal. Calcd for C ₂₅ H ₂₅ N ₃ O ₄ : C, 69.59; H, 5.84; N, 9.74 Found: C, 67.57; H, 6.06; N, 8.84
83	5-methoxy	-NH-	benzyl	mp 203-205°C, Mass: m/z 386.2 (M ⁺), Anal. Calcd for C ₂₇ H ₂₉ N ₃ O ₅ : C, 68.20; H, 6.15; N, 8.84 Found: C, 67.46; H, 6.14; N, 8.79
84	H	-NH-	benzyl	mp 204-206°C, Mass: m/z 356.3 (M ⁺), Anal. Calcd for C ₂₆ H ₂₇ N ₃ O ₄ : C, 70.10; H, 6.11; N, 9.43. Found: C, 68.48; H, 5.95; N, 9.26.
85	5-methoxy	-O-	H	mp 126-128°C, Mass: m/z 297.5 (M ⁺), Anal. Calcd for C ₁₈ H ₂₀ N ₂ O ₂ : C, 72.94; H, 6.80; N, 9.45. Found: C, 71.78; H, 6.71; N, 9.20. (isolated as the base)
86	H	-O-	H	mp 143-145°C, Mass: m/z 267.3 (M ⁺), Anal. Calcd for C ₁₇ H ₁₈ N ₂ O: C, 76.66; H, 6.81; N, 10.51. Found: C, 75.11; H, 6.61; N, 10.22 (isolated as the base)
87	5-fluoro	-O-	phenyl	mp 204-206°C, Mass: m/z 361.1 (M ⁺), Anal. Calcd for C ₂₅ H ₂₃ FN ₂ O ₅ : C, 66.66; H, 5.15; N, 6.22. Found: C, 66.83; H, 5.17; N, 6.30.

88	5-methoxy	-O-	naphth-1-yl	mp 196-198°C, Mass: m/z 423.1 (M^+), Anal. Calcd for $C_{30}H_{26}N_2O_6$: C, 70.30; H, 5.51; N, 5.47. Found: C, 68.11; H, 5.56; N, 5.52.
89	H	-O-	naphth-1-yl	mp 210-212°C, Mass: m/z 393.2 (M^+), Anal. Calcd for $C_{29}H_{24}N_2O_5$: C, 72.19; H, 5.43; N, 5.81. Found: C, 72.10; H, 5.40; N, 6.66.
90	5-methoxy	-O-	3-fluoro phenyl	mp 186-188°C, Mass: m/z 391.2 (M^+), Anal. Calcd for $C_{26}H_{25}FN_2O_6$: C, 64.99; H, 5.24; N, 5.83. Found: C, 63.10; H, 5.11; N, 5.67.
91	H	-O-	3-fluoro phenyl	mp 217-219°C in 75% yield, RMN consistent, Mass: m/z 361.1 (M^+), Anal. Calcd for $C_{25}H_{23}FN_2O_5$: C, 66.66; H, 5.15; N, 6.22. Found: C, 66.12; H, 5.22; N, 6.34.
92	5-methoxy	-O-	2-fluoro phenyl	mp 184-186°C, Mass: m/z 391.2 (M^+), Anal. Calcd for $C_{26}H_{25}FN_2O_6$: C, 64.99; H, 5.24; N, 5.83. Found: C, 65.06; H, 5.23; N, 5.85.
93	H	-O-	2-fluoro phenyl	mp 206-208°C, Mass: m/z 361.1 (M^+), Anal. Calcd for $C_{25}H_{23}FN_2O_5$: C, 66.66; H, 5.15; N, 6.22. Found: C, 66.30; H, 4.97; N, 6.21.
94	5-methoxy	-O-	4-fluoro phenyl	mp 184-186°C, Mass: m/z 391.2 (M^+), Anal. Calcd for $C_{26}H_{25}FN_2O_6$: C, 64.99; H, 5.24; N, 5.83. Found: C, 63.99; H, 4.95; N, 5.75.
95	H	-O-	4-fluoro phenyl	mp 222-224°C, Mass: m/z 361.1 (M^+), Anal. Calcd for $C_{25}H_{23}FN_2O_5$: C, 66.66; H, 5.15; N, 6.22. Found: C, 65.74; H, 4.81; N, 6.13.
96	5-methoxy	-O-	naphth-2-yl	mp 198-200°C, Mass: m/z 423.1 (M^+), Anal. Calcd for $C_{30}H_{26}N_2O_6$: C, 70.30; H, 5.51; N, 5.47. Found: C, 68.97; H, 5.43; N, 5.44.
97	H	-O-	naphth-2-yl	mp 219-221°C, Mass: m/z 393.2 (M^+), Anal. Calcd for $C_{29}H_{24}N_2O_5$: C, 72.19; H, 5.43; N, 5.81. Found: C, 71.65; H, 5.32; N, 5.91.
98	5-methoxy	-O-	benzyl	mp 204-206°C, Mass: m/z 387.2 (M^+), Anal. Calcd for $C_{27}H_{28}N_2O_6$: C, 68.05; H, 5.92; N, 5.87. Found: C, 67.26; H, 5.80; N, 5.86.

-63-

99	H	-O-	benzyl	mp 211-213°C, Mass: m/z 357.3 (M^+), Anal. Calcd for $C_{26}H_{26}N_2O_5$: C, 69.94; H, 5.86; N, 6.27. Found: C, 69.46; H, 5.75; N, 6.16.
100	5-hydroxy	-O-	phenyl	mp 188-190°C, Mass: m/z 359.2 (M^+), Anal. Calcd for $C_{25}H_{24}N_2O_6$: C, 66.95; H, 5.39; N, 6.24. Found: C, 63.56; H, 5.01; N, 5.86.
101	5-methoxy	-O-	pyrimid-5-yl	mp 191-193°C, Mass: m/z 375.2 (M^+), Anal. Calcd for $C_{24}H_{24}N_4O_6$: C, 62.06; H, 5.20; N, 12.06. Found: C, 61.66; H, 5.41; N, 10.87.
102	H	-O-	pyrimid-5-yl	mp 188-190°C, Mass: m/z 345.1 (M^+), Anal. Calcd for $C_{23}H_{22}N_4O_5$: C, 63.58; H, 5.10; N, 12.89. Found: C, 62.52; H, 5.28; N, 11.58.
103	5-methoxy	-O-	pyrid-4-yl	mp 124-126°C, Mass: m/z 374.2 (M^+), Anal. Calcd for $C_{23}H_{25}Cl_2N_3O_2$: C, 61.88; H, 5.64; N, 9.41. Found: C, 61.26; H, 5.70; N, 9.14. (isolated as the hydrochloride)
104	H	-O-	pyrid-4-yl	mp 147-149°C, Mass: m/z 344.2 (M^+), Anal. Calcd for $C_{22}H_{23}Cl_2N_3O$: C, 63.46; H, 5.56; N, 10.09. Found: C, 61.47; H, 5.33; N, 9.43. (isolated as the hydrochloride)
105	6-chloro	-O-	pyrid-4-yl	mp 150-152°C, Mass: m/z 378.2 (M^+), Anal. Calcd for $C_{22}H_{22}Cl_3N_3O$: C, 58.61; H, 4.91; N, 9.32. Found: C, 57.28; H, 4.61; N, 8.85.
106	5-methoxy	-O-	pyrid-3-yl	mp 178-180°C, Mass: m/z 374.2 (M^+), Anal. Calcd for $C_{25}H_{25}N_3O_6$: C, 64.78; H, 5.43; N, 9.06. Found: C, 63.02; H, 5.30; N, 8.87.
107	H	-O-	pyrid-3-yl	mp 190-192°C, Mass: m/z 344.1 (M^+), Anal. Calcd for $C_{24}H_{23}N_3O_5$: C, 66.50; H, 5.34; N, 9.69. Found: C, 65.69; H, 5.21; N, 9.20.
108	5-fluoro	-O-	pyrid-3-yl	mp 135-137°C, Mass: m/z 362.3 (M^+), Anal. Calcd for $C_{22}H_{22}Cl_2FN_3O$: C, 60.83; H, 5.10; N, 9.67. Found: C, 61.49; H, 5.31; N, 9.70. (isolated as the hydrochloride)

109	6-chloro	-O-	pyrid-3-yl	mp 160-162°C, Mass: m/z 378.1 (M^+), Anal. Calcd for $C_{22}H_{22}Cl_3N_3O$: C, 58.61; H, 4.91; N, 9.32. Found: C, 58.18; H, 4.89; N, 9.01. (isolated as the hydrochloride)
110	5-methoxy	-O-	pyrid-2-yl	mp 202-204°C, Mass: m/z 374.2 (M^+), Anal. Calcd for $C_{23}H_{25}Cl_2N_3O_2$: C, 61.88; H, 5.64; N, 9.41. Found: C, 60.57; H, 6.35; N, 10.89. (isolated as the hydrochloride)
111	H	-O-	pyrid-2-yl	mp 196-198°C, Mass: m/z 344.2 (M^+), Anal. Calcd for $C_{22}H_{23}Cl_2N_3O$: C, 63.46; H, 5.56; N, 10.09. Found: C, 63.69; H, 6.09; N, 11.62. (isolated as the hydrochloride)
112	6-chloro	-O-	pyrid-2-yl	mp 149-151°C, Mass: m/z 378.1 (M^+), Anal. Calcd for $C_{22}H_{22}Cl_3N_3O$: C, 58.61; H, 4.91; N, 9.32. Found: C, 61.96; H, 4.91; N, 9.73. (isolated as the hydrochloride)
113	5-methoxy	-O-	thiazol-2-yl	mp 180-182°C, Mass: m/z 380.3 (M^+), Anal. Calcd for $C_{23}H_{23}N_3O_6S$: C, 58.83; H, 4.93; N, 8.94. Found: C, 58.11; H, 4.79; N, 8.84.
114	H	-O-	thiazol-2-yl	mp 203-205°C, Mass: m/z 350.3 (M^+), Anal. Calcd for $C_{22}H_{21}N_3O_5S$: C, 60.12; H, 4.81; N, 9.56. Found: C, 59.73; H, 4.83; N, 9.36.
115	5-methoxy	-O-	2,6-difluoro phenylsulfon-yl	mp 137-139°C, Mass: m/z 473.1 (M^+), Anal. Calcd for $C_{26}H_{24}F_2N_2O_8S$: C, 55.51; H, 4.30; N, 4.97. Found: C, 55.90; H, 4.47; N, 5.12.
116	H	-O-	2,6-difluoro phenylsulfon-yl	mp 185-187°C, Mass: m/z 443.2 (M^+), Anal. Calcd for $C_{25}H_{22}F_2N_2O_7S$: C, 56.38; H, 4.16; N, 5.26. Found: C, 56.96; H, 4.39; N, 5.31.
117	5-methoxy	-NH-	pyrid-2-yl	mp 174-176°C, Mass: m/z 373.1 (M^+), Anal. Calcd for $C_{23}H_{26}Cl_2N_4O_6$: C, 62.02; H, 5.88; N, 12.57. Found: C, 61.45; H, 5.91; N, 12.22. (isolated as the hydrochloride)


-65-

118	H	-NH-	pyrid-2-yl	mp 168-170°C, Mass: m/z 343.1 (M^+), Anal. Calcd for $C_{22}H_{24}Cl_2N_4$: C, 63.61; H, 5.82; N, 13.48. Found: C, 62.18; H, 6.12; N, 12.11. (isolated as the hydrochloride)
119	6-chloro	-NH-	pyrid-2-yl	mp 164-166°C, Mass: m/z 377.1 (M^+), Anal. Calcd for $C_{22}H_{23}Cl_3N_4$: C, 58.74; H, 5.15; N, 12.45. Found: C, 57.75; H, 5.07; N, 11.94. (isolated as the hydrochloride)
120	5-methoxy	-NH-	pyrid-3-yl	mp 150-154°C, Mass: m/z 373.2 (M^+), Anal. Calcd for $C_{23}H_{26}Cl_2N_4O_6$: C, 62.02; H, 5.88; N, 12.57. Found: C, 61.30; H, 6.58; N, 10.87. (isolated as the hydrochloride)
121	H	-NH-	pyrid-3-yl	mp 140-142°C, Mass: m/z 343.2 (M^+), Anal. Calcd for $C_{22}H_{22}N_4$: C, 77.16; H, 6.47; N, 16.36. Found: C, 75.73; H, 6.54; N, 15.58. (isolated as the base)
122	6-chloro	-NH-	pyrid-3-yl	mp 172-174°C, Mass: m/z 377.2 (M^+), Anal. Calcd for $C_{22}H_{23}Cl_3N_4$: C, 58.74; H, 5.15; N, 12.45. Found: C, 57.05; H, 5.16; N, 11.84. (isolated as the hydrochloride)
123	5-methoxy	-NH-	pyrid-4-yl	mp 170-172°C, Mass: m/z 373.3 (M^+), Anal. Calcd for $C_{23}H_{26}Cl_2N_4O_6$: C, 62.02; H, 5.88; N, 12.57. Found: C, 61.05; H, 6.08; N, 11.97. (isolated as the hydrochloride)
124	H	-NH-	pyrid-4-yl	mp 174-176°C, Mass: m/z 343.4 (M^+), Anal. Calcd for $C_{22}H_{24}Cl_2N_4$: C, 63.61; H, 5.82; N, 13.48. Found: C, 62.32; H, 6.20; N, 12.44. (isolated as the hydrochloride)
125	6-chloro	-NH-	pyrid-4-yl	mp 158-160°C, Mass: m/z 377.2 (M^+), Anal. Calcd for $C_{22}H_{23}Cl_3N_4$: C, 58.74; H, 5.15; N, 12.45. Found: C, 57.17; H, 5.19; N, 11.69. (isolated as the hydrochloride)
126	5-methoxy-6-fluoro	-NH-	2,2,2-trifluoroethyl	mp 151-153°C, Mass: m/z 397.2 (M^+), Anal. Calcd for $C_{24}H_{24}F_4N_2O_6$: C, 56.25; H, 4.72; N, 5.46. Found: C, 56.38; H, 4.76; N, 5.53. (isolated as the maleate)

-66-

127	5-methoxy-6-fluoro	-NH-	2,2,3,3,3-pentafluoro propyl	mp 145-147°C, Mass: m/z 447.2 (M^+), Anal. Calcd for $C_{25}H_{24}F_6N_2O_6$: C, 53.38; H, 4.30; N, 4.98. Found: C, 53.36; H, 4.29; N, 5.00. (isolated as the maleate)
128	5-methoxy-6-fluoro	-O-	2,2,3,3-tetrafluoro propyl	mp 143-145°C, Mass: m/z 429.2 (M^+), Anal. Calcd for $C_{25}H_{25}F_5N_2O_6$: C, 55.14; H, 4.62; N, 5.14. Found: C, 55.10; H, 4.62; N, 5.18. (isolated as the maleate)
129	5-methoxy	-C(O)-	phenyl	mp 163-166°, Mass: m/z 385.2 (M^+), Anal. Calcd for $C_{27}H_{26}N_2O_6$: C, 68.34; H, 5.52; N, 5.90. Found: C, 66.64; H, 5.56; N, 5.90.
130	H	-C(O)-	phenyl	mp 168-170°C, Mass: m/z 355.3 (M^+), Anal. Calcd for $C_{26}H_{24}N_2O_5$: C, 70.26; H, 5.44; N, 6.30. Found: C, 69.51; H, 5.52; N, 6.22.
130 A	6-fluoro	-O-	pyrid-4-yl	mp: 123.4-124.9°C. Mass (ES ⁺): m/z 363.0 (M+1). Anal. Calcd. for $C_{22}H_{20}FN_3O$: C, 73.11; H, 5.58; N, 11.63. Found: C, 73.36; H, 5.41; N, 11.57. (isolated as the free base)
130 B	6-fluoro	-O-	pyrid-3-yl	mp 169.0-170.8°C. Mass (APCD): m/z 362.1 (M+1). Anal. Calcd for $C_{22}H_{20}F_1N_3O \cdot 1.0 C_4H_4O_4$: C, 65.40; H, 5.07; N, 8.80. Found: C, 65.45; H, 5.12; N, 8.70. (isolated as the maleate salt)
130 C	5-methoxy 6-fluoro	-O-	2,2,2 trifluoro ethyl	mp 151-153°C Mass: m/z 397.2 (M^+), Anal. Calcd for $C_{24}H_{24}F_4N_2O_6$: C, 56.25; H, 4.72; N, 5.46. Found: C, 56.38; H, 4.76; N, 5.53. (isolated as the maleate salt)

By the method of Example 67 the following compounds were prepared, isolated as the oxalate except where noted:



-67-

No.	Z'	X	R ₄	Data
131	3-chloro	-O-	phenyl	mp 222-224°C, Mass: m/z 338.2 (M ⁺), Anal. Calcd for C ₂₃ H ₂₂ ClNO ₅ : C, 64.56; H, 5.18; N, 3.27. Found: C, 64.24; H, 5.02; N, 3.89.
132	3-trifluoro methyl	-O-	phenyl	mp 220-222°C, Mass: m/z 372.2 (M ⁺), Anal. Calcd for C ₂₄ H ₂₂ F ₃ NO ₅ : C, 62.47; H, 4.81; N, 3.04. Found: C, 62.69; H, 4.78; N, 3.10.
133	4-methoxy	-O-	phenyl	mp 221-223°C, Mass: m/z 334.2 (M ⁺), Anal. Calcd for C ₂₄ H ₂₅ NO ₆ : C, 68.07; H, 5.95; N, 3.31. Found: C, 67.98; H, 5.92; N, 3.29.
134	3,4-dimethoxy	-O-	phenyl	Mp 209-211°C, Mass: m/z 364.2 (M ⁺), Anal. Calcd for C ₂₅ H ₂₇ NO ₇ : C, 66.21; H, 6.00; N, 3.09. Found: C, 66.28; H, 6.07; N, 3.27.
135	3-methoxy	-O-	phenyl	Mp 210-212°C, Mass: m/z 334.1 (M ⁺), Anal. Calcd for C ₂₄ H ₂₅ NO ₆ : C, 68.07; H, 5.95; N, 3.31. Found: C, 68.31; H, 5.78; N, 3.36.
136	3,4-dichloro	-O-	phenyl	mp 219-221°C, Mass: m/z 372.1 (M ⁺), Anal. Calcd for C ₂₃ H ₂₁ Cl ₂ NO ₅ : C, 59.75; H, 4.58; N, 3.03. Found: C, 58.98; H, 4.63; N, 3.66.
137	3-chloro	-O-	3-trifluoromethyl phenyl	mp 214-216°C, Mass: m/z 406.4 (M ⁺), Anal. Calcd for C ₂₄ H ₂₁ ClF ₃ NO ₅ : C, 58.13; H, 4.27; N, 2.82. Found: C, 58.28; H, 4.53; N, 2.86.
138	3-chloro	-O-	4-t-butyl phenyl	mp 221-223°C, Mass: m/z 394.2 (M ⁺), Anal. Calcd for C ₂₇ H ₃₀ ClNO ₅ : C, 67.00; H, 6.25; N, 2.89. Found: C, 66.36; H, 5.83; N, 2.94.
139	3-chloro	-O-	4-chloro phenyl	mp 212-214°C, Mass: m/z 372.1 (M ⁺), Anal. Calcd for C ₂₃ H ₂₁ Cl ₂ NO ₅ : C, 59.75; H, 4.58; N, 3.03. Found: C, 61.50; H, 4.77; N, 3.20.
140	3-chloro	-O-	4-methoxy phenyl	mp 207-209°C, Mass: m/z 368.2 (M ⁺), Anal. Calcd for C ₂₄ H ₂₄ ClNO ₆ : C, 62.95; H, 5.28; N, 3.06. Found: C, 63.17; H, 5.32; N, 3.19.

-68-

141	3-chloro	-O-	4-methyl phenyl	mp 206-208°C, Mass: m/z 352.4 (M ⁺), Anal. Calcd for C ₂₄ H ₂₄ ClNO ₅ : C, 65.23; H, 5.47; N, 3.17. Found: C, 67.52; H, 5.68; N, 3.30.
142	3-chloro	-O-	3,5-dichloro phenyl	mp 223-225°C, Mass: m/z 406.3 (M ⁺), Anal. Calcd for C ₂₃ H ₂₀ Cl ₃ NO ₅ : C, 55.61; H, 4.06; N, 2.82. Found: C, 56.08; H, 3.83; N, 2.26.
143	3-chloro	-O-	3,4-dichloro phenyl	mp 217-219°C, Mass: m/z 406.4 (M ⁺), Anal. Calcd for C ₂₃ H ₂₀ Cl ₃ NO ₅ : C, 55.61; H, 4.06; N, 2.82. Found: C, 55.73; H, 4.38; N, 3.02.
144	H	-O-	phenyl	mp 162-164°C, Mass: m/z 304.2 (M ⁺), Anal. Calcd for C ₂₃ H ₂₃ NO ₅ : C, 70.22; H, 5.89; N, 3.56. Found: C, 70.70; H, 5.38; N, 3.78.
145	4-chloro	-O-	phenyl	mp 222-224°C, Mass: m/z 338.2 (M ⁺), Anal. Calcd for C ₂₃ H ₂₂ ClNO ₅ : C, 64.56; H, 5.18; N, 3.27. Found: C, 63.65; H, 5.18; N, 3.25.
146	3-chloro	-S-	phenyl	mp 122-124°C, Mass: m/z 354.3 (M ⁺), Anal. Calcd for C ₂₃ H ₂₂ ClNO ₄ S: C, 62.23; H, 4.99; N, 3.15. Found: C, 63.08; H, 5.09; N, 3.15.
147	3-chloro	-SO ₂ -	phenyl	mp 110-112°C, Mass: m/z 386.1 (M ⁺), Anal. Calcd for C ₂₃ H ₂₂ ClNO ₆ S: C, 58.04; H, 4.66; N, 2.94. Found: C, 58.91; H, 4.78; N, 3.05.
148	H	-S-	phenyl	mp 111-113°C, Mass: m/z 320.1 (M ⁺), Anal. Calcd for C ₂₃ H ₂₃ NO ₄ S: C, 67.46; H, 5.66; N, 3.42. Found: C, 67.66; H, 5.77; N, 3.41.
149	H	-SO ₂ -	phenyl	mp 127-129°C, Mass: m/z 352.4 (M ⁺), Anal. Calcd for C ₂₃ H ₂₃ NO ₆ S: C, 62.57; H, 5.25; N, 3.17. Found: C, 62.75; H, 5.16; N, 3.26.
150	3-chloro	-S-	4-methyl phenyl	mp 222-224°C, Mass: m/z 368.1 (M ⁺), Anal. Calcd for C ₂₄ H ₂₄ ClNO ₄ S: C, 62.94; H, 5.28; N, 3.06. Found: C, 63.11; H, 5.35; N, 3.11.

151	3-chloro	-SO ₂ -	4-methyl phenyl	mp 226-228°C, Mass: m/z 400.1 (M ⁺), Anal. Calcd for C ₂₄ H ₂₄ ClNO ₆ S: C, 58.83; H, 4.94; N, 2.86. Found: C, 58.79; H, 4.94; N, 2.93.
152	3-chloro	-NH-	benzyl	mp 206-208°C, Mass: m/z 351.5 (M ⁺), Anal. Calcd for C ₂₄ H ₂₅ ClN ₂ O ₄ : C, 65.38; H, 5.72; N, 6.35. Found: C, 65.23; H, 5.86; N, 6.29.
153	3-chloro	-NH-	phenyl	mp 196-198°C, Mass: m/z 337.2 (M ⁺), Anal. Calcd for C ₂₃ H ₂₃ ClN ₂ O ₄ : C, 64.71; H, 5.43; N, 6.56. Found: C, 56.60; H, 4.90; N, 5.64.
154	3-chloro	-CH(OH)-	phenyl	mp 193-195°C, Mass: m/z 352.4 (M ⁺), Anal. Calcd for C ₂₄ H ₂₄ ClN ₂ O ₅ : C, 65.23; H, 5.47; N, 3.17. Found: C, 64.96; H, 5.60; N, 3.32.
155	3-chloro	-CH ₂ -	phenyl	mp 220-222°C, Mass: m/z 336.1 (M ⁺), Anal. Calcd for C ₂₄ H ₂₄ ClNO ₄ : C, 67.68; H, 5.68; N, 3.29. Found: C, 67.65; H, 5.83; N, 3.42.
156	3-chloro	-CH(F)-	phenyl	mp 182-184°C, Mass: m/z 354.3 (M ⁺), Anal. Calcd for C ₂₄ H ₂₃ ClFNO ₄ : C, 64.94; H, 5.22; N, 3.16. Found: C, 65.21; H, 5.26; N, 3.09.
157	3-chloro	-O-	4-fluoro phenyl	mp 218-220°C, Mass: m/z 356.2 (M ⁺), Anal. Calcd for C ₂₃ H ₂₁ ClFNO ₅ : C, 61.96; H, 4.75; N, 3.14. Found: C, 60.56; H, 4.67; N, 3.17.
158	3-trifluoromethyl	-O-	4-fluoro phenyl	mp 221-223°C, Mass: m/z 390.2 (M ⁺), Anal. Calcd for C ₂₄ H ₂₁ F ₄ NO ₅ : C, 60.13; H, 4.42; N, 2.92. Found: C, 59.18; H, 4.30; N, 2.91.
159	3-chloro	-O-	2-fluoro phenyl	mp 214-216°C, Mass: m/z 356.2 (M ⁺), Anal. Calcd for C ₂₃ H ₂₁ ClFNO ₅ : C, 61.96; H, 4.75; N, 3.14. Found: C, 61.42; H, 4.68; N, 3.21.
160	3-trifluoromethyl	-O-	2-fluoro phenyl	mp 218-220°C, Mass: m/z 390.2 (M ⁺), Anal. Calcd for C ₂₄ H ₂₁ F ₄ NO ₅ : C, 60.13; H, 4.42; N, 2.92. Found: C, 59.83; H, 4.34; N, 2.96.

161	3-chloro	-O-	3-fluoro phenyl	mp 219-221°C, Mass: m/z 356.2 (M ⁺), Anal. Calcd for C ₂₃ H ₂₁ ClFNO ₅ : C, 61.96; H, 4.75; N, 3.14. Found: C, 61.26; H, 4.74; N, 3.11.
162	3-triflororo methyl	-O-	3-fluoro phenyl	mp 221-223°C, Mass: m/z 390.2 (M ⁺), Anal. Calcd for C ₂₄ H ₂₁ F ₄ NO ₅ : C, 60.13; H, 4.42; N, 2.92. Found: C, 58.79; H, 4.28; N, 2.88.
163	3-chloro	-O-	naphth-2-yl	mp 229-231°C, Mass: m/z 388.1 (M ⁺), Anal. Calcd for C ₂₇ H ₂₄ ClNO ₅ : C, 67.85; H, 5.06; N, 2.93. Found: C, 67.71; H, 5.02; N, 3.03.
164	3-triflororo methyl	-O-	naphth-2-yl	mp 225-227°C, Mass: m/z 422.0 (M ⁺), Anal. Calcd for C ₂₈ H ₂₄ F ₃ NO ₅ : C, 65.75; H, 4.73; N, 2.74. Found: C, 65.72; H, 4.84; N, 2.88.
165	3-chloro	-O-	naphth-1-yl	mp 208-210°C, Mass: m/z 388.1 (M ⁺), Anal. Calcd for C ₂₇ H ₂₄ ClNO ₅ : C, 67.85; H, 5.06; N, 2.93. Found: C, 66.71; H, 5.11; N, 3.26.
166	3-triflororo methyl	-O-	naphth-1-yl	mp 211-213°C, Mass: m/z 422.0 (M ⁺), Anal. Calcd for C ₂₈ H ₂₄ F ₃ NO ₅ : C, 65.75; H, 4.73; N, 2.74. Found: C, 64.30; H, 4.76; N, 2.90.
167	3-chloro	-O-	H	mp 96-98°C, Mass: m/z 262.0 (M ⁺), Anal. Calcd for C ₁₅ H ₁₅ ClNO: C, 68.83; H, 6.16; N, 5.35. Found: C, 68.59; H, 5.99; N, 5.37. (isolated as the base)
168	3-triflororo methyl	-O-	H	mp 101-103°C, Mass: m/z 296.3 (M ⁺), Anal. Calcd for C ₁₆ H ₁₆ F ₃ NO: C, 65.07; H, 5.46; N, 4.74. Found: C, 65.06; H, 5.42; N, 4.80. (isolated as the base)
169	3-triflororo methyl	-O-	benzyl	mp 223-225°C, Mass: m/z 386.1 (M ⁺), Anal. Calcd for C ₂₅ H ₂₄ F ₃ NO ₅ : C, 63.15; H, 5.08; N, 2.94. Found: C, 63.22; H, 4.97; N, 3.02.
170	3-chloro	-O-	2,4-difluoro phenylsulfonyl	mp 201-203°C, Mass: m/z 438.0 (M ⁺), Anal. Calcd for C ₂₃ H ₂₀ ClF ₂ NO ₇ S: C, 52.32; H, 3.81; N, 2.65. Found: C, 52.26; H, 3.80; N, 2.71.

-71-

171	3-trifluoromethyl	-O-	2,4-difluorophenylsulfon-yl	mp 202-204°C, Mass: m/z 472.2 (M ⁺), Anal. Calcd for C ₂₄ H ₂₀ F ₅ NO ₇ S: C, 51.34; H, 3.59; N, 2.49. Found: C, 51.61; H, 3.65; N, 2.54.
172	3-chloro	-O-	thiazol-2-yl	mp 216-218°C, Mass: m/z 345.0 (M ⁺), Anal. Calcd for C ₂₀ H ₁₉ ClN ₂ O ₅ S: C, 55.23; H, 4.40; N, 6.44. Found: C, 55.15; H, 4.16; N, 6.43.
173	3-trifluoromethyl	-O-	thiazol-2-yl	mp 222-224°C, Mass: m/z 379.4 (M ⁺), Anal. Calcd for C ₂₁ H ₁₉ F ₃ N ₂ O ₅ S: C, 53.84; H, 4.08; N, 5.98. Found: C, 53.71; H, 3.95; N, 5.96.
174	3-chloro	-O-	pyrid-3-yl	mp 213-215°C, Mass: m/z 339.1 (M ⁺), Anal. Calcd for C ₂₂ H ₂₁ ClN ₂ O ₅ : C, 61.61; H, 4.93; N, 6.53. Found: C, 60.40; H, 4.89; N, 6.74.
175	3-trifluoromethyl	-O-	pyrid-3-yl	mp 221-223°C, Mass: m/z 373.1 (M ⁺), Anal. Calcd for C ₂₃ H ₂₁ F ₃ N ₂ O ₅ : C, 59.74; H, 4.57; N, 6.05. Found: C, 59.17; H, 4.47; N, 6.93.
176	3-methoxy	-O-	pyrid-3-yl	mp 101-103°C, Mass: m/z 335.2 (M ⁺), Anal. Calcd for C ₂₁ H ₂₄ Cl ₂ N ₂ O ₂ : C, 61.92; H, 5.93; N, 6.87. Found: C, 61.43; H, 6.07; N, 6.25. (isolated as the hydrochloride)
177	3-chloro	-O-	pyrid-4-yl	mp 154-156°C, Mass: m/z 339.1 (M ⁺), Anal. Calcd for C ₂₀ H ₂₁ Cl ₃ N ₂ O: C, 58.34; H, 5.14; N, 6.80. Found: C, 58.35; H, 5.18; N, 6.69. (isolated as the hydrochloride)
178	3-trifluoromethyl	-O-	pyrid-4-yl	mp 208-210°C, Mass: m/z 373.1 (M ⁺), Anal. Calcd for C ₂₁ H ₂₁ Cl ₂ F ₃ N ₂ O: C, 56.64; H, 4.75; N, 6.29. Found: C, 56.57; H, 4.68; N, 6.20. (isolated as the hydrochloride)
179	3-chloro	-O-	pyrimid-5-yl	mp 205-207°C, Mass: m/z 340.1 (M ⁺), Anal. Calcd for C ₂₁ H ₂₀ ClN ₃ O ₅ : C, 58.67; H, 4.68; N, 9.77. Found: C, 57.66; H, 4.70; N, 8.17.


-72-

180	3-trifluoro methyl	-O-	pyrimid-5-yl	mp 218-220°C, Mass: m/z 374.1 (M ⁺), Anal. Calcd for C ₂₂ H ₂₀ F ₃ N ₃ O ₅ : C, 57.02; H, 4.35; N, 9.06. Found: C, 56.55; H, 4.44; N, 8.89.
181	3-chloro	-O-	pyrid-2-yl	mp 93-95°C, Mass: m/z 339.1 (M ⁺), Anal. Calcd for C ₂₀ H ₂₁ Cl ₃ N ₂ O: C, 58.34; H, 5.14; N, 6.80. Found: C, 62.31; H, 5.30; N, 7.36. (isolated as the hydrochloride)
182	3-trifluoro methyl	-O-	pyrid-2-yl	mp 86-88°C, Mass: m/z 373.1 (M ⁺), Anal. Calcd for C ₂₁ H ₂₁ Cl ₂ F ₃ N ₂ O: C, 56.64; H, 4.75; N, 6.29. Found: C, 60.00; H, 4.92; N, 6.76. (isolated as the hydrochloride)
183	3-chloro	-NH-	pyrid-3-yl	mp 158-160°C, Mass: m/z 338.3 (M ⁺), Anal. Calcd for C ₂₂ H ₂₂ ClN ₃ O ₄ : C, 61.75; H, 5.18; N, 9.82. Found: C, 58.90; H, 4.64; N, 8.87.
184	3-trifluoro methyl	-NH-	pyrid-3-yl	mp 182-184°C, Mass: m/z 372.3 (M ⁺), Anal. Calcd for C ₂₃ H ₂₂ F ₃ N ₃ O ₄ : C, 59.86; H, 4.80; N, 9.10. Found: C, 58.33; H, 4.44; N, 8.60.
185	3-chloro	-NH-	pyrid-4-yl	mp 156-158°C, Mass: m/z 338.3 (M ⁺), Anal. Calcd for C ₂₀ H ₂₂ Cl ₃ N ₃ : C, 58.48; H, 5.39; N, 10.22. Found: C, 57.13; H, 5.49; N, 9.80. (isolated as the hydrochloride)
186	3-trifluoro methyl	-NH-	pyrid-4-yl	mp 142-144°C, Mass: m/z 372.3 (M ⁺), Anal. Calcd for C ₂₁ H ₂₂ Cl ₂ F ₃ N ₃ : C, 56.76; H, 4.99; N, 9.45. Found: C, 55.05; H, 4.88; N, 9.33. (isolated as the hydrochloride)
187	3-chloro	-NH-	pyrid-2-yl	mp 142-144°C, Mass: m/z 338.0 (M ⁺), Anal. Calcd for C ₂₀ H ₂₂ Cl ₃ N ₃ : C, 58.48; H, 5.39; N, 10.22. Found: C, 58.12; H, 5.39; N, 10.08. (isolated as the hydrochloride)
188	3-trifluoro methyl	-NH-	pyrid-2-yl	mp 144-146°C, Mass: m/z 372.1 (M ⁺), Anal. Calcd for C ₂₁ H ₂₂ Cl ₂ F ₃ N ₃ : C, 56.76; H, 4.99; N, 9.45. Found: C, 56.60; H, 5.04; N, 9.32. (isolated as the hydrochloride)

-73-

189	3-chloro	-O-	benzyl	MS m/e 351.9 (m+1)
190	3-trifluoromethyl	-NH-	phenyl	mp = 205-207°C; ms:m+1=371.1


By the method of Example 67 the following compounds were prepared, isolated as the oxalate except where noted:



No.	R ₁	Data
191	pyrid-4-yl	mp 176-178°C, Mass: m/z 305.2 (M ⁺), Anal. Calcd for C ₂₂ H ₂₂ N ₂ O ₅ : C, 66.99; H, 5.62; N, 7.10. Found: C, 67.55; H, 5.70; N, 7.24.
192	pyrid-3-yl	mp 198-200°C, Mass: m/z 305.2 (M ⁺), Anal. Calcd for C ₂₂ H ₂₂ N ₂ O ₅ : C, 66.99; H, 5.62; N, 7.10. Found: C, 64.98; H, 5.43; N, 6.86.
193	thien-2-yl	mp 234-236°C, Mass: m/z 310.2 (M ⁺), Anal. Calcd for C ₂₁ H ₂₁ NO ₅ S: C, 63.14; H, 5.29; N, 3.50. Found: C, 62.25; H, 5.18; N, 3.53.
194	imidazol-4-yl	mp 194-196°C, Mass: m/z 294.2 (M ⁺), Anal. Calcd for C ₂₀ H ₂₀ N ₃ O ₅ : C, 62.65; H, 5.52; N, 10.95. Found: C, 59.94; H, 5.30; N, 10.12.
195	naphth-2-yl	mp 223-225°C, Mass: m/z 354.4 (M ⁺), Anal. Calcd for C ₂₇ H ₂₅ NO ₅ : C, 73.12; H, 5.68; N, 3.16. Found: C, 73.38; H, 5.94; N, 3.40.
196	naphth-1-yl	mp 223-225°C, Mass: m/z 354.4 (M ⁺), Anal. Calcd for C ₂₇ H ₂₅ NO ₅ : C, 73.12; H, 5.68; N, 3.16. Found: C, 73.18; H, 5.52; N, 3.23.

5

By the method of Example 67 the following compounds were prepared, isolated as the oxalate except where noted:




No.	R ₁	Data
197	3-chlorophenyl	mp 240-242°C, Mass: m/z 336.0 (M ⁺), Anal. Calcd for C ₂₃ H ₂₀ ClNO ₅ : C, 55.23; H, 4.40; N, 6.44. Found: C, 55.15; H, 4.16; N, 6.43.
198	3-trifluoromethylphenyl	mp 255-257°C, Mass: m/z 370.0 (M ⁺), Anal. Calcd for C ₂₄ H ₂₀ F ₃ NO ₅ : C, 62.74; H, 4.38; N, 3.04. Found: C, 62.95; H, 4.27; N, 3.08.
199	5-methoxy-1H-indol-3-yl	mp 232-234°C, Mass: m/z 371.1 (M ⁺), Anal. Calcd for C ₂₆ H ₂₄ N ₂ O ₆ : C, 67.81; H, 5.25; N, 6.08. Found: C, 67.46; H, 4.44; N, 5.44.

-74-

200	1H-indol-3-yl	mp 221-223°C, Mass: m/z 341.1 (M^+), Anal. Calcd for $C_{25}H_{22}N_2O_5$: C, 69.75; H, 5.15; N, 6.50. Found: C, 71.99; H, 4.48; N, 6.40.
-----	---------------	--


By the method of Example 67 the following compounds were prepared, isolated as the oxalate except where noted:



No.	R ₁	R ₉	Data
201	3-chlorophenyl	phenyl	mp 225-227°C, Mass: m/z 361.1 (M^+), Anal. Calcd for $C_{25}H_{23}ClN_2O_4$: C, 66.59; H, 5.14; N, 6.21. Found: C, 66.21; H, 5.02; N, 6.14.
202	3-trifluoromethyl phenyl	phenyl	mp 216-218°C, Mass: m/z 395.1 (M^+), Anal. Calcd for $C_{25}H_{25}F_3N_2O_4$: C, 64.45; H, 4.78; N, 5.78. Found: C, 63.98; H, 4.67; N, 5.76.
203	5-methoxy-1H-indol-3-yl	phenyl	mp 208-210°C, Mass: m/z 394.2 (M^+), Anal. Calcd for $C_{28}H_{27}N_3O_5$: C, 69.26; H, 5.60; N, 8.62. Found: C, 67.78; H, 5.29; N, 8.42.
204	1H-indol-3-yl	phenyl	mp 227-229°C, Mass: m/z 364.3 (M^+), Anal. Calcd for $C_{27}H_{25}N_3O_4$: C, 71.19; H, 5.53; N, 9.22. Found: C, 70.02; H, 5.33; N, 8.95.
205	5-methoxy-1H-indol-3-yl	H	mp 170-172°C, Mass: m/z 318.2 (M^+), Anal. Calcd for $C_{22}H_{23}N_3O_5$: C, 64.53; H, 5.62; N, 10.26. Found: C, 56.16; H, 4.98; N, 8.75.

5


By the method of Example 67 the following compounds were prepared, isolated as the oxalate except where noted:



No.	R ₁	R ₉	Data
206	3-chlorophenyl	phenyl	mp 237-239°C, Mass: m/z 361.1 (M^+), Anal. Calcd for $C_{25}H_{23}ClN_2O_4$: C, 66.59; H, 5.14; N, 6.21. Found: C, 66.55; H, 5.16; N, 6.20.
207	3-trifluoromethyl phenyl	phenyl	mp 239-241°C, Mass: m/z 395.1 (M^+), Anal. Calcd for $C_{25}H_{25}F_3N_2O_4$: C, 64.45; H, 4.78; N, 5.78. Found: C, 64.59; H, 4.83; N, 5.83.
208	5-methoxy-1H-indol-3-yl	phenyl	mp 194-196°C, Mass: m/z 396.2 (M^+), Anal. Calcd for $C_{28}H_{27}N_3O_5$: C, 69.26; H, 5.60; N, 8.62. Found: C, 68.33; H, 5.37; N, 8.52.


-75-

209	1H-indol-3-yl	phenyl	mp 206-208°C, Mass: m/z 366.2 (M^+), Anal. Calcd for $C_{27}H_{25}N_3O_4$: C, 71.19; H, 5.53; N, 9.22. Found: C, 69.23; H, 5.42; N, 8.86.
210	5-methoxy-1H-indol-3-yl	H	mp 186-188°C, Mass: m/z 318.2 (M^+), Anal. Calcd for $C_{22}H_{23}N_3O_5$: C, 64.53; H, 5.66; N, 10.26. Found: C, 62.88; H, 4.61; N, 9.27.

Example 220N-(2-(3-Chlorophenyl)ethyl)-3-benzoylbenzylamine

5 Combine 3-benzoylbenzaldehyde (0.45 g, 2.1 mmol), and (3-chlorophenyl)ethylamine (0.3 ml, 2.1 mmol) and 3Å molecular sieves (1.0 g) in MeOH (30 ml). Heat to reflux. After 3 hours, cool, filter, and concentrate to give a residue. Dissolve the residue in dichloroethane (20 ml), add acetic acid (0.12 ml, 2.1 mmol) and sodium triacetoxyborohydride (0.6 g, 2.94 mmol) and stir at ambient temperature. After 2
10 hours, concentrate the reaction mixture and add dichloromethane (90 ml) and extract sequentially with distilled water (50 ml) and then brine (50 ml). Dry the organic layer over Na_2SO_4 and give a residue. Chromatograph the residue on silica gel eluting with EtOAc to give the title compound as the base.

 The oxalate using the method of Example 67 to give the title compound: mp 196-
15 198°C, Mass: m/z 350.4 (M^+), Anal. Calcd for $C_{24}H_{22}ClNO_5$: C, 65.53; H, 5.04; N, 3.18. Found: C, 65.27; H, 5.20; N, 3.13.

Example 221N-(2-(3-Chlorophenyl)ethyl)-3-ethoxybenzylamine


20

 Combine 3-ethoxybenzaldehyde (3.38 g, 22.5 mmol), 2-(3-chlorophenyl)ethylamine (2.33 g, 15.0 mol), and 3Å molecular sieves (2.88 g) in ethanol (230 ml). Stir the reaction mixture at reflux for 4 hours. Filter to remove the molecular sieves and then slowly add sodium borohydride (1.70 g, 45.0 mmol) to the filtrate and stir
25 at ambient temperature. After 15 hours, concentrate the reaction mixture to a residue, dissolve the residue in 1 N NaOH, and extract with dichloromethane. Combine organic extracts, wash with brine, dry over Na_2SO_4 , and concentrate to a residue. Chromatograph

-76-


the residue on silica gel eluting with ethyl acetate to give the title compound. The HCl salt was prepared in ethyl acetate to give the title compound: mp 178-180 °C; MS (ACPI): m/e 290.1 (M+1); Analysis for C₁₇H₂₁Cl₂NO: Calcd: C, 62.58; H, 6.49; N, 4.29; Found: C, 62.65; H, 6.53; N, 4.32.

- 5 By the method of Example 221 the following compounds were prepared, isolated as the maleate except where noted:



No.	Z''	R ₄	Data
222	3-chloro	propyl	mp 138-140 °C. MS (ACPI): m/e 304.1 (M+1). Analysis for C ₁₈ H ₂₃ Cl ₂ NO: Calcd: C, 63.53; H, 6.81; N, 4.12; found: C, 63.74; H, 6.81; N, 4.22. (isolated as the hydrochloride)
223	3-trifluoro methyl	propyl	mp 145-147 °C. MS (ACPI): m/e 338.1 (M+1). Analysis for C ₂₃ H ₂₆ F ₃ NO ₅ : Calcd: C, 60.92; H, 5.78; N, 3.09; found: C, 60.77; H, 5.60; N, 3.12.
224	3-trifluoro methyl	ethyl	mp 164-166 °C. MS (ACPI): m/e 324.2 (M+1). Analysis for C ₁₈ H ₂₁ ClF ₃ NO: Calcd: C, 60.09; H, 5.88; N, 3.89; found: C, 60.42; H, 5.80; N, 3.93. (isolated as the hydrochloride)
225	2-phenyl	2,2,2-trifluoro ethyl	mp 181-183 °C. MS (ACPI): m/e 386.2 (M+1). Analysis for C ₂₇ H ₂₆ F ₃ NO ₅ : Calcd: C, 64.67; H, 5.23; N, 2.79; found: C, 64.52; H, 5.01; N, 2.85.
226	4-phenyl	2,2,2-trifluoro ethyl	mp 39 °C. MS (ACPI): m/e 386.2 (M+1). (Exception- one equivalent of triethylamine in the reaction) (isolated as the free base)

- 10 By the method of Example 221 the following compounds were prepared, isolated as the maleate except where noted:



No.	Z'	R ₄	Data
227	5-chloro	ethyl	mp 153-156 °C, MS (ACPI): m/e 329.1 (M+1). Analysis for C ₂₃ H ₂₅ ClN ₂ O ₅ : Calcd: C, 62.09; H, 5.66; N, 6.30; found: C, 62.27; H, 5.38; N, 6.19
228	5-chloro	propyl	mp 163-166 °C. MS (ACPI): m/e 343.1 (M+1). Analysis for C ₂₄ H ₂₇ ClN ₂ O ₅ : Calcd: C, 62.81; H, 5.93; N, 6.10; found: C, 63.07; H, 5.80; N, 6.07.

229	5-chloro	2,2,2-trifluoroethyl	mp 178-181 °C, MS (ACPI): m/e 383.1 (M+1). Analysis for C ₂₃ H ₂₂ ClF ₃ N ₂ O ₅ : Calcd: C, 55.37; H, 4.44; N, 5.62; found: C, 55.71; H, 4.39; N, 5.66.
230	5-chloro	3-fluoropropyl	mp 167-170 °C, MS (ACPI): m/e 361.1 (M+1). Analysis for C ₂₄ H ₂₆ ClF ₃ N ₂ O ₅ : Calcd: C, 60.44; H, 5.49; N, 5.87; found: C, 60.30; H, 5.25; N, 5.78.
231	5-chloro	2,2,3,3,3-pentafluoro propyl	mp 170-173 °C, MS (ACPI): m/e 433.1 (M+1). Analysis for C ₂₄ H ₂₂ ClF ₅ N ₂ O ₅ : Calcd: C, 52.52; H, 4.04; N, 5.10; found: C, 52.49; H, 4.06; N, 5.16.
232	5-chloro	2,2,3,3-tetrafluoro propyl	mp 163-167 °C, MS (ACPI): m/e 415.1 (M+1). Analysis for C ₂₄ H ₂₃ ClF ₄ N ₂ O ₅ : Calcd: C, 54.30; H, 4.37; N, 5.28; found: C, 54.47; H, 4.36; N, 5.33.
233	5-methoxy	2,2,2-trifluoroethyl	mp 179-182 °C, MS (ACPI): m/e 379.1 (M+1). Analysis for C ₂₄ H ₂₅ F ₃ N ₂ O ₆ : Calcd: C, 58.30; H, 5.10; N, 5.67; found: C, 58.26; H, 5.09; N, 5.69.
234	6-chloro	2,2,3,3-tetrafluoro propyl	mp 156-160 °C, MS (ACPI): m/e 415.1 (M+1). Analysis for C ₂₄ H ₂₃ ClF ₄ N ₂ O ₅ : Calcd: C, 54.30; H, 4.37; N, 5.28; found: C, 54.31; H, 4.34; N, 5.31.
235	5-cyano	2,2,2-trifluoroethyl	mp 176-178 °C, MS (ACPI): m/e 374.0 (M+1). Analysis for C ₂₀ H ₁₉ ClF ₃ N ₃ O: Calcd: C, 58.61; H, 4.67; N, 10.25; found: C, 58.52; H, 4.61; N, 10.17. (isolated as the hydrochloride)
236	5-methyl sulfonyl	2,2,2-trifluoro ethyl	mp 193-195 °C, MS (ACPI): m/e 429.9 (M+1). Analysis for C ₂₄ H ₂₅ F ₃ N ₂ O ₇ S: Calcd: C, 53.13; H, 4.64; N, 5.16; found: C, 53.12; H, 4.58; N, 5.20.
237	5-cyano	3,3,3-trifluoro propyl	mp 150-154 °C, MS (ACPI): m/e 387.9 (M+1). Analysis for C ₂₅ H ₂₄ F ₃ N ₃ O ₅ : Calcd: C, 59.64; H, 4.80; N, 8.35; found: C, 59.55; H, 4.77; N, 8.38.
238	5-methyl sulfonyl	3,3,3-trifluoro propyl	mp 178-181 °C, MS (ACPI): m/e 440.9 (M+1). Analysis for C ₂₅ H ₂₇ F ₃ N ₂ O ₇ S: Calcd: C, 53.95; H, 4.89; N, 5.03; found: C, 53.87; H, 4.86; N, 5.04.
239	4-fluoro	2,2,2-trifluoro ethyl	mp 199-202 °C, MS (ACPI): m/e 367.2 (M+1). Analysis for C ₁₉ H ₁₉ ClF ₄ N ₂ O: calcd: C, 56.65; H, 4.75; N, 6.95; found: C, 56.82; H, 4.65; N, 6.84. (isolated as the hydrochloride)
240	4-fluoro	2,2,3,3,3-pentafluoro propyl	mp 118-121 °C, MS (ACPI): m/e 417.2 (M+1). Analysis for C ₂₄ H ₂₂ F ₆ N ₂ O ₅ : Calcd: C, 54.14; H, 4.16; N, 5.26; found: C, 54.39; H, 4.25; N, 5.30.
241	4-fluoro	2,2,3,3-tetrafluoro propyl	mp 188-191 °C, MS (ACPI): m/e 399.0 (M+1). Analysis for C ₂₀ H ₂₀ ClF ₅ N ₂ O: Calcd: C, 55.24; H, 4.64; N, 6.44; found: C, 55.03; H, 4.53; N, 6.34. (isolated as the hydrochloride)
242	7-fluoro	2,2,2-trifluoro ethyl	mp 157-160 °C, MS (ACPI): m/e 367.2 (M+1). Analysis for C ₂₃ H ₂₂ F ₄ N ₂ O ₅ : Calcd: C, 57.26; H, 4.60; N, 5.81; found: C, 57.34; H, 4.39; N, 6.11.

243	7-fluoro	2,2,3,3,3-pentafluoro propyl	mp 166-168 °C. MS (ACPI): m/e 417.2 (M+1). Analysis for C ₂₄ H ₂₂ F ₆ N ₂ O ₅ : Calcd: C, 54.14; H, 4.16; N, 5.26; found: C, 53.99; H, 3.98; N, 5.61.
244	7-fluoro	2,2,3,3-tetrafluoro propyl	mp 170-173 °C. MS (ACPI): m/e 399.2 (M+1). Analysis for C ₂₄ H ₂₄ F ₅ N ₂ O ₅ : Calcd: C, 56.03; H, 4.51; N, 5.45; found: C, 55.73; H, 4.30; N, 5.66.
245	5-amido	3,3,3-trifluoro propyl	mp 143-147 °C. MS (ACPI): m/e 406.1 (M+1). Analysis for C ₂₁ H ₂₂ F ₃ N ₃ O ₂ : Calcd: C, 62.22; H, 5.47; N, 10.36; found: C, 61.96; H, 5.42; N, 10.13. (isolated as the base)
246	5-amido	2,2,2-trifluoro ethyl	mp 125-130 °C. MS (ACPI): m/e 392.1 (M+1). Analysis for C ₂₀ H ₂₁ ClF ₃ N ₃ O ₂ : Calcd: C, 56.15; H, 4.95; N, 9.82; found: C, 55.80; H, 4.93; N, 9.71. (isolated as the hydrochloride)
247	6-phenyl	2,2,2-trifluoro ethyl	mp 117-120 °C. MS (ACPI): m/e 425.1 (M+1). Analysis for C ₂₆ H ₂₃ F ₃ N ₂ O: Calcd: C, 70.74; H, 5.46; N, 6.60; found: C, 70.75; H, 5.42; N, 6.66. (isolated as the base)
248	6-methyl	2,2,3,3,3-pentafluoro propyl	m.p. 168-170 °C. MS (ACPI): m/e 413.2 (M+1). Analysis for C ₂₁ H ₂₂ F ₅ N ₂ O: Calcd: C, 56.82; H, 4.77; N, 5.30; found: C, 57.21; H, 4.46; N, 5.33
249	6-phenyl	2,2,3,3,3-pentafluoro propyl	mp 110.5-113.5 °C. MS (ACPI): m/e 475.1 (M+1). Analysis for C ₂₆ H ₂₃ F ₅ N ₂ O: Calcd: C, 65.82; H, 4.89; N, 5.90; found: C, 65.70; H, 4.84; N, 5.93. (isolated as the base)
250	6-phenyl	2,2,3,3-tetrafluoro propyl	mp 94-98 °C. MS (ACPI): m/e 457.1 (M+1). Analysis for C ₂₆ H ₂₄ F ₄ N ₂ O: calcd: C, 68.41; H, 5.30; N, 6.14; found: C, 68.18; H, 5.28; N, 6.06 (isolated as the base)
251	6-methyl	2,2,2-trifluoro ethyl	mp 176-178 °C. MS (ACPI): m/e 363.1 (M+1). Analysis for C ₂₀ H ₂₂ ClF ₃ N ₂ O: Calcd: C, 60.23; H, 5.56; N, 7.02; found: C, 60.16; H, 5.43; N, 6.98. (isolated as the hydrochloride)
252	6-methyl	2,2,3,3-tetrafluoro propyl	mp 156-158 °C. MS (ACPI): m/e 395.1 (M+1). Analysis for C ₂₁ H ₂₃ ClF ₄ N ₂ O: Calcd: C, 58.54; H, 5.38; N, 6.50; found: C, 58.60; H, 5.32; N, 6.55. (isolated as the hydrochloride)
253	6-ethoxy carbonyl	2,2,3,3-tetrafluoro propyl	mp 166-168 °C. MS (ACPI): m/e 453.1 (M+1). Analysis for C ₂₃ H ₂₅ ClF ₄ N ₂ O ₃ : Calcd: C, 56.50; H, 5.15; N, 5.73; found: C, 56.18; H, 5.00; N, 5.66. (isolated as the hydrochloride)
254	6-ethoxy carbonyl	2,2,2-trifluoro ethyl	mp 169.5-171.5 °C. MS (ACPI): m/e 421.2 (M+1). Analysis for C ₂₆ H ₂₇ F ₃ N ₂ O ₇ : Calcd: C, 58.21; H, 5.07; N, 5.22; found: C, 58.43; H, 4.85; N, 5.27.
255	6-cyano	2,2,2-trifluoro ethyl	mp 175-177 °C. MS (ACPI): m/e 374.1 (M+1). Analysis for C ₂₄ H ₂₂ F ₃ N ₃ O ₅ : Calcd: C, 58.90; H, 4.53; N, 8.59; found: C, 58.62; H, 4.48; N, 8.50.

256	6-cyano	2,2,3,3-tetrafluoropropyl	mp 167-169 °C. MS (ACPI): m/e 406.1 (M+1). Analysis for C ₂₅ H ₂₃ F ₄ N ₃ O ₅ : Calcd: C, 57.58; H, 4.45; N, 8.06; found: C, 57.31; H, 4.35; N, 8.08.
257	6-amido	2,2,2-tetrafluoroethyl	mp 102 °C. MS (ACPI): m/e 392.2 (M+1). Analysis for C ₂₀ H ₂₀ F ₃ N ₃ O ₂ : Calcd: C, 61.38; H, 5.15; N, 10.74; found: C, 61.68; H, 5.11; N, 10.65. (isolated as the base)
258	6-amido	2,2,3,3-tetrafluoropropyl	mp 120 °C. MS (ACPI): m/e 424.3 (M+1). Analysis for C ₂₁ H ₂₁ F ₄ N ₃ O ₂ : Calcd: C, 59.57; H, 5.00; N, 9.92; found: C, 59.33; H, 4.82; N, 9.79. (isolated as the base)
259	6-trifluoromethoxy	2,2,3,3-tetrafluoropropyl	mp 132-134 °C. MS (ACPI): m/e 465.1 (M+1). Analysis for C ₂₁ H ₂₀ ClF ₇ N ₂ O ₂ : Calcd: C, 50.36; H, 4.03; N, 5.59; found: C, 50.25; H, 3.96; N, 5.58. (isolated as the hydrochloride)
260	6-trifluoromethoxy	2,2,2-trifluoroethyl	mp 160-164 °C. MS (ACPI): m/e 433.1 (M+1). Analysis for C ₂₀ H ₁₉ ClF ₆ N ₂ O ₂ : Calcd: C, 51.24; H, 4.08; N, 5.98; found: C, 51.26; H, 3.99; N, 5.96. (isolated as the hydrochloride)
260 A	7-chloro	2,2,3,3-tetrafluoropropyl	mp 153.6-154.4 °C. MS (APCI): m/e 415.1 (M+1). Anal. Calcd. for C ₂₀ H ₁₉ ClF ₄ N ₂ O•1.0HCl: C, 53.23; H, 4.47; N, 6.21. Found: C, 52.89; H, 4.40; N, 6.18. (isolated as the hydrochloride)
260 B	7-chloro	2,2,2-trifluoroethyl	mp 193.4-194.9 °C. Mass (ES+): m/z 383.17 (M+1). Anal. Calcd. for C ₁₉ H ₁₈ ClF ₃ N ₂ O•1.0HCl: C, 54.43; H, 4.57; N, 6.68. Found: C, 54.66; H, 4.39; N, 6.66. (isolated as the hydrochloride)

Example 261N-(2-(7-Fluoro-1H-indol-3-yl)ethyl)-3-(2,2,3,3-tetrafluoropropoxy)benzylamine hydrochloride

- 5 Add acetyl chloride (2.4 mL, 33.8 mmol) dropwise to anhydrous ethanol (50 mL) and stir the solution for 10 min at ambient temperature and add to a solution of N-(2-(7-fluoro-1H-indol-3-yl)ethyl)-3-(2,2,3,3-tetrafluoropropoxy)benzylamine (12.0 g, 30.1 mmol) in ethyl acetate. Concentrate the resulting solution under reduced pressure to give a yellow solid. Recrystallize the yellow solid from ethyl acetate/ethanol/diethyl ether to give
- 10 the title compound: mp 142-143 °C. MS(m/e): 399 (M+1), 397 (M-1). Calculated for C₂₀H₁₉F₅N₂O•HCl: Calcd: C, 55.24; H, 4.64; N, 6.44. Found: C, 55.44; H, 4.66; N, 6.46.

-80-

Example 262

(N-(2-(7-Fluoro-1H-indol-3-yl)ethyl)-3-(2,2,3,3-tetrafluoropropoxy)benzylamine hydrochloride L(+)-tartrate

- 5 Add L-(+)-tartaric acid (49 mg, 0.33 mmol) and methanol to a solution of (N-(2-(7-fluoro-1H-indol-3-yl)ethyl)-3-(2,2,3,3-tetrafluoropropoxy)benzylamine (130 mg, 0.33 mmol) in ethyl acetate. Evaporate the solvent to give a gum. Crystallize the gum from diethyl ether/ethyl acetate to give the title compound: mp 192-194 °C.

Example 263

- 10 N-(2-(7-Fluoro-1H-indol-3-yl)ethyl)-3-(2,2,2-trifluoroethoxy)benzylamine hydrochloride

- Add dropwise acetyl chloride (2.3 mL, 32.4 mmol) to anhydrous ethanol (50 mL) and stir the solution for 10 min at ambient temperature and add to a solution N-(2-(7-fluoro-1H-indol-3-yl)ethyl)-3-(2,2,2-trifluoroethoxy)benzylamine (10.7 g, 29.2 mmol) in diethyl ether. Concentrate the resulting solution under reduced pressure to give a yellow
15 solid. Recrystallize the yellow solid from ethyl acetate/methanol to give the title compound: mp 163-164 °C; MS(m/e): 367 (M+1), 365 (M-1); Calculated for C₁₉H₁₈F₄N₂O HCl: Calcd: C, 56.65; H, 4.75; N, 6.95. Found: C, 56.45; H, 4.54; N, 6.90.


Example 264

- 20 N-(2-(7-Fluoro-1H-indol-3-yl)ethyl)-3-(2,2,2-trifluoroethoxy)benzylamine L(+)-tartrate

- Add L-(+)-tartaric acid (295 mg, 1.96 mmol) in methanol to a solution of N-(2-(7-fluoro-1H-indol-3-yl)ethyl)-3-(2,2,2-trifluoroethoxy)benzylamine (720 mg, 1.96 mmol) in ethyl acetate. Concentrate under reduced pressure the resulting solution to give a clear colorless oil. Crystallize the oil from diethyl ether to give the title compound: mp 118-119
25 °C. MS(m/e): 367 (M+1), 365 (M-1). Calculated for C₁₉H₁₈F₄N₂O·C₄H₆O₆: Calcd: C, 53.49; H, 4.68; N, 5.42. Found: C, 53.21; H, 4.55; N, 5.41.

Example 270

N-(2-(5-Fluoro-1H-indol-3-yl)ethyl)-3-propoxybenzylamine hydrochloride



30


- Combine 3-propoxybenzaldehyde (2.96 g, 18.0 mmol), 5-fluorotryptamine hydrochloride (2.58 g, 12.0 mol), triethylamine (1.15 g), and 3 Å molecular sieves (2.27 g) in ethanol (200 ml). Stir the reaction mixture at reflux for 4 hours. Filter to remove the

-81-

molecular sieves and then slowly add sodium borohydride (1.36 g, 36.0 mmol) to the filtrate and stir at ambient temperature. After 15 hours, concentrate the reaction mixture to a residue, dissolve the residue in 1 N NaOH, and extract with dichloromethane.

Combine organic extracts, wash with brine, dry over Na₂SO₄, and concentrate to a residue. Chromatograph the residue on silica gel eluting with ethyl acetate yielded 3.31 g of an oil. The HCl salt was prepared in diethyl ether: mp 197-199°C; MS (ACPI): m/e 327.2 (M+1); Analysis for C₂₀H₂₄ClFN₂O: Calcd: C, 66.20; H, 6.67; N, 7.72; found: C, 66.06; H, 6.63; N, 7.76.

By the method of Example 270 the following compounds were prepared, isolated as the maleate except where noted:




No.	Z'	R ₄	Data
271	5-fluoro	ethyl	mp 196-198 °C, MS (ACPI): m/e 313.1 (M+1). Analysis for C ₁₉ H ₂₂ ClFN ₂ O: Calcd: C, 65.42; H, 6.36; N, 8.03; found: C, 65.48; H, 6.30; N, 8.04. (isolated as the hydrochloride)
272	5-trifluoro methyl	ethyl	mp 156-160 °C, MS (ACPI): m/e 363.1 (M+1). Analysis for C ₂₄ H ₂₅ F ₃ N ₂ O ₅ : Calcd: C, 60.25; H, 5.27; N, 5.85; found: C, 60.47; H, 5.26; N, 5.79.
273	5-trifluoro methyl	propyl	mp 169-172 °C, MS (ACPI): m/e 377.1 (M+1). Analysis for C ₂₅ H ₂₇ F ₃ N ₂ O ₅ : Calcd: C, 60.97; H, 5.53; N, 5.69; found: C, 60.95; H, 5.54; N, 5.70
274	5-trifluoro methyl	2,2,2-trifluoro ethyl	mp 180-184 °C. MS (ACPI): m/e 417.1 (M+1). Analysis for C ₂₄ H ₂₂ F ₆ N ₂ O ₅ : Calcd: C, 54.14; H, 4.16; N, 5.26; found: C, 53.99; H, 4.07; N, 5.61.
275	5-trifluoro methyl	3,3,3-trifluoro propyl	mp 158-161 °C, MS (ACPI): m/e 431.1 (M+1). Analysis for C ₂₅ H ₂₄ F ₆ N ₂ O ₅ : Calcd: C, 54.95; H, 4.43; N, 5.13; found: C, 54.84; H, 4.46; N, 5.03.
276	4-methoxy	2,2,3,3-tetrafluoro propyl	mp 144-147 °C. MS (ACPI): m/e 411.1 (M+1). Analysis for C ₂₅ H ₂₆ F ₄ N ₂ O ₆ : Calcd: C, 57.03; H, 4.98; N, 5.32; found: C, 56.84; H, 4.94; N, 5.34.
277	5-cyano	2,2,3,3-tetrafluoro propyl	mp 172-174 °C. MS (ACPI): m/e 406.2 (M+1). Analysis for C ₂₅ H ₂₃ F ₄ N ₃ O ₅ : Calcd: C, 57.58; H, 4.45; N, 8.06; found: C, 57.91; H, 4.13; N, 8.34.
278	5-cyano	2,2,3,3,3-pentafluoro propoxy	mp 168-170 °C. MS (ACPI): m/e 424.1 (M+1). Analysis for C ₂₅ H ₂₂ F ₅ N ₃ O ₅ : Calcd: C, 55.66; H, 4.11; N, 7.79; found: C, 55.54; H, 4.16; N, 7.71.

279	5-(4-fluorophenyl)	2,2,3,3-tetrafluoro propyl	p 161-165 °C. MS (ACPI): m/e 475.1 (M+1). Analysis for $C_{26}H_{24}ClF_5N_2O$: Calcd: C, 61.12; H, 4.73; N, 5.48; found: C, 61.18; H, 4.64; N, 5.50. (isolated as the hydrochloride)
280	5-(4-fluorophenyl)	2,2,3,3,3-pentafluoro propoxy	(isolated as the hydrochloride) mp 168-171 °C. MS (ACPI): m/e 493.1 (M+1). Analysis for $C_{26}H_{23}ClF_6N_2O$: Calcd: C, 59.04; H, 4.38; N, 5.30; found: C, 59.15; H, 4.28; N, 5.30. (isolated as the hydrochloride)
281	5-phenyl	2,2,3,3-tetrafluoro propyl	mp 148-151 °C. MS (ACPI): m/e 457.1 (M+1). Analysis for $C_{26}H_{25}ClF_4N_2O$: Calcd: C, 63.35; H, 5.11; N, 5.68; found: C, 63.16; H, 4.99; N, 5.67. (isolated as the hydrochloride)
282	5-phenyl	2,2,3,3,3-pentafluoro prop	mp 65-70 °C, dec. MS (ACPI): m/e 475.1 (M+1). Analysis for $C_{26}H_{24}ClF_5N_2O$: Calcd: C, 61.12; H, 4.73; N, 5.48; found: C, 60.98; H, 4.66; N, 5.41. (isolated as the hydrochloride)
283	5-(4-fluorophenyl)	2,2,2-trifluoro ethyl	mp 214-216 °C. MS (ACPI): m/e 443.1 (M+1). Analysis for $C_{25}H_{23}ClF_4N_2O$: Calcd: C, 62.70; H, 4.84; N, 5.85; found: C, 62.47; H, 4.71; N, 5.79. (isolated as the hydrochloride)
284	5-phenyl	2,2,2-trifluoro ethyl	mp 171-174 °C, dec. MS (ACPI): m/e 425.1 (M+1). Analysis for $C_{25}H_{24}ClF_3N_2O$: Calcd: C, 65.15; H, 5.25; N, 6.08; found: C, 65.46; H, 5.17; N, 6.10. (isolated as the hydrochloride)
285	4-phenyl	2,2,3,3,3-pentafluoro propyl	mp 55 °C, dec. MS (ACPI): m/e 475.1 (M+1). Analysis for $C_{26}H_{24}ClF_5N_2O$: Calcd: C, 61.12; H, 4.73; N, 5.48; found: C, 61.11; H, 4.83; N, 5.40. (isolated as the hydrochloride)
286	4-phenyl	2,2,2-trifluoro ethyl	mp 60 °C, dec. MS (ACPI): m/e 425.1 (M+1). Analysis for $C_{25}H_{24}ClF_3N_2O$: Calcd: C, 65.15; H, 5.25; N, 6.08; found: C, 65.08; H, 5.42; N, 5.93. (isolated as the hydrochloride)
287	4-phenyl	2,2,3,3-tetrafluoro propyl	mp 56 °C, dec. MS (ACPI): m/e 457.1 (M+1). Analysis for $C_{26}H_{25}ClF_4N_2O$: Calcd: C, 63.35; H, 5.11; N, 5.68; found: C, 63.60; H, 5.35; N, 5.48. (isolated as the hydrochloride)
288	7-fluoro	pyrid-4-yl	mp 212-214 °C. MS (ACPI): m/e 362.2 (M+1). (isolated as the oxalate)
289	7-fluoro	pyrid-3-yl	mp 167-169 °C. MS (ACPI): m/e 362.3 (M+1). (isolated as the oxalate)
299	7-phenyl	2,2,2-trifluoro ethyl	mp 116-120 °C. MS (ACPI): m/e 425.3 (M+1). Analysis for $C_{29}H_{27}F_3N_2O_5$: Calcd: C, 64.44; H, 5.03; N, 5.18; found: C, 64.47; H, 4.96; N, 5.24.
300	7-phenyl	2,2,3,3-tetrafluoro propyl	mp 108-111 °C. MS (ACPI): m/e 457.3 (M+1). Analysis for $C_{30}H_{28}F_4N_2O_5$: Calcd: C, 62.93; H, 4.93; N, 4.89; found: C, 63.02; H, 4.91; N, 4.96.


Example 301

N-(2-(6-Chloro-1H-indol-3-yl)ethyl)-N-methyl-3-(2,2,3,3-tetrafluoropropoxy)benzylamine maleate




5 Add 3-(2,2,3,3-tetrafluoropropoxy)benzaldehyde (232.6 mg, 0.98 mmol) to a solution of N-(2-(6-chloro-1H-indol-3-yl)ethyl)-N-methylamine (205.6 mg, 0.98 mmol) and sodium triacetoxy borohydride (305.3 mg, 1.37 mmol) in dichloroethane (50 ml). Stir at ambient temperature. After 24 hours, evaporate to residue and dissolve the residue in 10 1N NaOH then extract with dichloromethane. Combine the organic extracts, washed with brine, dry (Na₂SO₄), filter, and evaporate to a residue. Chromatograph the residue on silica gel eluting with ethyl acetate to give the title compound. The maleate salt was prepared in diethyl ether: mp 125-128 °C. MS (ACPI): m/e 429.3 (M+1). Analysis for C₂₅H₂₅ClF₄N₂O₅: Calcd: C, 55.10; H, 4.62; N, 5.14; found: C, 55.13; H, 4.59; N, 5.09.

15 By the method of Example 301 the following compounds were prepared:




No.	Z'	R ₄	Data
302	5-methoxy	2,2,2-trifluoro ethyl	mp 144-147 °C. MS (ACPI): m/e 393.1 (M+1). Analysis for C ₂₃ H ₂₅ F ₃ N ₂ O ₆ : Calcd: C, 57.26; H, 5.22; N, 5.81; found: C, 56.89; H, 5.16; N, 5.82. (isolated as the oxalate)
303	4-methoxy	2,2,3,3-tetrafluoro propyl	mp 104-109 °C. MS (ACPI): m/e 425.2 (M+1). Analysis for C ₂₄ H ₂₆ F ₄ N ₂ O ₆ : Calcd: C, 56.03; H, 5.09; N, 5.45; found: C, 55.85; H, 5.05; N, 5.43. (isolated as the oxalate)
304	4-fluoro	2,2,2-trifluoro ethyl	mp 199-202 °C. MS (ACPI): m/e 367.2 (M+1). Analysis for C ₁₉ H ₁₉ ClF ₄ N ₂ O: Calcd: C, 56.65; H, 4.75; N, 6.95; found: C, 56.82; H, 4.65; N, 6.84. (isolated as the hydrochloride)
305	6-phenyl	2,2,3,3-tetrafluoro propyl	mp 94-98 °C. MS (ACPI): m/e 457.1 (M+1). Analysis for C ₂₆ H ₂₄ F ₄ N ₂ O: Calcd: C, 68.41; H, 5.30; N, 6.14; found: C, 68.18; H, 5.28; N, 6.06. (isolated as the base)

-84-

Example 306N-(2-(6-Carboxy-1H-indol-3-yl)ethyl)-3-(2,2,3,3-tetrafluoropropoxy)benzylamine

Combine N-(2-(6-ethoxycarbonyl-1H-indol-3-yl)ethyl)-N-(2,2,3,3-tetrafluoropropoxybenzyl)amine (1.09 g, 2.4 mmol) and 2N NaOH (4.8 ml) in ethanol (4.8 ml). Heat to reflux. After 2 hours, cool to ambient temperature, evaporate under vacuum to remove the ethanol, and then neutralize with 5N HCl (1.92 ml) to give a solid. Collect the solid by filtration and dry under vacuum to give the title compound as a white powder: mp 186 °C, dec, MS (ACPI): m/e 425.1 (M+1).

10

Example 307N-(2-(6-Carboxy-1H-indol-3-yl)ethyl)-3-(2,2,2-trifluoroethoxy)benzylamine

The method of Example 306 gives the title compound:
mp 232-235 °C. MS (ACPI): m/e 393.2 (M+1).

15

Example 3105-Phenoxy-1H-indole

Combine potassium hydroxide (3 g, 0.054 mol) and phenol (15 g, 0.16 mol), and heat to 110 °C until the potassium hydroxide is dissolved. Cool the mixture to room temperature and add 5-fluoro-2-nitrotoluene (7.75 g, 0.05 mol) in one aliquot. Heat the reaction mixture to 130 °C for 30 min, cool to room temperature, and then pour into 10% NaOH (200 mL). Extract the aqueous solution with ether (2 X 100 mL), combine the organic layers and wash with 10% NaOH (2 X 100 mL), water (2 X 100 mL), dry over Na₂SO₄ and concentrate *in vacuo*. Chromatograph on silica gel eluting with hexanes / ethyl acetate to give 2-nitro-5-phenoxytoluene as a solid: ¹H NMR (300 MHz, CDCl₃) 2.59 (s, 3H), 6.81- 6.85 (m, 2H), 7.06-7.09 (m, 2H), 7.22-7.26 (m, 1H), 7.40-7.45 (m, 2H), 8.03-8.06 (m, 1H).

Combine 2-nitro-5-phenoxytoluene (1.15 g, 5.0 mmol) and tris(dimethylamino)methane (0.87 g, 6.0 mmol) in 10 mL dry toluene and heat to reflux under nitrogen. After 2 hours, cool the reaction mixture to room temperature and

30

-85-

evaporate the toluene under reduced pressure to form a residue. Dissolve the residue in 15 mL EtOAc, mix with Pd/C (10%, 100 mg), stir at room temperature under 1 atmosphere of hydrogen for 1.5 days. Filter off catalyst and concentrate the filtrate. Chromatograph on silica gel eluting with hexanes / EtOAc give the title compound as a solid: ¹H NMR (300MHz, CDCl₃) 6.49-6.50 (m, 1H), 6.93-7.03 (m, 4H), 7.22-7.27 (m, 5H), 8.15 (br, 1H).

- By the method of Example 310 the following compounds were prepared: a) 4-(p-Tolyloxy)-2-methylnitrobenzene: ¹H NMR (300 MHz, CDCl₃) 2.35 (s, 3H), 2.57 (s, 3H), 6.77-6.80 (m, 2H), 6.93-7.03 (m, 2H), 7.18-7.24 (m, 2H), 8.00-8.03 (m, 1H);
- b) 5-p-Tolyloxy-1H-indole: ¹H NMR (300 MHz, CDCl₃) 2.31 (s, 3H), 6.48-6.49 (m, 1H), 6.87-6.96 (m, 3H), 7.07-7.10 (m, 2H), 7.20-7.35 (m, 3H), 8.15 (br, 1H);
- c) 4-(o-Tolyloxy)-2-methylnitrobenzene: ¹H NMR (400 MHz, CDCl₃) 2.16 (s, 3H), 2.57 (s, 3H), 6.50-6.78 (m, 2H), 6.93-7.03 (m, 3H), 7.18-7.35 (m, 1H), 8.00-8.03 (m, 1H);
- d) 5-o-Tolyloxy-1H-indole: ¹H NMR (400 MHz, CDCl₃) 2.31 (s, 3H), 6.45-6.46 (m, 1H), 6.78-6.80 (m, 1H), 6.90-6.00 (m, 2H), 7.01-7.10 (m, 2H), 7.13-7.24 (m, 2H), 7.32-7.34 (m, 1H), 8.11 (br, 1H);
- e) 4-(m-Tolyloxy)-2-methylnitrobenzene: ¹H NMR (300 MHz, CDCl₃) 2.37 (s, 3H), 2.60 (s, 3H), 6.80-6.88 (m, 4H), 7.03-7.06 (m, 1H), 7.27-7.32 (m, 1H), 8.03-8.06 (m, 1H);
- f) 5-m-Tolyloxy-1H-indole: 6.0 g (54%) was obtained (red oil). ¹H NMR (300 MHz, CDCl₃) 2.25 (s, 3H), 6.51-6.52 (m, 1H), 6.76-6.98 (m, 4H), 7.14-7.39 (m, 4H), 8.17 (br, 1H);
- g) 4-(4-Fluorophenoxy)-2-methylnitrobenzene: ¹H NMR (300 MHz, CDCl₃) 2.60 (s, 3H), 6.80-6.82 (m, 2H), 7.03-7.12 (m, 4H), 8.03-8.06 (m, 1H); and
- h) 5-(4-Fluorophenoxy)-1H-indole: 2.68 g (26%) was obtained (red oil). ¹H NMR (300 MHz, CDCl₃) 6.50-6.52 (m, 1H), 6.91-7.01 (m, 5H), 7.24-7.38 (m, 3H), 8.18 (br, 1H). a) 5-p-tolyloxy-1H-indole.

Example 311
2-Oxo-(5-phenoxy-1H-indol-3-yl)acetyl chloride

Combine 5-phenoxy-indole (1.57 g, 7.5 mmol) and anhydrous ether (35 mL) and add oxalyl chloride (1.07 g, 8.25 mmol) in 8 mL ether. A precipitate forms. Stir the reaction over night. Collect the precipitate, dry *in vacuo*, to give the title compound: ¹H NMR (300 MHz, DMSO-d₆) 6.99-7.15 (m, 4H), 7.37-7.42 (m, 2H), 7.60 (d, 1H, *J* = 8.7 Hz), 7.75 (d, 1H, *J* = 2.4 Hz), 8.47 (d, 1H, *J* = 3.2 Hz), 12.49, (br, 1H).

-86-

By the method of Example 311 the following compounds were prepared: a) 2-Oxo-(5-*p*-tolylloxy-1H-indol-3-yl)acetyl chloride;

- b) 2-Oxo-(5-*o*-tolylloxy-1H-indol-3-yl)acetyl chloride: ¹H NMR (300 MHz, CDCl₃) 2.83 (s, 3H), 6.86-6.89 (m, 1H), 7.03-7.16 (m, 5H), 7.26-7.27 (m, 1H), 7.40-7.44 (m, 1H), 7.87 (m, 1H), 8.20-8.32 (m, 2H), 8.90 (br, 1H);
- c) 2-Oxo-(5-*m*-tolylloxy-1H-indol-3-yl)acetyl chloride; and
- d) 2-Oxo-((4-fluorophenoxy)-1H-indol-3-yl)acetyl chloride.

Example 312

10 2-Oxo-2-(5-phenoxy-1H-indol-3-yl)acetamide

Combine 2-oxo-(5-phenoxy-1H-indol-3-yl)acetyl chloride (2.15 g, 7.18 mmol), and ammonium hydroxide (28-30%, 32ml, 680 mmol) and stir for 2 hours. Pour the reaction mixture into 10% (aq.) HCl, extract with dichloromethane, combine the organic layers and dry over Na₂SO₄, evaporate solvent *in vacuo* to give 1.94 g (96%) of title compound: ¹H NMR (300 MHz, CDCl₃) 4.87 (s, 2H), 7.51-7.91 (m, 7H), 8.13-8.24 (m, 3H).

- By the method of Example 312 the following compounds were prepared: a) 2-Oxo-2-(5-*p*-tolylloxy-1H-indol-3-yl)acetamide;
- b) 2-Oxo-2-(5-*o*-tolylloxy-1H-indol-3-yl)acetamide; and
- 20 c) 2-Oxo-2-(5-*m*-tolylloxy-1H-indol-3-yl)acetamide.

Example 314

5-Phenoxytryptamine oxalate

Add 2-oxo-2-(5-phenoxy-1H-indol-3-yl)acetamide (1.9 g, 6.86 mmol) in THF (60 mL) dropwise to a solution of LiAlH₄-THF (1.0 M, 41 mL, 41.0 mmol) in THF at room temperature. Heat the reaction mixture to reflux for 4 hours and cool to room temperature. Quench the reaction mixture with water (6 mL), followed by NaOH (2N, 3 mL). Collect the precipitate by filtration, and wash with ether (3 x 50 mL). Dry the filtrate over Na₂SO₄, concentrate *in vacuo*, purify the residue by flash chromatography (dichloromethane/MeOH/NH₄OH) to obtain 1.0 g (59%) of free amine of the title compound. The oxalic salt of the title compound gives: m.p. 156-157 °C; ¹H NMR (300 MHz, DMSO-d₆) 2.94 (t, 2H, *J* = 7.3 Hz), 3.00 (t, 2H, *J* = 7.3 Hz), 5.00 (br, 2H), 6.83-7.04 (m, 4H), 7.26-7.41 (m, 5H), 11.05 (br, 1H); MS (ELECTROSPRAY), *m/e*: 341.1 (M-1); Anal. Calcd. C₁₈H₁₈N₂O₅: C, 63.15; H, 5.30; N, 8.18. Found: C, 62.97; H, 5.25; N, 8.20.

By the method of Example 314 the following compounds were prepared and isolated as the oxalate, unless otherwise noted: a) 5-*p*-Tolylloxytryptamine: ¹H NMR (300

-87-

MHz, CDCl₃) 2.31 (s, 3H), 2.83 (t, 2H, *J* = 6.4 Hz), 2.98 (t, 2H, *J* = 6.3 Hz), 6.86-6.96 (m, 3H), 7.07-7.10 (m, 3H), 7.24-7.33 (m, 2H), 8.02 (br, 1H) (isolated as the base);

b) 5-o-Tolyloxytryptamine: m.p. 187-188 °C. ¹H NMR (300 MHz, DMSO-d₆) 2.27 (s, 3H), 2.90-3.05 (m, 4H), 6.66-6.68 (m, 1H), 6.76-6.79 (m, 1H), 6.93-6.98 (m, 1H), 7.06-7.16 (m, 2H), 7.24-7.39 (m, 3H), 7.66 (br, 2H), 11.05 (br, 1H); MS (ELECTROSPRAY) m/e: 265.1 (M-1-C₂H₂O₄); Anal. Calcd. C₁₉H₂₂N₂O₅: C, 64.04; H, 5.66; N, 7.86. Found: C, 63.90; H, 5.72; N, 7.83; and

c) 5-m-Tolyloxytryptamine: m.p. 164-165 °C; ¹H NMR (250 MHz, DMSO-d₆) 2.26 (s, 3H), 2.89-3.07 (m, 4H), 4.52 (br, 2H), 6.68-6.72 (m, 2H), 6.82-6.86 (m, 2H), 7.17-7.42 (m, 4H), 11.06 (br, 1H); MS (ELECTROSPRAY) m/e: 265.1 (M-1-C₂H₂O₄).

Example 315

6-Chloro-7-fluoro-1H-indole

Combine boron trichloride (36.0 mL, 1.0 M solution in heptane, 36 mmol) and 1,2-dichloroethane (40 mL) and cool to 5°C. Add dropwise a solution of 2-fluoro-3-chloroaniline (4.36 g, 30.0 mmol) in 20 mL 1,2-dichloroethane. Warm the reaction mixture to room temperature and stir for 30 min. Add to the reaction mixture, chloroacetonitrile (2.71 g, 36.0 mmol), followed by TiCl₄ (6.83 g, 3.84 mL, 36.0 mmol). Heat the reaction mixture to reflux overnight. Cool the reaction mixture to room temperature, add 55.0 mL of 2.5 N HCl, heat to 85 °C for 30 min. Cool to room temperature, extract with dichloromethane (3 x 25 mL), combine the organic layers, wash with brine, dry over Na₂SO₄, concentrate *in vacuo* to give 5.1 g of 1-(2-amino-2-fluoro-3-chlorophenyl)-2-chloroethanone: ¹H NMR (300 MHz, CDCl₃) 4.63 (s, 2H), 6.49 (br, 2H), 6.62-6.69 (m, 1H), 7.36-7.39 (m, 1H).

Dissolve 1-(2-amino-2-fluoro-3-chlorophenyl)-2-chloroethanone in 50 mL (10% water in 1,4-dioxane, v/v) and add NaBH₄ (0.86 g, 22.8 mmol) cautiously at room temperature. Reflux the reaction mixture for about 4 hour, cool to room temperature. Add 35 mL of 1N HCl and stir at room temperature for half an hour, extract with dichloromethane (20 mL X 3), combine the organic layers and wash with H₂O and brine, dry over Na₂SO₄, and concentrate *in vacuo*. Chromatograph on silica gel eluting with EtOAc / hexanes to give the title compound 0.94 g (24%): ¹H NMR (300 MHz, CDCl₃) 6.55-6.58(m, 1H), 7.04-7.10 (m, 1H), 7.22-7.33 (m, 2H), 8.38 (br, 1H).

By the method of Example 315 the following compounds were prepared: a) 5,7-Difluoro-1H-indole : ¹H NMR (300 MHz, CDCl₃) 6.55-6.56 (m, 1H), 6.71-6.78 (m, 1H), 7.01-7.11 (m, 1H), 7.26-7.28 (m, 1H), 8.34 (br, 1H);

b) 6,7-Difluoro-1H-indole: ¹H NMR (300 MHz, CDCl₃) 6.53-6.56 (m, 1H), 6.90-6.99 (m, 1H), 7.22-7.31 (m, 2H), 8.39 (br, 1H);

-88-

c) 5,6,7-Trifluoro-1H-indole: ^1H NMR (300 MHz, CDCl_3) 6.52-6.55 (m, 1H), 7.13-7.20 (m, 1H), 7.26-7.27 (m, 1H), 8.35 (br, 1H); and

d) 4,5,7-Trifluoro-1H-indole: ^1H NMR (300 MHz, $\text{DMSO}-d_6$) 6.68-6.71 (m, 1H), 7.20-7.29 (m, 1H), 7.57-7.59 (m, 1H), 12.07 (br, 1H); MS (electrospray) m/e: 170.0 (M-1).

e) 4,7-Difluoro-1H-indole: ^1H NMR (400 MHz, $\text{dmso}-d_6$): 11.91 (br s, 1H), 7.44 (t, 1H, $J=2.8$ Hz), 6.84-6.90 (m, 1H), 6.69-6.74 (m, 1H), 6.54-6.56 (m, 1H); MS (ES-): m/e 152.0 (M-1).

10

Example 316

3-Formyl-6-chloro-7-fluoro-1H-indole

Add phosphorus oxychloride (0.94 g, 6.16 mmol) to DMF (12 mL, cooled in an ice bath) with vigorous stirring. After about 10 minutes, add 6-chloro-7-fluoro indole (0.93 g, 5.6 mmol) in anhydrous DMF (4 mL), stir at 0°C for 1 hour, warm to room temperature and stir overnight at room temperature (~16 hrs). Treat with 14.0 mL of 2N NaOH (4 eq.) with vigorous stirring. Heat the reaction to 80°C for half an hour then cool. Pour the reaction into cold water with vigorous stirring to give a solid. Collect the solid by filtration and dry overnight in a vacuum oven at room temperature, to give the title compound: ^1H NMR (300 MHz, $\text{CD}_3\text{COCD}_3/\text{CDCl}_3$) 7.09 (t, 1H, $J=7.7$ Hz), 7.83-7.86 (m, 2H), 9.89 (s, 1H). By the method of Example 316 the following compounds were prepared: a) 3-Formyl-5,7-difluoro-1H-indole: ^1H NMR (300 MHz, CD_3COCD_3) 6.98-7.06 (m, 1H), 7.71-7.75 (m, 1H), 8.35 (s, 1H), 10.04 (s, 1H);

b) 3-Formyl-6,7-difluoro-1H-indole: ^1H NMR (300 MHz, CDCl_3) 7.10-7.19 (m, 1H), 7.86-7.88 (m, 1H), 7.98-8.03 (m, 1H), 8.95 (br, 1H), 10.06 (s, 1H);

c) 3-Formyl-5,6,7-trifluoro-1H-indole: ^1H NMR (300 MHz, CD_3COCD_3) 7.87-7.93 (m, 1H), 8.42 (s, 1H), 10.07 (s, 1H); and

d) 3-Formyl-4,5,7-trifluoro-1H-indole: ^1H NMR (300 MHz, $\text{DMSO}-d_6$) 7.46-7.55 (m, 1H), 8.49 (s, 1H), 10.02 (d, 1H, $J=3.7$ Hz), 13.19 (br, 1H).

e) 3-Formyl-4,7-difluoro-1H-indole: ^1H NMR (400 MHz, $\text{dmso}-d_6$): δ 13.03 (br s, 1H), 10.00 (d, 1H, $J=3.2$ Hz), 8.36 (s, 1H), 7.07-7.13 (m, 1H), 6.94-7.00 (m, 1H); MS (APCI): m/e 182.0 (M+1).

f) 3-Formyl-4,5,6,7-tetrafluoro-1H-indole: ^1H NMR (400 MHz, $\text{dmso}-d_6$): δ 13.33 (br s, 1H), 9.94 (d, 1H, $J=4.4$ Hz), 8.49 (s, 1H); MS (ES-): m/e 216.0 (M-1).

Example 3173-(2-Nitrovinyl)-6-chloro-7-fluoro-1H-indole

Combine 3-formyl-6-chloro-7-fluoro-1H-indole (1.00 g, 5.06 mmol), ammonium acetate (292 mg, 3.8 mmol, 0.75 eq.) (dry by treating with toluene and remove the toluene *in vacuo*), and nitromethane (6.17 g, 101.2 mmol, 20 eq.). Warm to 65°C. When the reaction is complete (by TLC), add silica gel and remove the nitromethane *in vacuo*. Load the silica gel on top of short column of silica gel and elute with 25% acetone in hexanes to give, after evaporation, the title compound.

By the method of Example 317 the following compounds were prepared: a) 3-(2-Nitrovinyl)-5,7-difluoro-1H-indole: ¹H NMR (300 MHz, CDCl₃) 6.68-6.81 (m, 1H), 7.16-7.21 (m, 1H), 7.60 (d, 1H, *J* = 13.5 Hz), 7.73 (d, 1H, *J* = 2.7 Hz), 8.18 (d, 1H, *J* = 13.5 Hz), 10.95 (br, 1H);

b) 3-(2-Nitrovinyl)-6,7-difluoro-1H-indole: ¹H NMR (300 MHz, CDCl₃) 6.93-7.00 (m, 1H), 7.30-7.35 (m, 1H), 7.58 (d, 1H, *J* = 13.5 Hz), 7.69 (d, 1H, *J* = 2.9 Hz), 8.10 (d, 1H, *J* = 13.5 Hz), 11.18 (br, 1H); MS (electrospray) *m/e*: 225 (M+1), 223 (M-1);

c) 3-(2-Nitrovinyl)-5,6,7-trifluoro-1H-indole..

d) 3-(2-Nitrovinyl)-4,5,7-trifluoro-1H-indole.

e) 3-(2-Nitrovinyl)-4,7-difluoro-1H-indole: MS (ES-): *m/e* 223.0 (M-1).

f) 3-(2-Nitrovinyl)-4,5,6,7-tetrafluoro-1H-indole: MS(ES-): *m/e* 259.0 (M-1).

Example 3186-Chloro-7-fluorotryptamine

Add dropwise 3-(2-nitrovinyl)-6-chloro-7-fluoro-1H-indole (1.20 g, 5.06 mmol) in anhydrous THF to a solution of lithium aluminum hydride (30.0 mL, 30.0 mmol, 1.0 M solution in THF). Heat to reflux for 2 hour and then cool to room temperature. Quench by carefully adding 1N NaOH to give a suspension. Filter the suspension through celite and rinse repeatedly with ether. Evaporate the filtrate *in vacuo* to give a residue. Chromatograph the residue on silica gel eluting with dichloromethane, methanol, and ammonium hydroxide (10:1:01) to give, after evaporation, the title compound: ¹H NMR (300 MHz, CDCl₃) 2.87 (t, 2H, *J* = 6.6 Hz), 3.02 (t, 2H, *J* = 6.7 Hz), 7.03-7.08 (m, 2H), 7.26-7.29 (m, 1H), 8.51 (br, 1H).

By the method of Example 318 the following compounds were prepared: a) 5,7-Difluorotryptamine: ¹H NMR (300 MHz, CDCl₃) 2.46 (t, 2H, *J* = 6.5 Hz), 3.01 (t, 2H, *J* = 6.4 Hz), 6.69-6.77 (m, 1H), 7.03-7.11 (m, 2H), 8.29 (br, 1H);

b) 6,7-Difluorotryptamine: ¹H NMR (300 MHz, CDCl₃) 2.87 (t, 2H, *J* = 6.6 Hz), 3.02 (t, 2H, *J* = 6.7 Hz), 6.88-6.97 (m, 1H), 7.04 (m, 1H), 7.20-7.25 (m, 1H), 8.64 (br, 1H);

-90-

c) 5,6,7-Trifluorotryptamine: ^1H NMR (300 MHz, CDCl_3) 2.83 (t, 2H, $J = 6.6$ Hz), 3.00 (t, 2H, $J = 6.7$ Hz), 7.08-7.14 (m, 2H), 8.71 (br, 1H); MS (electrospray), m/e : 215.0 (M+1); and


d) 4,5,7-Trifluorotryptamine: ^1H NMR (300 MHz, CDCl_3) 2.93 (t, 2H, $J = 6.6$ Hz), 3.03 (t, 2H, $J = 6.4$ Hz), 6.73-6.82 (m, 1H), 7.02 (s, 1H), 8.58 (br, 1H); MS (electrospray), m/e : 215.0 (M+1), 213.0 (M-1).

f) 4,7-Difluorotryptamine: ^1H NMR (400 MHz, $\text{dms}-d_6$): 11.57 (br s, 1H), 7.19 (s, 1H), 6.80-6.85 (m, 1H), 6.61-6.67 (m, 1H), 2.79 (s, 4H). MS (ES+): m/e 197.0 (M+1) 180.0 (M-NH₂).

g) 4,5,6,7-Tetrafluorotryptamine: ^1H NMR (400 MHz, $\text{dms}-d_6$): δ 7.31 (s, 1H), 2.78 (s, 4H); MS (ES+): m/e 233.0 (M+1) 216.0 (M-16).


Example 319

N-(2-(5-Phenoxy-1H-indol-3-yl)ethyl)-3-phenoxybenzylamine



Combine 5-phenoxytryptamine (0.400 g, 1.59 mmol), 3-phenoxybenzaldehyde (0.377 g, 1.90 mmol) and molecular sieves 4 Å (0.40 g) in methanol (15 mL) and stir for 4h. Filter and wash the molecular sieves several times with MeOH. Add NaBH_4 (61.5 mg, 1.59 mmol) in portions to the filtrate and stir at room temperature for 1h. Remove the MeOH under vacuum to give a residue, dilute the residue with dichloromethane/water, separate the layers, extract the aqueous layer with dichloromethane, combine the organic layers, and dry over Na_2SO_4 . Concentrate *in vacuo* and chromatograph on silica gel eluting with dichloromethane / MeOH the title compound. Form the oxalate salt of the title compound: m.p. 196-198 °C; ^1H NMR (300 MHz, $\text{DMSO}-d_6$) 2.95-3.15 (m, 4H), 4.15 (s, 2H), 6.85-7.46 (m, 18H), 11.06 (br, 1H); MS (ELECTROSPRAY) m/e : 435.3 (M+1); HRMS (ES+) calcd for $\text{C}_{29}\text{H}_{27}\text{N}_2\text{O}_2$ (M+H) 435.2084 found 435.2073.

By the method of Example 319 the following compounds were prepared, isolated as the oxalate except where noted:



No.	Z'	R ₄	Data
320	5- <i>p</i> -tolylloxy	phenyl	m.p. 204-206 °C; ¹ H NMR (250 MHz, DMSO- <i>d</i> ₆) 2.25 (s, 3H), 2.97-3.12 (m, 4H), 4.01 (br, 2H), 4.16 (s, 2H), 6.78-6.84 (m, 3H), 7.00-7.10 (m, 10H), 7.13-7.43 (m, 4H), 11.05 (br, 1H); MS (ELECTROSPRAY) m/e: 449.1 (M+1-C ₂ H ₂ O ₄); Analysis calcd: C ₃₂ H ₃₀ N ₂ O ₆ : C, 71.36; H, 5.61; N, 5.20. Found: C, 71.22; H, 5.59; N, 5.28
321	5- <i>o</i> -tolylloxy	phenyl	m.p. 191-192 °C; ¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) 2.28 (s, 3H), 2.99-3.15 (m, 4H), 4.17 (s, 2H), 6.63-6.66 (m, 1H), 6.756.79 (m, 1H), 6.92-7.42 (m, 15H), 9.50 (br, 2H), 11.05 (br, 1H); MS(ELECTROSPRAY) m/e: 449.1 (M+1-C ₂ H ₂ O ₄); Anal. calcd. C ₃₂ H ₃₀ N ₂ O ₆ : C, 71.36; H, 5.61; N, 5.20. Found C, 71.11; H, 5.59; N, 5.18
322	5- <i>m</i> -tolylloxy	phenyl	m.p. 174-175 °C. ¹ H NMR (250 MHz, DMSO- <i>d</i> ₆) 2.51(s, 3H), 3.00-3.13 (m, 4H), 4.15 (s, 2H), 6.81-7.03 (m, 7H), 7.11-7.42 (m, 11H), 11.05 (br, 1H): MS (ELECTROSPRAY) m/e: 449.1 (M+1-C ₂ H ₂ O ₄)
323	6-chloro-7-fluoro	2,2,2-trifluoro ethyl	mp 186-187 °C. ¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) 3.13 (s, 4H), 4.15 s, 2H), 4.78 (q, 2H, <i>J</i> = 8.7 Hz), 7.07-7.12 (m, 2H), 7.21-7.24 (m, 1H), 7.37-7.45 (m, 4H), 9.44 (br, 1H), 11.72(br, 1H): ms (electrospray) m/e: 401.2 (M+1-HCl), 399.2 (M-1-HCl); Anal. calcd. C ₁₉ H ₁₇ ClF ₄ N ₂ O·HCl: C, 52.19; H, 4.15; N, 6.41. Found: C, 52.15; H, 4.14; N, 6.38 (isolated as the hydrochloride)
324	6-chloro-7-fluoro	2,2,3,3-tetrafluoro propyl	mp. 155-156 °C; ¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) 3.13 (s, 4H), 4.16 (s, 2H), 4.61 (t, 2H, <i>J</i> = 13.5 Hz), 6.70 (tt, 1H, <i>J</i> = 51.9 Hz, <i>J</i> = 5.5 Hz), 7.08-7.10 (m, 2H), 7.11-7.12 (m, 1H), 7.21-7.45 (m, 4H), 9.41 (br, 1H), 11.72(br, 1H); MS (electrospray) m/e: 433.2 (M+1-HCl), 431.2 (M-1-HCl); Anal. calcd. C ₂₀ H ₁₈ ClF ₅ N ₂ O·HCl: C, 51.19; H, 4.08; N, 5.97. Found: C, 51.27; H, 4.10; N, 6.07 (isolated as the hydrochloride)
325	5,7-difluoro	2,2,2-trifluoro ethyl	m.p.: 179-180 °C; ¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) 3.11 (s, 4H), 4.16 (s, 2H), 4.77 (q, 2H, <i>J</i> = 8.7 Hz), 6.93-6.97 (m, 1H), 7.00-7.14 (m, 1H), 7.21-7.43 (m, 5H), 9.41 (br, 1H), 11.61 (br, 1H); ms (electrospray) m/e: 385.2 (M+1-HCl), 383.0 (M-1-HCl); Anal. calcd. C ₁₉ H ₁₇ F ₅ N ₂ O·HCl·0.1H ₂ O: C, 54.00; H, 4.34; N, 6.63. Found: C, 53.71; H, 4.24; N, 6.70 (isolated as the hydrochloride)

326	5,7-difluoro	2,2,3,3-tetrafluoro propyl	mp. 109-110 °C; ¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) 2.71-2.84 (m, 4H), 3.71 (s, 2H), 4.53 (t, 2H, <i>J</i> = 13.5 Hz), 6.67 (tt, 1H, <i>J</i> = 51.9 Hz, <i>J</i> = 5.5 Hz), 6.87-7.02 (m, 4H), 7.12-7.28 (m, 3H), 11.40 (br, 1H); MS (electrospray) <i>m/e</i> : 417.0 (M+1), 415.0 (M-1); Anal. calcd. C ₂₀ H ₁₈ F ₆ N ₂ O·0.1H ₂ O: C, 57.45; H, 4.39; N, 6.70. Found: C, 57.24; H, 4.08; N, 6.68
327	6,7-difluoro	2,2,2-trifluoro ethyl	m.p.: 164-165 °C; ¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) 3.13 (s, 4H), 4.16 (s, 2H), 4.77 (q, 2H, <i>J</i> = 9.1 Hz), 7.00-7.13 (m, 2H), 7.20-7.23 (m, 1H), 7.33-7.43 (m, 4H), 9.36 (br, 1H), 11.57 (br, 1H); MS (electrospray) <i>m/e</i> : 385.2 (M+1-HCl), 383.3 (M-1-HCl); Anal. calcd. C ₁₉ H ₁₇ F ₅ N ₂ O·HCl: C, 54.23; H, 4.31; N, 6.66. Found: C, 53.86; H, 4.28; N, 6.58 (isolated as the hydrochloride)
328	6,7-difluoro	2,2,3,3-tetrafluoro propyl	m.p.: 214-215 °C; ¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) 3.02-3.17 (m, 4H), 4.16 (s, 2H), 4.59 (t, 2H, <i>J</i> = 13.5 Hz), 6.68 (tt, 1H, <i>J</i> = 51.9 Hz, <i>J</i> = 5.5 Hz), 7.00-7.17 (m, 5H), 7.21-7.42 (m, 4H), 11.65 (br, 1H); MS (electrospray) <i>m/e</i> : 417.0 (M+1-C ₄ H ₄ O ₄), 415.0 (M-1-C ₄ H ₄ O ₄); Anal. calcd. C ₂₀ H ₁₈ F ₆ N ₂ O·C ₄ H ₄ O ₄ ·0.9H ₂ O: C, 52.54; H, 4.37; N, 5.11. Found: C, 52.14; H, 3.95; N, 5.49 (isolated as the maleate)
329	5,6,7-trifluoro	2,2,2-trifluoro ethyl	m.p.: 111-112 °C; ¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) 2.72-2.81 (m, 4H), 3.71 (s, 2H), 4.68 (q, 2H, <i>J</i> = 8.8 Hz), 6.87-7.00 (m, 3H), 7.22-7.40 (m, 3H), 11.58 (br, 1H); MS (electrospray) <i>m/e</i> : 403.1 (M+1), 401.2 (M-1). Anal. calcd. C ₁₉ H ₁₆ F ₆ N ₂ O: C, 56.72; H, 4.01; N, 6.96. Found: C, 56.61; H, 3.92; N, 6.96 (isolated as the base)
330	5,6,7-trifluoro	2,2,3,3-tetrafluoro propyl	m.p.: 223-224 °C; ¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) 3.11 (s, 4H), 4.15 (s, 2H), 4.61 (t, 2H, <i>J</i> = 13.5 Hz), 6.70 (tt, 1H, <i>J</i> = 51.9 Hz, <i>J</i> = 5.5 Hz), 7.08-7.12 (m, 1H), 7.19-7.25 (m, 1H), 7.36-7.43 (m, 3H), 7.52-7.58 (m, 1H), 9.50 (br, 1H), 11.78 (br, 1H); MS (electrospray) <i>m/e</i> : 435.1 (M+1-HCl), 433.1 (M-1-HCl); Anal. calcd. C ₂₀ H ₁₈ F ₆ N ₂ O·HCl·0.1H ₂ O: C, 50.83; H, 3.88; N, 5.93. Found: C, 50.60; H, 3.74; N, 6.07 (isolated as the hydrochloride)

331	4,5,7-trifluoro	2,2,2-trifluoro ethyl	m.p.: 243-244 °C; ¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) 3.16-3.21 (m, 4H), 4.18 (s, 2H), 4.75 (q, 2H, <i>J</i> = 8.8 Hz), 7.11-7.25 (m, 3H), 7.39-7.45 (m, 3H), 9.37 (br, 1H), 11.90 (br, 1H); MS (electrospray) <i>m/e</i> : 403.1 (M+1-HCl), 401.0 (M-1-HCl); Anal. calcd. C ₁₉ H ₁₆ F ₆ N ₂ O·HCl: C, 52.00; H, 3.91; N, 6.38. Found: C, 51.83; H, 3.62; N, 6.55 (isolated as the hydrochloride)
332	4,5,7-trifluoro	2,2,3,3-tetrafluoro propyl	m.p.: 261-262 °C; ¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) 3.18 (s, 4H), 4.17 (s, 2H), 4.61 (t, 2H, <i>J</i> = 13.5 Hz), 6.69 (tt, 1H, <i>J</i> = 51.9 Hz, <i>J</i> = 5.5 Hz), 7.09-7.13 (m, 1H), 7.17-7.26 (m, 2H), 7.32-7.42 (m, 3H), 9.37 (br, 1H), 11.92 (br, 1H); MS (electrospray) <i>m/e</i> : 435.1 (M+1-HCl), 433.1 (M-1-HCl); Anal. calcd. C ₂₀ H ₁₇ F ₇ N ₂ O·HCl, C, 51.02; H, 3.85; N, 5.95. Found: C, 50.62; H, 3.79; N, 6.00 (isolated as the hydrochloride)
333	7-cyano	2,2,2-trifluoro ethyl	mp. 241-242 °C; ¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) 3.15 (s, 4H), 4.17 (s, 2H), 4.78 (q, 2H, <i>J</i> = 8.7 Hz), 7.10-7.22 (m, 3H), 7.33-7.43 (m, 3H), 7.60-7.62 (m, 1H), 7.95-7.97 (m, 1H), 9.29 (br, 2H), 11.90 (br, 1H); MS (electrospray) <i>m/e</i> : 374.2 (M+1-HCl), 372.0 (M-1-HCl); Anal. calcd. C ₂₀ H ₁₈ F ₃ N ₃ O·HCl·0.2 H ₂ O: C, 58.10; H, 4.73; N, 10.16. Found: C, 57.91; H, 4.56; N, 10.08.
334	7-cyano	2,2,3,3-tetrafluoro ethyl	mp. 212-213 °C; ¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) 3.16 (s, 4H), 4.16 (s, 2H), 4.61 (t, 2H, <i>J</i> = 13.6 Hz), 6.69 (tt, 1H, <i>J</i> = 51.9 Hz, <i>J</i> = 5.5 Hz), 7.09-7.22 (m, 3H), 7.33-7.43 (m, 3H), 7.60-7.63 (m, 1H), 7.96-7.98 (m, 1H), 9.34 (br, 2H), 11.92 (br, 1H); MS (electrospray) <i>m/e</i> : 406.2 (M+1-HCl), 404.0 (M-1-HCl); Anal. calcd. C ₂₁ H ₁₉ F ₄ N ₃ O·HCl, C, 57.08; H, 4.56; N, 9.51. Found: C, 57.12; H, 4.61; N, 9.53.

Example 3352-Fluoro-3-phenoxybenzaldehyde

Cool a solution of 2,2,6,6-tetramethylpiperidine (5.1 mL, 30.0 mmol) in THF (40 mL) to -78 °C. Add dropwise *n*-Butyllithium (18.7 mL, 30.0 mmol, 1.6M in hexanes) and stir for 10 min at -78 °C. Add dropwise 2-fluorophenyl phenyl ether (4.7 g, 25.0 mmol), stir 2 h at -78 °C. Add *N,N*-dimethylformamide (2.3 mL, 30.0 mmol) dropwise over 15 min. Stir the resulting mixture for 3 h at -78 °C and allow to warm to ambient temperature

-94-

over 16 h. Quench the reaction mixture with water (50 mL), extract with ethyl acetate, dry over Na₂SO₄, filter and concentrate under reduced pressure to give an oil. Crystallize the oil with hexanes to give a solid, collect and recrystallize from hexanes/ethyl acetate/methylene chloride to give the title compound: mp 75-77 °C; MS(m/e): 216 (M⁺);

5 Calculated for C₁₃H₉FO₂: Calcd: C, 72.22; H, 4.20. Found: C, 72.41; H, 4.23.

Purification of the mother liquors by silica gel chromatography (2-3% ethyl acetate/hexanes) gives an additional amount of the title compound: MS(m/e): 216 (M⁺).

By the method of Example 335 the following compound was prepared: a) 6-Fluoro-3-phenoxybenzaldehyde: MS(m/e): 216 (M⁺).

10

Example 336

3-Ethoxybenzaldehyde

Combine 3-hydroxybenzaldehyde (5.6 g, 46 mmol) and 1-iodoethane (10.7 g, 69 mmol) in DMSO (25 mL) and warm to 80°C. Treat with of cesium carbonate
15 (22.4 g, 69 mmol) in portions. During the addition the temperature begin to rise so the bath is removed. Stir the reaction at 80°C for 1 hour, pour into 200 mL brine and extract twice with 150 mL diethyl ether. Wash the combine extracts twice with 200 mL brine, dry over MgSO₄ and concentrate under vacuum to give an oil. Purification by chromatography (SiO₂; 2.5% EtOAc in hexanes) affords 5.73 g (38
20 mmol; 83%) of the title compound as an oil: ¹H NMR (CDCl₃) 9.94 (s, 1H), 7.42-7.41 (m, 2H), 7.36-7.35 (m, 1H), 7.16-7.13 (m, 1H), 4.10-4.04 (q, 2H), 1.64-1.40 (t, 3H).

By the method of Example 336 the following compounds were prepared: a) 3-Propoxybenzaldehyde: ¹H NMR (CDCl₃) 9.95 (s, 1H), 7.43-7.41 (m, 2H), 7.37-7.36 (m,
25 1H), 7.17-7.14 (m, 1H), 9.98-3.95 (t, 2H), 1.84-1.79 (m, 2H), 1.05-1.02 (t, 3H).

Example 337

p-Toluene-3-(2,2,3,3-tetrafluoropropoxy)tosylate

Add pyridine (1.9 L) (dried over molecular sieves 4 Å) to a round-bottomed flask
30 (5 L), under inert atmosphere and equipped with a mechanical agitator and add 2,2,3,3-tetrafluoro-1-propanol (604.5 g, 4.58 mol). Cool the mixture to 0°C with an ice-bath. Add p-toluenesulfonyl chloride (960 g, 5.04 mol) over 20 min in 4 portions to the reaction mixture and stir. After 20 min., cooling on an ice-bath), a precipitate is formed. Stir the reaction mixture for 1 h at 0°C and 2 h at 20°C. Pour the reaction mixture, with agitation,
35 over an ice-water mixture (1.44 L) and leave overnight (18 h) at 20°C. The crude tosylate derivative separates from the aqueous mixture as an oily material (1.34 kg) containing 14% w/w pyridine which corresponds to 1.15 kg of the tosylate (87.8%). The crude

-95-

material is carried to the next reaction step without further purification: ^1H -RMN is consistent.

Example 338

5 3-(2,2,3,3,3-Pentafluoropropoxy)benzaldehyde

Combine 3-hydroxybenzaldehyde (137.6 g, 1.127 mol), p-toluene-3-(2,2,3,3,3-pentafluoropropoxy) tosylate (243 g, 0.799 mol), potassium carbonate (220 g, 1.597 mol) and dimethylformamide (2451 mL) in a double-wall 4 L reactor equipped with a reflux condenser and a mechanical agitator, and heat at 110°C for 46.5 h under argon atmosphere. Cool the reaction mixture to room temperature and filter through a bed of 400 g of silica gel. Elute silica gel bed with 2.451 mL of ethyl acetate. Pour the combined organic layers over 7.3 L of ice-water. Add 10 N sodium hydroxide (500 mL) to this mixture and stir for 1 h. Separate the aqueous phase and extract with ethyl acetate (1000 mL). Pool the organic phase, wash with water (1000 mL) and brine (750 mL). Evaporation of the organic solvents under reduced pressure provides 159.79 g of a brown oily material containing the crude title compound. Purification by fractional distillation (two successive cycles) under reduced pressure (2 mm Hg) using a distillation apparatus equipped with a 30 cm length adiabatic column to gives a fraction of 52.4 g of the expected product (96.2% area by HPLC).

20


Example 339

3-(3,3,3-Trifluoropropoxy)benzaldehyde


Combine 3-hydroxybenzaldehyde (130.2 g, 1.066 mol), 3,3,3-trifluoropropoxy tosylate (143 g, 0.533 mol), potassium carbonate (147.35 g, 1,066 mol) and absolute ethanol (1430 mL) in a three-necked round-bottomed flask equipped with a reflux condenser and a magnetic stirred and reflux for 4 h under argon atmosphere. Concentrate the reaction mixture under reduced pressure. Pour the concentrated mixture over 1N sodium hydroxide (2145 mL), stir for 30 min and extract with dichloromethane (2145 mL). Decant the organic layer wash with 1N sodium hydroxide (2145 mL). After separation, wash the organic layer successively twice by 1 L water (pH aqueous phase = 7), dry over 30 g magnesium sulfate, evaporate the dichloromethane organic layer to dryness under reduced pressure to yield 55.4 g of the title compound (0.254 mol, 47.6% yield) as a slightly yellow oily material.

30


-96-

Example 340N-(2-(6-Fluoro-1H-indol-3-yl)ethyl)-2-fluoro-3-phenoxy-benzylamine

- Combine 6-fluorotryptamine (419 mg, 2.35 mmol) and 2-fluoro-3-phenoxybenzaldehyde (610 mg, 2.82 mmol) in absolute ethanol (6 mL). Heat the mixture to 65 °C to give a homogeneous solution. Add 3 Å molecular sieves (400 mg) to the mixture and heat to reflux temperatures for 5 h. Cool the reaction mixture to ambient temperature and add sodium borohydride (267 mg, 7.1 mmol). Stir the mixture for 18 h at ambient temperature. Cool the reaction mixture on a water bath, quench with acetone, dilute with ethanol and acetone, and filter the molecular sieves. Concentrate the filtrate under reduced pressure, dilute with 1N NaOH, extract with ethyl acetate, wash with brine, dry (Na₂SO₄), filter and concentrate under reduced pressure to give 1.0 g of an oil. Chromatograph on silica gel eluting with 1%, 4% 2N ammonia in methanol/methylene chloride gives a clear colorless oil. Formation of the hydrochloride salt in ethyl acetate/methanol gives the hydrochloride of the title compound: mp 173-174.5 °C; MS(m/e): 379 (M+1), 377 (M-1); Calculated for C₂₃H₂₀F₂N₂O·HCl: Calcd: C, 66.59; H, 5.10; N, 6.75. Found: C, 66.50; H, 5.09; N, 6.73.

Example 34120 N-(2-(6-Fluoro-1H-indol-3-yl)ethyl)-6-fluoro-3-phenoxy-benzylamine

- The method of Example 340 gives the hydrochloride of the title compound: mp 183.5 - 185.5 °C; MS(m/e): 379 (M+1), 377 (M-1); Calculated for C₂₃H₂₀F₂N₂O·HCl: Calcd: C, 66.59; H, 5.10; N, 6.75. Found: C, 66.54; H, 5.11; N, 6.68.
- 25 By the method of Example 340 the following compounds were prepared, isolated as the hydrochloride except where noted:




No.	Z'	R ₄	Data
342	5-methoxy	ethyl	ISMS 325 (M+1); Analysis for C ₂₀ H ₂₅ ClN ₂ O ₂ 0.2EtOH 0.1H ₂ O: calcd: C, 65.88; H, 7.16; N, 9.53; found: C, 65.90; H, 6.97; N, 7.16; ¹ H NMR (DMSO-d ₆) 10.85 (s, 1H), 9.43 (bs, 2H), 7.42-7.22 (m, 4H), 7.18-7.10 (m, 2H), 7.05-7.0 (m, 1H), 6.32-6.15 (m, 1H), 4.3-4.15 (m, 2H), 4.15-4.05 (q, 2H), 3.85 (s, 3H), 3.15 (s, 4H), 1.45-1.35 (t, 3H)
343	5-methoxy	propyl	ISMS 339 (M+1); Analysis for C ₂₁ H ₂₇ ClN ₂ O ₂ : calcd: C, 67.28; H, 7.23; N, 7.47; found: C, 67.28; H, 7.30; N, 7.13; ¹ H NMR (DMSO-d ₆) 10.85 (s, 1H), 9.43 (bs, 2H), 7.35-7.15 (m, 4H), 7.1-7.05 (m, 2H), 7.0-6.92 (m, 1H), 6.7-6.6 (m, 1H), 4.3-4.16 (m, 1H), 4.15-4.05 (q, 2H), 3.85 (s, 3H), 3.15 (s, 4H), 1.45-1.35 (t, 3H)
344	5-fluoro	2,2,2-trifluoro ethyl	ISMS 367 (M+1); Analysis for C ₁₉ H ₁₉ ClF ₄ N ₂ O: calcd: C, 56.65; H, 4.75; N, 6.95; found: C, 56.37; H, 4.83; N, 6.81 (base)
345	5-methoxy	2,2,2-trifluoro ethyl	Analysis for C ₂₀ H ₂₂ ClF ₃ N ₂ O ₂ : calcd: C, 57.91; H, 5.34; N, 6.75; found: C, 57.72; H, 5.17; N, 6.61; ISMS 379 (M+1)
346	5-fluoro	2,2,3,3,3-pentafluoro propyl	ISMS 417 (M+1); Analysis for C ₂₀ H ₁₈ F ₆ N ₂ O C ₂ H ₂ O ₄ : calcd: C, 51.18; H, 3.98; N, 5.53; found: C, 51.18; H, 3.91; N, 5.51 (isolated as the oxalate)
347	5-methoxy	2,2,3,3,3-pentafluoro propyl	ISMS 429 (M+1); Analysis for C ₂₁ H ₂₁ F ₅ N ₂ O ₂ 1.2C ₂ H ₂ O ₄ 0.8H ₂ O: calcd: C, 51.02; H, 4.57; N, 5.09; found: C, 50.64; H, 4.23; N, 5.15 (isolated as the oxalate)
348	5-methoxy	2,2,3,3-tetrafluoro propyl	ISMS 411 (M+1) Analysis for C ₂₁ H ₂₂ F ₄ N ₂ O ₂ C ₂ H ₂ O ₄ 0.1H ₂ O: calcd: C, 55.0; H, 4.86; N, 5.58; found: C, 54.74; H, 4.74; N, 5.58 (isolated as the oxalate)
349	5-methoxy	3,3,3-trifluoro propyl	Analysis for C ₂₁ H ₂₃ F ₃ N ₂ O ₂ HCl: calcd: C, 58.81; H, 5.64; N, 6.53; found: C, 58.42; H, 5.44; N, 6.51; ISMS 393 (M+1)
350	5-fluoro	2,2,3,3-tetrafluoro propyl	Analysis for C ₂₀ H ₂₀ F ₄ N ₂ O HCl: calcd: C, 57.63; H, 5.08; N, 6.72; found: C, 57.49; H, 5.04; N, 6.76; ISMS 381 (M+1)
351	4-chloro-5-methoxy	2,2,2-trifluoro ethyl	Analysis for C ₂₀ H ₂₀ ClF ₃ N ₂ O ₂ HCl: calcd: C, 53.47; H, 4.71; N, 6.24; found: C, 53.33; H, 4.65; N, 6.21; ISMS 413 (M+1)
352	4-chloro-5-methoxy	2,2,3,3-tetrafluoro propyl	Analysis for C ₂₁ H ₂₁ ClF ₄ N ₂ O ₂ HCl: calcd: C, 52.40; H, 4.61; N, 5.82; found: C, 52.25; H, 4.50; N, 5.80; ISMS 445 (M+1)

-98-


353	4-chloro-5-methoxy	3,3,3-trifluoropropyl	Analysis for $C_{21}H_{22}ClF_3N_2O_2$ HCl: calcd: C, 54.4; H, 5.00; N, 6.05; found: C, 54.18; H, 4.86; N, 6.06; ISMS 427 (M+1)
-----	--------------------	-----------------------	---

By the method of Example 340 the following compounds were prepared, isolated as the hydrochloride except where noted:




No.	Z'	R ₄	Data
360	3-chloro	2,2,2-trifluoroethyl	ISMS 344 (M+1); Analysis for $C_{17}H_{18}ClF_3NO$ calcd: C, 53.70; H, 4.77; N, 3.68; found: C, 53.61; H, 4.96; N, 3.66
361	3-trifluoromethyl	2,2,2-trifluoroethyl	ISMS 378 (M+1); Analysis for $C_{20}H_{19}F_6NO_5$: calcd: C, 51.40; H, 4.10; N, 3.0; found: C, 51.26; H, 4.06; N, 3.07 (isolated as the oxalate)
362	3-chloro	2,2,3,3,3-pentafluoropropyl	ISMS 394 (M+1); Analysis for $C_{18}H_{18}ClF_5NO$: calcd: C, 50.25; H, 4.22; N, 3.26; found: C, 50.38; H, 4.03; N, 3.45
363	3-trifluoromethyl	2,2,3,3,3-pentafluoropropyl	ISMS 428 (M+1); Analysis for $C_{19}H_{17}F_8NO$ $C_2H_2O_4$: calcd: C, 48.75; H, 3.70; N, 2.70; found: C, 48.76; H, 3.67; N, 2.79 (isolated as the oxalate)
364	3-chloro	2,2,3,3-tetrafluoropropyl	Analysis for $C_{18}H_{18}ClF_4NO$ $C_2H_2O_4$: calcd: C, 51.57; H, 4.33; N, 3.01; found: C, 51.92; H, 4.29; N, 3.08; ISMS 376 (M+1)
365	3-trifluoromethyl	2,2,3,3-tetrafluoropropyl	Analysis for $C_{19}H_{18}F_7NO$ $C_2H_2O_4$: calcd: C, 50.51; H, 4.04; N, 2.81; found: C, 50.48; H, 4.02; N, 2.85; ISMS 410 (M+1) (isolated as the oxalate)
366	3-trifluoromethyl	3,3,3-trifluoropropyl	Analysis for $C_{19}H_{19}F_6NO$ HCl: calcd: C, 53.34; H, 4.71; N, 3.27; found: C, 53.23; H, 4.73; N, 3.28; ISMS 392 (M+1)
367	3-chloro	3,3,3-trifluoropropyl	Analysis for $C_{18}H_{19}ClF_3NO$ HCl: calcd: C, 54.84; H, 5.11; N, 3.55; found: C, 54.74; H, 5.02; N, 3.11; ISMS 358 (M+1)

-99-

Example 370N-(2-(6-Fluoro-1H-indol-3-yl)ethyl)-3-(2,2,2-trifluoroethoxy)benzylamine

- Combine 6-fluorotryptamine oxalate (350 mg, 1.3 mmol), N,N-diisopropylethylamine (506 mg, 3.9 mmol), 3-(2,2,2-trifluoroethoxy)benzaldehyde (266 mg, 1.3 mmol), and 4Å molecular sieves (4g) in EtOH (30 mL) and reflux for 7 hours. Decant the liquid into a separate flask and treat with NaBH₄ (148 mg, 3.9 mmol). Stir 1 hour, concentrate the mixture in vacuo to give a residue. Partition the residue between 25 mL 5 N NaOH and 25 mL dichloromethane. Extract the aqueous layer with 25 mL dichloromethane, combine the organic layers, dry over MgSO₄, and concentrate to approximately a 20 mL volume. Chromatograph on silica gel eluting with 1% MeOH in CHCl₃ mixed with conc. NH₄OH to give the title compound. Combine an EtOAc solution of the title compound with an EtOAc solution of one equivalent of oxalic acid to give a solid, filter, and dry under vacuum to give the oxalate salt of the title compound: ISMS 367 (M+1); Analysis for C₁₉H₁₉ClF₄N₂O: calcd: C, 55.27; H, 4.42; N, 6.14; found: C, 55.17; H, 4.38; N, 6.09.

By the method of Example 370 the following compounds were prepared, isolated as the hydrochloride except where noted:



20

No.	Z'	R ₄	Data
372	5-fluoro	2,2,3,3-tetrafluoro propyl	ISMS 399 (M+1); Analysis for C ₁₉ H ₁₇ F ₈ NO C ₂ H ₂ O ₄ H ₂ O: calcd: C, 53.51; H, 4.41; N, 5.67; found: C, 53.12; H, 4.21; N, 5.63 (isolated as the oxalate)
373	6-fluoro	2,2,3,3,3-pentafluoro propyl	Analysis for C ₂₀ H ₁₈ F ₆ N ₂ O HCl: calcd: C, 53.05; H, 4.23; N, 6.19; found: C, 52.88; H, 4.05; N, 6.12; ISMS 417 (M+1)
374	6-chloro-5-methoxy	2,2,2-trifluoro ethyl	Analysis for C ₂₀ H ₂₀ ClF ₃ N ₂ O ₂ HCl: calcd: C, 53.47; H, 4.71; N, 6.24; found: C, 53.65; H, 4.85; N, 6.45; ISMS 413 (M+1) (Form the salt in 50 mL 50/50 THF/EtOH using polyvinyl pyridine hydrochloride)

-100-

No.	Z'	R ₄	Data
375	6-chloro-5-methoxy	2,2,3,3-tetrafluoro propyl	Analysis for C ₂₁ H ₂₁ ClF ₄ N ₂ O ₂ HCl: calcd: C, 52.40; H, 4.61; N, 5.82; found: C, 52.15; H, 4.51; N, 5.69; ISMS 445 (M+1)
376	6-fluoro	2,2,3,3-tetrafluoro propyl	Analysis for C ₂₀ H ₁₉ F ₅ N ₂ O HCl: calcd: C, 55.24; H, 4.64; N, 6.44; found: C, 55.06; H, 4.63; N, 6.44; ISMS 399 (M+1)
377	6-fluoro	3,3,3-trifluoro propyl	Analysis for C ₂₀ H ₂₀ F ₄ N ₂ O HCl: calcd: C, 54.83; H, 5.11; N, 3.55; found: C, 54.74; H, 5.02; N, 3.11; ISMS 381 (M+1)
378	5-trifluoro methoxy	2,2,3,3,3-pentafluoro propyl	Analysis for C ₂₁ H ₁₈ F ₈ N ₂ O ₂ HCl: calcd: C, 48.62; H, 3.69; N, 5.40; found: C, 48.55; H, 3.48; N, 5.33; ISMS 483 (M+1)
379	5-trifluoro methoxy	2,2,3,3-tetrafluoro propyl	Analysis for C ₂₁ H ₁₉ F ₇ N ₂ O ₂ HCl: calcd: C, 50.36; H, 4.02; N, 5.59; found: C, 50.27; H, 3.92; N, 5.63; ISMS 465 (M+1)
380	5-trifluoro methoxy	2,2,2-trifluoro ethyl	Analysis for C ₂₀ H ₁₈ F ₆ N ₂ O ₂ HCl: calcd: C, 51.24; H, 4.08; N, 5.98; found: C, 51.33; H, 4.09; N, 6.26; ISMS 433 (M+1)

Example 381N-t-Butoxycarbonyl-2-(5-m-tolyloxy-1H-indol-3-yl)ethylamine

The method of Example 20 gives the title compound: ¹H NMR (300 MHz, CDCl₃)


- 5 1.41 (s, 9H), 2.30 (s, 3H), 2.89 (t, 2H, *J* = 6.7 Hz), 3.41 (m, 2H), 6.74-6.85 (m, 3H), 6.93-6.99 (m, 1H), 7.07-7.35 (m, 4H), 8.05 (br, 1H).

Example 382N-Methyl-2-(5-m-tolyl)tryptamine

- 10 The method of Example 21 gives the title compound and formation of the oxalate salt gave: m.p. 182-183 °C; ¹H NMR (250 MHz, DMSO-d₆) 2.26 (s, 3H), 2.59 (s, 3H), 2.98-3.18 (m, 4H), 6.68-6.72 (m, 2H), 6.82-6.86 (m, 2H), 7.17-7.22 (m, 1H), 7.29-7.42 (m, 3H), 11.06 (br, 1H); MS (ELECTROSPRAY) *m/e*: 281.2 (M+1-C₂H₂O₄); Anal. calcd. C₂₀H₂₂N₂O₅: C, 64.85; H, 5.99; N, 7.56. Found: C, 65.01; H, 5.74; N, 7.71.

15

-101-

Example 383N-Methyl-N-(2-(5-*m*-tolxyloxy-1H-indol-3-yl)ethyl)-3-phenoxybenzylamine

The method of Example 301 gives the title compound and formation of the oxalate salt gave: m.p. 142-144 °C; ¹H NMR (250 MHz, DMSO-d₆) 2.24 (s, 3H), 2.634 (s, 3H), 3.01-3.12 (m, 4H), 3.92 (br, 2H), 4.16 (s, 2H), 6.65-6.70 (m, 2H), 6.81-6.84 (m, 2H), 6.99-7.03 (m, 3H), 7.12-7.26 (m, 6H), 7.34-7.43 (m, 4H), 11.00 (br, 1H); MS (ELECTROSPRAY) m/e: 463.4 (M+1-C₂H₂O₄); Anal. calcd. C₃₃H₃₂N₂O₆: C, 71.72; H, 5.84; N, 5.07. Found: C, 71.44; H, 5.89; N, 4.99.

10

Example 3845-Nitrotryptamine

Warm mixture of 5-nitroindole (10 g, 62 mmol) and 200 mL glacial acetic acid to 70°C and treat with Eschenmoser's salt (12 g, 65 mmol). Concentrate after 1 hour the reaction under vacuum to dryness. Mix the residue with 200 mL toluene, reconcentrate to dryness then partition between 200 mL concentrated ammonium hydroxide and 200 mL EtOAc. When all solids dissolved, Separate the layers and extract the aqueous layer 200 mL EtOAc. Dry the combined organic layer over MgSO₄ and concentrate to give N,N-dimethyl-5-nitrotryptamine as a solid.

Dissolve the N,N-dimethyl-5-nitrotryptamine obtained above in 200 mL dry DMSO, treat with iodomethane (7.7 mL, 17.5 g, 124 mmol), and stir for 1 hour at ambient temperature. Add KCN (40 g, 621 mmol) and 18-crown-6 (0.5 g). Warm the reaction to 110°C for 45 minutes, cool, poured onto ice then saturate with NaCl. Extract the quenched reaction mixture with EtOAc, combine the extracts, and wash 3 times with brine. Dry over MgSO₄ and concentrate under vacuum. chromatograph on silica gel eluting with 1% MeOH in CHCl₃ to give (5-nitro-1H-indol-3-yl)acetonitrile as a solid: FDMS 201 (M+); Analysis for C₁₀H₇N₃O₂: calcd: C, 59.70; H, 3.51; N, 20.89; found: C, 59.32; H, 3.52; N, 20.56.

Dissolve (5-nitro-1H-indol-3-yl)-acetonitrile (9 g, 44.7 mmol) in 250 mL dry THF and treat with 90 mL 1 M BH₃ in THF at ambient temperature. Stir overnight and quench the reaction cautiously by the dropwise addition of 10 mL water. Concentrate to dryness under vacuum and partition the residue between 5 N


30

-102-

HCl and EtOAc. Extract the aqueous layer with EtOAc and combine with the original EtOAc layer. Treat the aqueous layer with 5 N NaOH and extract 3 times with 10 % MeOH in EtOAc. Purify by flushing the extracts through a pad of 100 g SCX ion exchange resin, rinsing with 2 liters of MeOH which was discarded, and then eluting with the 2 M NH_3 in MeOH and concentrating to give the title compound as a dark solid: ISMS 206 (M+1); Analysis for $\text{C}_{20}\text{H}_{18}\text{F}_6\text{N}_2\text{O}_2 \cdot 0.3\text{H}_2\text{O} \cdot 0.1\text{C}_7\text{H}_8$: calcd: C, 57.34; H, 5.74; N, 19.29; found: C, 57.30; H, 5.38; N, 19.08; ^1H NMR (DMSO- d_6) 11.9-11.2 (bs, 1H), 8.50-8.49 (d, 1H), 7.95-7.92 (m, 1H), 7.47-7.45 (m, 1H), 7.38 (s, 1H), 2.79 (s, 4H), 2.2-1.3 (bs, 2H).


Example 3856-Nitrotryptamine

The method of Example 384 gives (6-nitro-1H-indol-3-yl)-acetonitrile: ISMS 200 (M-1); Analysis for $\text{C}_{10}\text{H}_7\text{N}_3\text{O}_2 \cdot 0.1\text{H}_2\text{O}$: calcd: C, 59.17; H, 3.58; N, 20.70; found: C, 59.04; H, 3.28; N, 20.39 which gives the title compound: ISMS 206 (M+1); ^1H NMR (DMSO- d_6) 11.5 (bs, 2H), 8.26 (s, 1H), 7.84-7.81 (m, 1H), 7.68-7.66 (m, 1H), 7.57 (s, 1H), 2.80-74 (m, 4H) (indole N-H not observable).

Example 390N-(2-(5-Nitro-1H-indol-3-yl)ethyl)-3-phenoxybenzylamine


The method of Example 340 gives the title compound, salt formation in 10 mL EtOH with 0.25 mL 5 N HCl and 40 mL toluene then concentrating to a solid give the hydrochloride of the title compound: Analysis for $\text{C}_{23}\text{H}_{21}\text{N}_3\text{O}_3 \cdot \text{HCl} \cdot 0.2 \text{ EtOH}$: calcd: C, 64.62; H, 5.17; N, 9.75; found: C, 64.89; H, 5.40; N, 9.75; ISMS 388 (M+1).

By the method of Example 390 the following compounds were prepared, isolated as the hydrochloride except where noted:



-103-


No.	Z'	R ₄	Data
391	5-nitro	2,2,2-trifluoro ethyl	ISMS 444 (M+1); Analysis for C ₂₀ H ₂₀ ClF ₄ N ₃ O ₃ ·0.1 H ₂ O: calcd: C, 52.87; H, 4.48; N, 9.74; found: C, 52.63; H, 4.34; N, 9.67
392	5-nitro	2,2,3,3-tetrafluoro propyl	ISMS 444 (M+1); Analysis for C ₂₀ H ₂₀ ClF ₄ N ₃ O ₃ : calcd: C, 52.01; H, 4.36; N, 9.10; found: C, 51.94; H, 4.19; N, 8.93
393	6-nitro	2,2,2-trifluoro ethyl	ISMS 394 (M+1); ¹ H NMR (CDCl ₃ -freebase) 8.47 (bs, 1H), 8.31-8.30 (m, 1H), 8.01-7.98 (m, 1H), 7.63-7.61 (m, 1H), 7.32-7.31 (m, 1H), 7.24-7.21 (m, 1H), 6.94-6.92 (m, 1H), 6.88 (s, 1H), 6.80-6.77 (m, 1H), 4.33-4.26 (m, 2H), 3.79 (s, 2H), 3.00-2.93 (m, 4H), 1.54 (s, 1H)
394	6-nitro	2,2,3,3-tetrafluoro propyl	ISMS 426 (M+1); Analysis for C ₂₀ H ₁₉ F ₄ N ₃ O ₃ : calcd: C, 52.01; H, 4.36; N, 9.10; found: C, 51.96; H, 4.16; N, 8.76
395	6-nitro	2,2,3,3,3-pentafluoro propyl	ISMS 444 (M+1); Analysis for C ₂₀ H ₁₈ F ₅ N ₃ O ₃ : calcd: C, 50.06; H, 3.99; N, 8.76; found: C, 49.76; H, 3.86; N, 8.67

Example 396N-(2-(5-Amino-1H-indol-3-yl)ethyl)-3-phenoxybenzylamine

- 5 Combine N-(2-(5-nitro-1H-indol-3-yl)ethyl)-3-phenoxybenzylamine hydrochloride (250 mg, 0.64 mmol) and NiCl₂·6H₂O (460 mg, 1.9 mmol) in 30 mL MeOH and treat with NaBH₄ (73 mg, 1.9 mmol). After 1 hour concentrate to dryness, partition between EtOAc and concentrated NH₄OH. Extract the aqueous layer with EtOAc, combine the organic layer, dry over MgSO₄ and concentrate to
- 10 dryness. Chromatograph on silica gel eluting with a stepwise gradient 20/75/5 THF/hexanes/Et₃N then 40/55/5 THF/hexanes/Et₃N gives the title compound as an oil. Additional chromatograph on silica gel eluting with 1% MeOH in CHCl₃ mixed with conc. NH₄OH gives the title compound as an oil. Treatment with in 10 mL EtOH with 0.25 mL 5 N HCl and 40 mL toluene then concentrating give the
- 15 title compound as the hydrochloride: Analysis for C₂₃H₂₃N₃O 2.6 HCl 0.6 EtOH: calcd: C, 59.66; H, 5.83; N, 9.07; found: C, 59.30; H, 5.48; N, 8.82; ISMS 358 (M+1).

-104-

By the method of Example 396 the following compounds were prepared, isolated as the hydrochloride except where noted:



No.	Z'	R ₄	Data
397	5-amino	2,2,2-trifluoro ethyl	Analysis for C ₁₉ H ₂₀ F ₃ N ₃ O 2HCl 0.2 CHCl ₃ 0.3 CH ₃ OH: calcd: C, 49.85; H, 5.02; N, 8.94; found: C, 50.05; H, 4.99; N, 8.73; ISMS 364 (M+)
398	5-amino	2,2,3,3-tetrafluoro propyl	¹ H NMR (DMSO-d ₆) 11.3 (bs, 1H), 10.25 (bs, 3H), 9.6 (bs, 2H), 7.6 (s, 1H), 7.5-7.35 (m, 4H), 7.3-7.2 (m, 1H), 7.2-7.0 (m, 2H), 6.9-6.5 (d, 1H), 4.65-4.5 (t, 2H), 4.25 (s, 2H), 3.3 (s, 4H); Analysis for C ₂₀ H ₂₁ F ₄ N ₃ O 2HCl: calcd: C, 51.29; H, 4.95; N, 8.97; found: C, 51.26; H, 4.98; N, 8.26
399	6-amino	2,2,2-trifluoro ethyl	ISMS 363 (M+); C ₁₉ H ₂₂ Cl ₂ F ₃ N ₃ O·0.4 H ₂ O: calcd: C, 51.45; H, 5.18; N, 9.48; found: C, 51.45; H, 5.10; N, 9.63
400	6-amino	2,2,3,3-tetrafluoro propyl	ISMS 393 (M+); C ₂₀ H ₂₃ Cl ₂ F ₄ N ₃ O·0.2 H ₂ O: calcd: C, 50.90; H, 5.00; N, 8.90; found: C, 50.73; H, 4.82; N, 8.65

5

Example 401

6-Fluorotryptamine

Combine 6-fluoroindole (108 g, 0.8 mol) and dichloromethane (324 ml). Cool in an ice bath. Add trifluoroacetic acid (308 ml) over a few minutes (exothermic). Add a solution of Z-1-dimethylamino-2-nitroethylene (94.7 g, 0.816 mol) in dichloromethane (600 ml) during 40 minutes while maintaining the temperature at about 0-5°C. After 45 minutes, warm to about 20°C. After 2 hours, pour over 1.2 L ice water and stirring overnight with seeding to give a solid. Collect the solid by filtration, wash first with 100 ml of a mixture dichloromethane-cyclohexane 1/1, then with 750 ml of water and dry at 40°C to give 3-(2-nitrovinyl)-6-fluoroindole.


Combine LiAlH₄ (48.8 g, 1.286 moles, 5 equiv.) and THF (848 ml) and cool in an ice-water bath to about 6 °C while keeping the temperature below 32°C. Add a solution of 3-(2-nitrovinyl)-6-fluoroindole (53 g, 0.257 mol, 1 equiv.) in THF (694 ml) while keeping the temperature below about 31°C. Allow to stir at ambient temperature. After 2.5 hours, quench with a mixture of 49 ml of water and 49 ml of THF, then with 49 ml of

-105-

NaOH 15% and finally with 49 ml of water. Keep the temperature below ~32 °C during the quench. Stir for 1.5 hours, filter through a celite bed and wash with THF. Evaporate to residue, dissolve in 750 ml of diethyl ether and cool in an ice-water bath. Add a solution of HCl/diethyl ether to give a solid. Stir for 1 hour, collect the solid by filtration, wash with diethyl ether, and dry under reduced pressure at 45°C to give the hydrochloride of the title compound.

Example 402

N-(2-(6-Fluoro-1H-indol-3-yl)ethyl)-3-(2,2,3,3-tetrafluoropropoxy)benzylamine



10

Combine 6-fluorotryptamine hydrochloride (90 g, 0.419 mol) and water (900 ml). Add an aqueous solution of NaOH (2N, 230 ml) and dichloromethane (900 ml). After 1 hour, separate the organic layer, extract the aqueous layer with dichloromethane, combine the organic layers, wash water, dry over MgSO₄, and evaporate to a residue. Combine the residue and toluene (200 ml) and evaporate to give 78.45 g of a brown oil. Combine the above 78.45 g with another 41.4 g batch to provide 6-fluorotryptamine. Combine 6-fluorotryptamine (119.85) and ethanol (3.325 L), add 2,2,3,3-tetrafluoropropylbenzaldehyde (176 g, 0.745 moles, 1.2 equiv.) and 150 g of molecular sieve 3Å. Heat to reflux. After 2 hours, cool to RT room temperature and add NaBH₄ (35.2 g, 0.93 mol, 1.5 equiv.). After 1 hour, filter through celite and wash with 500 ml of ethanol. Evaporate the filtrate under reduced pressure to give an oily residue. Partition the residue between water and dichloromethane. Separate the layers, extract the aqueous later with dichloromethane, combine organic layers, wash with brine and dry over MgSO₄. Filter and evaporate under reduced pressure to give the title compound.

25

The HCl salt is formed as follows: Combine N-(2-(6-fluor-1H-indol-3-yl)ethyl)-3-(2,2,3,3-tetrafluoropropyl)benzylamine (387 g, 0.97 moles) and diethyl ether (3.95 L) of at room temperature. Add dropwise a solution of HCl/Et₂O (298 ml) over 15 minutes until the pH is about 3 to give a solid. Stir for 1 hour and collect the solid, wash with ether, and dry under reduced pressure for at 40°C to give the title compound as the hydrochloride.

30

-106-

Example 410(5-Bromo-1H-indol-3-yl)acetonitrile

The method of Example 384 using 5-bromoindole gives the title compound: ISMS 234 (M-1); Analysis for $C_{10}H_7BrN_2 \cdot 0.1H_2O$: calcd: C, 50.70; H, 3.06; N, 11.83; found: C, 50.69; H, 2.90; N, 11.64; 1H NMR ($CDCl_3$) 8.22 (s, 1H), 7.70-7.69 (m, 1H), 7.33-7.31 (m, 1H), 7.24 (s, 1H), 7.23-7.22 (m, 1H), 3.78-3.77 (m, 4H).

Example 4115-Bromotryptamine

10 Dissolve 5-bromo-1H-indole-3-carbonitrile (9.5 g, 40.4 mmol) in 200 mL dry THF and treat with 80 mL 1 M BH_3 in THF at ambient temperature. Stir overnight, the reaction and cautiously quench by dropwise addition of 5 mL water. Concentrate to dryness under vacuum and the residue. Partition between 1 N HCl and EtOAc. Extract the organic layer with 1 N HCl which was combined with the
15 original aqueous layer. The Treat the aqueous layer with 5 N NaOH and extract with EtOAc. Saturate with NaCl and extract again with EtOAc. Combine the extracts dry over $MgSO_4$ and concentrate to dryness to give 4.72 g (19.7 mmol, 49%) of an oil which crystallized.

Conversion to the oxalate salt by treating an EtOAc solution of the compound with
20 a solution of one equivalent of oxalic acid. Filter the resulting solid and dry under vacuum: Analysis for $C_{10}H_{11}BrN_2 \cdot C_2H_2O_2 \cdot H_2O$: calcd: C, 43.08; H, 4.10; N, 8.37; found: C, 43.26; H, 3.91; N, 8.20; ISMS 240 (M+1).

Example 4135-Methoxycarbonyl-1H-indole

25 Combine 5-carboxyindole (7.2 g, 44.7 mmol) in 400 mL dichloromethane and 100 mL MeOH and treat dropwise with 35 mL 2 M TMS diazomethane in hexanes. Stir overnight at ambient temperature. Concentration under vacuum to give the title compound as a solid: Analysis for $C_{10}H_9NO_2 \cdot 0.1H_2O$: calcd: C, 67.86; H, 5.24; N, 7.91; found: C, 68.03; H, 5.15; N, 7.98; 1H NMR ($CDCl_3$) 8.44 (bs, 1H), 8.412-8.409 (m, 1H), 7.91-7.88 (m, 1H), 7.46-7.38 (m, 1H), 7.26-7.24 (m, 1H), 6.64-6.63 (m, 1H), 3.92 (s, 3H); ISMS 176 (M+1).

Example 4143-Formyl-5-methoxycarbonyl-1H-indole

35 Place anhydrous DMF (25 mL) in a flask under an atmosphere of nitrogen, cool to 10°C and treat dropwise with $POCl_3$ (8.22 g, 54 mmol) while keeping the

-107-

temperature below 15°C. Add a solution of 5-methoxycarbonyl-1H-indole in 30 mL DMF portionwise keeping the temperature below 20°C. Remove the cooling bath and stir the mixture at ambient temperature for 1 hour then pour onto ice. Addition of 50 mL 5 N NaOH precipitated a solid which is filtered and rinsed with water and EtOAc to give the title compound: ¹H NMR (DMSO-d₆) 9.95 (s, 1H), 8.76 (s, 1H), 8.4 (s, 1H), 7.9-7.8 (m, 1H), 7.5-7.7 (d, 1H), 3.85 (s, 3H), 1.7 (s, 1H); ISMS 204 (M+1).

Example 415

10 3-(2-Nitroethyl)-5-methoxycarbonyl-1H-indole

The method of Example 317 to give the title compound: ¹H NMR (DMSO-d₆) 12.5 (bs, 1H), 8.38-8.37 (m, 1H), 8.37-8.34 (m, 1H), 8.23 (s, 1H), 7.87-7.84 (m, 1H), 7.80-7.77 (m, 1H), 7.57-7.55 (d, 1H), 3.85 (s, 3H); ISMS 246 (M+1).

Example 416

15 3-(2-Nitroethyl)-5-methoxycarbonyl-1H-indole

Treat a solution of 3-(2-nitrovinyl)-5-methoxycarbonyl-1H-indole (57 mg, 0.23 mmol) in 9 mL THF and 2 mL MeOH with NaBH₄ (26 mg, 0.69 mmol). Stir at ambient temperature overnight, concentrate to dryness and partition between conc. NH₄OH (10 mL) and dichloromethane. Extract the aqueous layer with dichloromethane, acidify with conc. HCl and extract twice with dichloromethane. Combine the organic layers, concentrate, and chromatograph on silica gel eluting with 1% MeOH in CHCl₃ to give the title compound as a solid: ¹H NMR (CDCl₃) 8.35 (bs, 1H), 8.32 (s, 1H), 7.92-7.90 (m, 1H), 7.38-7.36 (d, 1H), 7.12-7.11 (m, 1H), 4.69-4.65 (t, 2H), 3.93 (s, 3H), 3.51-3.48 (t, 2H); ISMS 248 (M+).

Example 417

5-Methoxycarbonyltryptamine

Combine 3-(2-nitroethyl)-5-methoxycarbonyl-1H-indole (280 mg, 1.1 mmol), PtO₂ (200 mg) and 15 mL MeOH and hydrogenate at atmospheric pressure overnight. Filter reaction mixture through a pad of celite, concentrate the filtrate, and chromatograph on silica gel eluting with 5% MeOH in CHCl₃ mixed with conc. NH₄OH to give the title compound as an oil: ISMS 219 (M+1); ¹H NMR (CDCl₃) 9.01 (s, 1H), 8.36 (s, 1H), 7.88-7.85 (m, 1H), 7.32-7.24 (m, 1H), 7.05 (s, 1H), 3.91 (s, 3H), 3.05-3.01 (m, 2H), 2.93-2.89 (m, 2H), 1.22 (bs, 2H).

-108-

Example 4182-(2-(5-Benzyloxy-1H-indol-3-yl)ethyl)isoindole-1,3-dione

Combine 5-benzyloxytryptamine hydrochloride (1 g, 3.3 mmol), phthalic anhydride (.56 g, 4.0 mmol) and N,N-diisopropylethylamine (0.86 g, 6.6 mmol) in 25 mL anhydrous pyridine and reflux for 1 hour, cool to room temperature and treat with 4 g 3Å molecular sieves. Refluxing was continued for 60 hours then the mixture was filtered. Concentrate under vacuum to give a residue which is mixed with 25 mL CHCl₃ and filtered to give a solid. Purification of the filtrate by chromatography on silica gel eluting with 1% MeOH in CHCl₃ to give an additional amount of title compound: ISMS 397 (M+1); Analysis for C₂₅H₂₀N₂O₃ 0.3H₂O C₇H₈: calcd: C, 75.09; H, 5.25; N, 6.82; found: C, 75.00; H, 5.22; N, 6.96.

By the method of Example 418 the following compounds were prepared: a) 2-(2-(5-Hydroxy-1H-indol-3-yl)ethyl)isoindole-1,3-dione: (4.5 mmol, 95%); ¹H NMR (DMSO-d₆) 10.47 (s, 1H), 8.59 (bs, 1 H), 7.84-7.78 (m, 4H), 7.09-7.06 (d, 1H), 7.03-7.02 (d, 1H), 6.85-6.84 (d, 1H), 6.56-6.54 (m, 1H), 3.79-3.75 (t, 2H), 2.91-2.87 (m, 2H).

Example 4192-(2-(5-Hydroxy-1-triisopropylsilylanyl-1H-indol-3-yl)ethyl)isoindole-1,3-dione

Combine a mixture of an oil dispersion of KH (40%, 1 g) in 30 mL anhydrous THF and a suspension of 2-(2-(5-benzyloxy-1H-indol-3-yl)ethyl)isoindole-1,3-dione (1.2 g, 3 mmol) in 30 mL THF portionwise. Stir for 1 hour at ambient temperature, cooled to 0°C, add triisopropylsilyltrifluoromethanesulfonate (1.85 g, 6 mmol) and stir an additional 1 hour at ambient temperature. Pour the reaction into a rapidly stirring solution of saturated NaHCO₃ and extract with 2 x 50 mL EtOAc. Combine the organic layer, dry over MgSO₄ and concentrate to dryness and chromatograph on silica gel eluting with 1% MeOH in CHCl₃ to give 2-(2-(5-benzyloxy-1-triisopropylsilylanyl-1H-indol-3-yl)ethyl)isoindole-1,3-dione as an oil.

Combine 2-(2-(5-benzyloxy-1-triisopropylsilylanyl-1H-indol-3-yl)ethyl)isoindole-1,3-dione and EtOAc (40 mL) and hydrogenated overnight with 1 g 5% Pd/C at atmospheric pressure. Filter through celite, concentrate to dryness, and chromatograph on silica gel eluting stepwise with a gradient 10% EtOAc in hexanes to 30% EtOAc in hexanes to give the title compound as a solid: FDMS 462 (M+1) Analysis for C₂₇H₂₄N₂O₃Si H₂O: calcd: C, 69.55; H, 7.44; N, 6.01; Found: C, 69.44; H, 7.17; N, 6.00.

-109-

Example 4202-(2-(5-Propoxy-1-triisopropylsilanyl-1H-indol-3-yl)ethyl)isoindole-1,3-dione

Combine 2-(2-(5-hydroxy-1-triisopropylsilanyl-1H-indol-3-yl)ethyl)isoindole-1,3-dione (0.7g, 1.5 mmol), cesium carbonate (1 g, 3 mmol) and 1-iodopropane (.4 g, 2.3 mmol) in DMF (25 mL) and stir at ambient temperature overnight. Pour the reaction mixture into 50% EtOAc in hexanes and wash three times with brine. Dry the organic layer over MgSO₄ and concentrate under vacuum to give an oil. Chromatograph the oil on silica gel eluting with 5% EtOAc in hexanes to give the title compound: ISMS 505 (M+1); ¹H NMR (CDCl₃) 7.80-7.78 (m, 2H), 7.67-7.65 (m, 2H), 7.30-7.27 (d, 1H), 7.12-7.11 (d, 1H), 7.02 (s, 1H), 6.77-6.74 (m, 1H), 4.01-3.96 (m, 4H), 3.12-3.08 (m, 2H), 1.86-1.81 (m 2H), 1.64-1.57 (m, 3H), 1.08-1.04 (m, 21H).

Example 4215-Propoxy-1-triisopropylsilanyltryptamine

Combine 2-(2-(5-propoxy-1-triisopropylsilanyl-1H-indol-3-yl)ethyl)isoindole-1,3-dione (416 mg, 0.8 mmol) in 20 mL EtOH and 1 mL hydrazine hydrate. Reflux for 3 hours, filter through celite and concentrate to residue. Dissolve the residue in 10 mL MeOH and load onto a 12 g SCX ion exchange cartridge and rinse sequentially with MeOH, DMF, then MeOH. Elute the product with 2 M NH₃ in MeOH to give the title compound as an oil: ISMS 375 (M+1); ¹H NMR (CDCl₃) 7.34-7.32 (d, 1H), 7.02 (s, 1H), 7.00-6.99 (d, 1H), 6.80-6.77 (m, 1H), 3.97-3.94 (m, 2H), 3.01-2.98 (m, 2H), 2.86-2.83 (m, 2H), 1.88-1.76 (m 2H), 1.70-1.58 (m, 3H), 1.3 (bs, 2H), 1.14-1.08 (m, 18H), 1.06-1.02 (t, 3H).

Example 4226-Benzyloxytryptamine

Add to a mixture of LAH (6.2 g, 163.1 mmol) and 300 mL dry THF a solution of 3-(2-nitrovinyl)-6-benzyloxy-1H-indole (9 g, 30.6 mmol) in 200 mL THF. Reflux the mixture overnight and then cool to 0°C and quench sequentially with 6.2 mL water, 6.2 mL 15% aqueous NaOH and 18.6 mL water. After stirring 2 hours, filter through celite and concentrate to give 7.9 g (96%) of the title compound as an oil: ¹H NMR (CDCl₃) 8.06 (bs, 1H), 7.47-7.43 (m, 3H), 7.38-7.35 (m, 2H), 7.32-7.28 (m, 1H), 6.88-6.84 (m, 3H), 5.08 (s, 2H), 3.01-2.97 (m, 2H), 2.87-2.83 (m, 2H), 1.6 (bs, 2H).

-110-

Example 423N-t-Butoxycarbonyl-2-(6-benzyloxy-1H-indol-3-yl)ethylamine

The method of Example 20 gives the title compound: ¹H NMR (CDCl₃)
7.84 (bs, 1H), 9.36 (s, 2H), 8.91 (s, 1H), 7.38-7.33 (m, 2H), 7.28-7.26 (m, 1H),
5 7.20-7.18 (m, 1H), 7.09-7.07 (m, 1H), 6.94-6.93 (m, 1H), 6.68-6.67 (m, 1H), 6.50-
6.47 (m, 1H), 4.79-4.72 (m, 2H), 4.13 (s, 2H), 3.05-3.02 (m, 4H).

Example 425N-t-Butoxycarbonyl-2-(6-hydroxy-1H-indol-3-yl)ethylamine

10 The method of Example 471 gives the title compound.

Example 4282-(2-(5-Ethoxy-1H-indol-3-yl)ethyl)isoindole-1,3-dione

Combine 2-(2-(5-hydroxy-1H-indol-3-yl)ethyl)isoindole-1,3-dione (900
15 mg, 2.9 mmol), cesium carbonate (960 mg, 2.9 mmol) and 1-iodoethane (920 mg,
5.9 mmol) in N-methylpyrrolidinone (5 mL) and stir at ambient temperature for 4
hours, pour into brine and extract twice with EtOAc. Wash the combined extracts
three times with brine, dry over MgSO₄, and concentrate under vacuum to give an
oil. Chromatograph the oil on silica gel eluting with 20% EtOAc in hexanes to
20 give the title compound as a white solid: ISMS 335 (M+1); Analysis for
C₂₀H₁₈N₂O₃: calcd: C, 71.84; H, 5.43; N, 8.38; found: C, 71.97; H, 5.47; N, 8.36.

By the method of Example 428 the following compounds were prepared:

a) 2-(2-(5-Isopropoxy-1H-indol-3-yl)ethyl)isoindole-1,3-dione: ISMS 348
(M+); ¹H NMR (CDCl₃) 7.94 (bs, 1H), 7.82-7.80 (m, 2 H), 7.70-7.67 (m, 2H), 7.21-
25 7.19 (d, 1H), 7.18 (s, 1H), 7.05-7.04 (d, 1H), 6.82-6.79 (m, 1H), 4.55-4.49 (m, 1H),
3.99-3.95 (m, 2H), 3.11-3.07 (m, 2H), 1.64-1.33 (d, 6H);

b) 2-(2-(5-(2,2,2-Trifluoroethoxy)-1H-indol-3-yl)ethyl)isoindole-1,3-dione:
ISMS 389 (M+1); Analysis for C₂₀H₁₅F₃N₂O₃: calcd: C, 61.86; H, 3.89; N, 7.21;
found: C, 61.77; H, 3.83; N, 7.20;

30 c) 2-(2-(5-Butoxy-1H-indol-3-yl)ethyl)isoindole-1,3-dione: ISMS 363
(M+1); Analysis for C₂₂H₂₂N₂O₃: calcd: C, 72.91; H, 6.11; N, 7.73; found: C,
72.76; H, 6.09; N, 7.42; ¹H NMR (CDCl₃) 7.86-7.81 (m, 3H), 7.72-7.68 (m, 2 H),
7.23-7.20 (m, 1H), 7.16-7.15 (m, 1H), 7.08-7.07 (m, 1H), 6.85-6.84 (m, 1H),
6.40-3.98 (m, 4H), 3.13-3.09 (m, 2H), 1.83-1.76 (m, 2H), 1.56-1.48 (m, 2H),
35 1.01-0.98 (t, 3H);

-111-

d) 2-(2-(5-Nitro-1H-indol-3-yl)ethyl)isoindole-1,3-dione: ISMS 334 (M-1); Analysis for $C_{18}H_{13}N_3O_4 \cdot 0.1H_2O$: calcd: C, 64.13; H, 3.95; N, 12.47; found: C, 64.05; H, 3.82; N, 12.27.

By the method of Example 421 the following compounds were prepared:

- 5 a) 5-Ethoxytryptamine: ISMS 205 (M+1); Analysis for $C_{12}H_{16}N_2O \cdot H_2O$: calcd: C, 69.33; H, 7.95; N, 13.48; found: C, 69.62; H, 7.75; N, 13.30;
- b) 5-Isopropoxytryptamine: ISMS 219 (M+1); 1H NMR ($CDCl_3$) 8.57 (bs, 1H), 7.20-7.18 (d, 1 H), 7.08-7.07 (d, 1H), 6.95 (s, 1H), 6.84-6.82 (m, 1H), 4.54-4.48 (m, 1H), 3.01-2.98 (m, 2H), 2.86-2.83 (m, 2H), 1.38 (bs, 2H), 1.35-1.33 (d, 6H);
- 10 c) 5-(2,2,2-Trifluoroethoxy)tryptamine: ISMS 258 (M+); 1H NMR ($CDCl_3$) 8.33 (bs, 1H), 7.26-7.24 (d, 1 H), 7.09-7.08 (d, 1H), 7.03-7.02 (m, 1H), 6.90-6.87 (m, 1H), 4.40-4.34 (m, 2H), 3.03-3.00 (m, 2H), 2.87-2.84 (m, 2H), 1.44 (bs, 2H);
- d) 5-Butyloxytryptamine: 1H NMR ($CDCl_3$) 8.08 (bs, 1H), 7.23-7.21 (d, 1 H), 7.03-7.02 (d, 1H), 7.03-7.02 (m, 1H), 6.98-6.83 (m, 1H), 4.01-3.98 (m, 2H),
- 15 3.02-2.99 (m, 2H), 2.87-2.84 (m, 2H), 1.82-1.74 (m, 2H), 1.56-1.50 (m, 2H), 1.32 (bs, 2H), 1.00-0.96 (t, 3H);

Example 429

5-Benzenesulfonyl-1H-indole

- 20 Place a 35% oil dispersion of KH (6 g) in a flask under nitrogen, rinse with 50 mL hexanes and dry under vacuum. Cool the solid suspension in 100 mL anhydrous DMF to 0°C. Add dropwise over 10 minutes a solution of 5-bromoindole (10.3 g, 52.5 mmol) in 25 mL DMF. Stir the mixture 1 hour at 0°C then treat with triisopropylsilyltrifluoromethane sulfonate (32.2 g, 105.1 mmol).
- 25 Remove the cooling bath and stir the reaction 72 hours before pouring into 500 mL water and extracting with EtOAc. Dilute the combined extracts with hexanes, wash with brine then dry over $MgSO_4$. Concentration under vacuum and chromatograph on silica gel eluting with 1% EtOAc in hexanes to give 5-Bromo-1-triisopropylsilyl-1H-indole as a colorless oil: 1H NMR ($CDCl_3$) 7.73-7.72 (d, 1H), 7.36-7.34 (d, 1 H), 7.24-7.23 (d, 1H), 7.21-7.19 (m, 1H), 6.55-6.54 (m, 1H),
- 30 1.72-1.61 (m, 3H), 1.13-1.10 (m, 18H).

- Cool a solution of 5-bromo-1-triisopropylsilyl-1H-indole (9 g, 25.5 mmol) in 550 mL anhydrous THF to
- 75°C under argon and treat with 1.7 M t-butyl lithium (33mL, 56.2 mmol) while
- 35 keeping the temperature below
- 60°C. After the addition, recool the reaction mixture to about –73°C before adding a solution of phenylsulfonyl fluoride (4.6 g, 28.7 mmol) in 30 mL THF. Stir the

-112-

reaction at -78°C for 1 hour then quench with saturated NaHCO_3 followed by brine. Separate the layers and extract the aqueous layer with EtOAc . Treat the combined organic layers with 1 M tetrabutylammonium fluoride (35 mL) in THF for 1 hour at ambient temperature, then concentrate to dryness. Combine the residue with EtOAc , wash twice with 1 N HCl , dry over MgSO_4 , and concentrate to an oil. Chromatograph the oil on silica get eluting stepwise with 50% CHCl_3 in hexanes followed by 50% CHCl_3 in MeOH give an oily solid. Triturate the oily solid with CHCl_3 to give the title compound as a solid: Analysis for $\text{C}_{14}\text{H}_{11}\text{NO}_2\text{S}$ H_2O : calcd: C, 64.89; H, 4.36; N, 5.41; found: C, 64.76; H, 4.45; N, 5.33; ISMS 257 (M^+).

Example 430

2-(2-(5-Amino-1H-indol-3-yl)ethyl)isoindole-1,3-dione

Combine a mixture of 2-(2-(5-nitro-1H-indol-3-yl)ethyl)isoindole-1,3-dione (1.8 g, 5.4 mmol), PtO_2 (500 mg), 100 mL MeOH and 100 mL THF and hydrogenate at atmospheric pressure overnight. Filter the reaction through a pad of celite and concentrate to dryness. Redissolve the residue in 50/50 chloroform/dichloromethane and refilter through a pad of celite. Concentration under vacuum give the title compound as a dark solid: ISMS 306 (M^+); Analysis for $\text{C}_{18}\text{H}_{13}\text{N}_3\text{O}_4$ $0.1\text{C}_7\text{H}_8$ 0.2 dichloromethane: calcd: C, 68.70; H, 4.89; N, 12.58; found: C, 69.08; H, 4.75; N, 12.69; ^1H NMR (CDCl_3) 7.9-7.8 (m, 3H), 7.75-7.65 (m, 2 H), 7.2-7.1 (m, 1H), 7.05-7.0 (m, 2H), 6.7-6.6 (m, 1H), 4.0-3.9 (m, 2H), 3.4 (bs, 2H), 3.1-3.0 (m, 2H).

Example 431

2-(2-(5-Benzoylamino-1H-indol-3-yl)ethyl)isoindole-1,3-dione

Combine 2-(2-(5-amino-1H-indol-3-yl)ethyl)isoindole-1,3-dione (0.5 g, 1.64 mmol) and 4-dimethylaminopyridine (0.3 g, 2.5 mmol) and dissolve in 30 mL dichloromethane and cool to 0°C . Treat the reaction mixture with benzoyl chloride (276 mg, 1.96 mmol) and stir overnight during which the temperature was allowed to warm to room temperature. Concentrate to residue and chromatograph the residue on silica gel eluting with 0.5% MeOH in CHCl_3 to give the title compound as a solid: ISMS 410 (M^+); ^1H NMR (CDCl_3) 7.86-7.85 (m, 2H), 7.79 (s, 1 H), 7.72-7.68 (m, 2H), 7.60-7.57 (m, 2H), 7.46-7.42 (m, 1H), 7.4-7.36 (m, 3H), 7.1 3-7.11 (d, 1H), 6.89-6.88 (m, 1H), 3.88-3.84 (t, 2H), 3.00-2.97 (t, 2H).

By the method of Example 431 the following compounds were prepared:

a) 2-(2-(5-Methanesulfonylamino-1H-indol-3-yl)ethyl)isoindole-1,3-dione: ISMS 384 (M^+); ^1H NMR (CDCl_3) 10.84 (s, 1H), 9.21 (s, 1H), 7.83-7.76 (m, 4H),

-113-

7.39-7.38 (m, 1 H), 7.27-7.24 (m, 1H), 7.17-7.16 (m, 1H), 6.96-6.93 (m, 1H), 3.83-3.80 (m, 2H), 2.98-2.94 (m, 2H), 2.79 (s, 3H), 3.88-3.84 (t, 2H), 3.00-2.97 (t, 2H).

By the method of Example 421 the following compounds were prepared:

- 5 a) 5-Benzoylamino-tryptamine: ^1H NMR (CD_3OD) 7.94-7.92 (m, 2H), 7.85 (s, 1H), 7.54-7.47 (m, 3H), 7.34-7.29 (m, 2 H), 7.08 (s, 1H), 4.86 (s, 2H), 3.33 (s, 2H), 2.95-2.86 (m, 4H); and
- b) 5-Methanesulfonylamino-tryptamine: ISMS 253 (M^+); ^1H NMR (CD_3OD) 7.46-7.45 (d, 1H), 7.31-7.28 (d, 1H), 7.08 (s, 1 H), 7.04-7.01 (m, 1H), 4.86 (s, 4H),
10 2.89-2.83 (m, 7H).

Example 432

5-Ethoxycarbonyl-1H-indole

- Combine 5-carboxyindole (4.8 g, 29.8 mmol) in 150 mL THF and
15 carbonyldiimidazole (9.7 g, 59.6 mmol) and stir overnight at ambient temperature. Treat the reaction mixture with 25 mL EtOH and 1.2 g (29.8 mmol) of a 60% oil dispersion of NaH and stir for 2 hours. Concentration under vacuum gives a residue. Partition the residue between 150 mL EtOAc and 100 mL brine. Separate the layers, dry the organic layer over MgSO_4 , filter, and concentrated to an oil.
- 20 Chromatograph on silica gel eluting with 1% MeOH in CHCl_3 to give 7.2 g of an oil. Crystallize the oil from toluene gives the title compound: Analysis for $\text{C}_{11}\text{H}_{11}\text{NO}_2$: calcd: C, 69.83; H, 5.86; N, 7.40; found: C, 69.82; H, 5.90; N, 7.38; ISMS 190 ($\text{M}+1$).

Example 433

5-(N-Butylamido)-1H-indole

- Dissolve mixture of 5-carboxyindole (5 g, 31 mmol) in 150 mL THF and treat with carbonyldiimidazole (5 g, 31 mmol) and stir overnight at ambient temperature. Treat the reaction mixture with n-butylamine 4.5 g (62 mmol) and
30 reflux for 1 hour. Concentration under vacuum gives a residue which is dissolved in EtOAc. Wash sequentially with 5 N HCl, 5 N NaOH, and then brine. Dry the organic layer over MgSO_4 and concentrated to give the title compound as an oil: ^1H NMR (CDCl_3) 8.54 (bs, 1H), 8.07-8.06 (m, 1H), 7.63-7.61 (m, 1H), 7.39-7.37 (m, 1H), 7.26-7.24 (m, 1H), 6.60-6.59 (m, 1H), 6.14 (bs, 1H), 3.5-3.45 (m, 2H), 1.64-
35 1.57 (m, 2H), 1.47-1.37 (m, 2H), .97-0.93 (m, 3H); EIMS 217 ($\text{M}+1$).

Example 4345-(N-Propylamido)-1H-indole

The method of Example 433 gives the title compound: ^1H NMR (CDCl_3) 8.07 (bs, 1H), 8.07 (s, 1H), 7.63-7.60 (m, 1H), 7.38-7.36 (m, 1H), 7.25-7.24 (m, 1H), 6.59-6.58 (m, 1H), 6.21 (bs, 1H), 3.46-3.41 (m, 2H), 1.69-1.60 (m, 2H), 1.00-0.96 (m, 3H); EIMS 203 ($\text{M}+1$).

By the method of Example 414 the following compounds were prepared:

- a) 3-Formyl-5-benzenesulfonyl-1H-indole: ISMS 286 ($\text{M}+1$); ^1H NMR (DMSO-d_6) 9.83 (s, 1H), 8.55 (s, 1H), 7.89-7.86 (m, 2H), 7.61 (s, 2H), 7.59-7.52 (m, 3H), 1.70 (s, 3H).
- b) 3-Formyl-5-ethoxycarbonyl-1H-indole: Analysis for $\text{C}_{12}\text{H}_{11}\text{NO}_3$: calcd: C, 66.35; H, 5.10; N, 6.45; found: C, 65.97; H, 5.17; N, 6.46; ISMS 218 ($\text{M}+1$);
- c) 3-Formyl-N-butylamido-1H-indole: Analysis for $\text{C}_{14}\text{H}_{16}\text{N}_2\text{O}_2 \cdot 0.1\text{H}_2\text{O}$: calcd: C, 68.33; H, 6.64; N, 11.38; found: C, 68.35; H, 6.24; N, 11.30; ISMS 245 ($\text{M}+1$);
- d) 3-Formyl-5-(N-propylamido)-1H-indole: Analysis for $\text{C}_{13}\text{H}_{14}\text{N}_2\text{O}_2$: calcd: C, 67.81; H, 6.13; N, 12.16; found: C, 67.42; H, 6.04; N, 12.10; ^1H NMR (DMSO-d_6) 9.95 (s, 1H), 8.6 (s, 1H), 8.48-8.45 (t, 1H), 8.36-8.35 (m, 1H), 7.76-7.73 (m, 1H), 7.52-7.50 (d, 1H), 3.32 (bs, 1H), 3.24-3.19 (m, 2H), 1.58-1.48 (m, 2H), 0.90-0.86 (m, 3H); EIMS 230 ($\text{M}+$);
- e) 3-Formyl-6-benzyloxy-1H-indole: ^1H NMR (DMSO-d_6) 11.93 (s, 1H), 9.83 (s, 1H), 8.12-8.11 (m, 1H), 7.92-7.90 (m, 1H), 7.45-7.27 (m, 5H), 7.04-7.03 (m, 1H), 6.92-6.89 (m, 1H), 5.11 (s, 2H).

By the method of Example 415 the following compounds were prepared: a)

- 5-Benzenesulfonyl-3-(2-nitrovinyl)-1H-indole: Analysis for $\text{C}_{16}\text{H}_{12}\text{N}_2\text{O}_4\text{S} \cdot 0.1\text{H}_2\text{O}$: calcd: C, 58.42; H, 3.83; N, 8.31; found: C, 58.63; H, 3.52; N, 8.02; ISMS 229 ($\text{M}+1$);
- b) 3-(2-Nitrovinyl)-5-ethoxycarbonyl-1H-indole: Analysis for $\text{C}_{16}\text{H}_{12}\text{N}_2\text{O}_4\text{S} \cdot 0.1\text{H}_2\text{O}$: calcd: C, 58.42; H, 3.83; N, 8.31; found: C, 58.63; H, 3.52; N, 8.02; ISMS 229 ($\text{M}+1$);
- c) 3-(2-Nitro-vinyl)-N-butylamido-1H-indole: Analysis for $\text{C}_{15}\text{H}_{17}\text{N}_3\text{O}_3$: calcd: C, 62.71; H, 5.96; N, 14.62; found: C, 62.46; H, 5.81; N, 14.38; ISMS 288 ($\text{M}+1$);

-115-

d) 3-(2-Nitro-vinyl)-N-propylamido 1H-indole: ISMS 273 M(+1); ¹H NMR (DMSO-d₆) 12.38 (s, 1H), 8.62-8.59 (t, 1H), 8.43-8.39 (d, 1H), 8.37 (s, 1H), 8.31-8.30 (d, 1H), 8.18-8.15 (d, 1H), 7.84-7.82 (m, 1H), 7.55-7.53 (d, 1H), 3.31-3.24 (m, 2H), 1.61-1.52 (m, 2H), 0.92-0.89 (t, 3H); Analysis for C₁₄H₁₅N₃O₃ 0.1H₂O: calcd: C, 61.12; H, 5.57; N, 15.28; found: C, 61.06; H, 5.38; N, 15.05;

e) 3-(2-Nitro-vinyl)-6-benzyloxy-1H-indole: ¹H NMR (DMSO-d₆) 11.85 (bs, 1H), 8.32-8.29 (m, 1H), 8.09 (s, 1H), 7.94-7.91 (m, 1H), 7.83-7.81 (m, 1H), 7.45-7.43 (m, 2H), 7.38-7.31 (m, 2H), 7.29-7.27 (m, 1H), 7.05-7.04 (m, 1H), 6.92-6.89 (m, 1H), 5.13 (s, 2H).

10

By the method of Example 416 the following compounds were prepared:

a) 5-Benzenesulfonyl-3-(2-nitroethyl)-1H-indole: Analysis for C₁₆H₁₄N₂O₄S 0.1H₂O: calcd: C, 57.85; H, 4.31; N, 8.43; found: C, 57.72; H, 4.22; N, 8.25; ISMS 329 (M-1);

b) 3-(2-Nitroethyl)-5-ethoxycarbonyl-1H-indole: Analysis for C₁₃H₁₄N₂O₄: calcd: C, 59.54; H, 5.38; N, 10.68; found: C, 59.23; H, 5.25; N, 10.53; ISMS 263 (M+1);

c) 3-(2-Nitroethyl)-N-butylamido-1H-indole: Analysis for C₁₅H₁₉N₃O₃: calcd: C, 62.27; H, 6.62; N, 14.52; found: C, 61.98; H, 6.39; N, 14.42; ISMS 290 (M+1); and

d) 3-(2-Nitroethyl)-N-propylamido-1H-indole: ¹H NMR (CDCl₃) 8.52 (bs, 1H), 8.06 (s, 1H), 7.58-7.55 (m, 1H), 7.35-7.33 (m, 1H), 7.10-7.09 (m, 1H), 6.23 (bs, 1H), 4.65-4.61 (t, 2H), 3.48-3.43 (m, 4H), 1.71-1.62 (m, 2H), 1.01-0.98 (t, 3H); Analysis for C₁₄H₁₇N₃O₃ 0.1H₂O: calcd: C, 60.68; H, 6.26; N, 15.16; found: C, 60.88; H, 6.05; N, 15.07.

25

By the method of Example 421 the following compounds were prepared:

a) 5-Benzenesulfonyltryptamine: ISMS 301 (M+1); ¹H NMR (HCl-DMSO-d₆) (s, 1H), 8.3 (s, 1H), 8.2 (bs, 2H), 8.0-8.9 (m, 2H), 7.4-7.2 (m, 5H), 7.1-7.0 (m, 1H), 3.2-3.0 (s, 4H);

30

b) 5-Ethoxycarbonyltryptamine (isolated as the oxalate salt): Analysis for C₁₃H₁₆N₂O₂ C₂H₂O₄: calcd: C, 55.90; H, 5.63; N, 8.69; found: C, 56.07; H, 5.54; N, 8.29; ISMS 233 (M+1); and

c) 5-N-Butylamidotryptamine: Analysis for C₁₅H₂₁N₃O 0.3 H₂O: calcd: C, 68.05; H, 8.22; N, 15.87; found: C, 68.36; H, 8.11; N, 15.49; ISMS 260 (M+1); and

35

-116-

- d) 5-N-Propylamidotryptamine:(isolated as the oxalate salt): Analysis for $C_{14}H_{19}N_3O$ $C_2H_2O_4$ 0.1EtOAc: calcd: C, 57.23; H, 6.38; N, 12.21; found: C, 57.48; H, 6.53; N, 12.12; 1H NMR (DMSO- d_6) 11.2 (s, 1H), 8.4 (t, 1H), 8.2 (s, 1H), 7.75-7.65 (m, 1H), 7.6 (bs, 4H), 7.4-7.35 (m, 1H), 7.3-7.25 (d, 1H), 3.3-3.2 (m, 2H), 3.15-3.0 (m, 4H), 1.6-1.45 (m, 2H), 0.9-0.8 (t, 3H); ISMS 246 (M+1).

Example 435

N-t-Butoxycarbonyl-2-(6-butoxy-1H-indol-3-yl)ethylamine

- Combine N-t-butoxycarbonyl-2-(6-hydroxy-1H-indol-3-yl)ethylamine (250 mg, 0.9 mmol), cesium carbonate (295 mg, 0.9 mmol) and 1-iodobutane (200 mg, 1.1 mmol) and N-methylpyrrolidinone (10 mL) and stir at ambient temperature for 2 hours and pour into 75 mL brine. Extract the mixture twice with 25 mL EtOAc. Wash the combined extracts with brine 2X50 mL, dry over $MgSO_4$, and concentrate under vacuum to give an oil. Chromatograph the oil on silica gel eluting with 30% EtOAc in hexanes to give the title compound as a solid: ISMS 333 (M+1) ; Analysis for $C_{19}H_{28}N_2O_3$: calcd: C, 68.65; H, 8.49; N, 8.43; found: C, 68.83; H, 8.18; N, 8.33.

- By the method of Example 435 the following compounds were prepared: a) N-t-Butoxycarbonyl-2-(6-ethoxy-1H-indol-3-yl)ethylamine: ISMS 305 (M+1); Analysis for $C_{17}H_{24}N_2O_3$: calcd: C, 67.08; H, 7.95; N, 9.20; found: C, 66.85; H, 7.79; N, 9.14.

Example 436

6-Butoxytryptamine

- Combine N-t-butoxycarbonyl-2-(6-butoxy-1H-indol-3-yl)ethylamine (430 mg, 1.3 mmol), 1 mL anisole and 5 mL trifluoroacetic acid and stir at room temperature for 2 hours. Concentrate the reaction to dryness under vacuum, mix with 10 mL concentrated NH_4OH and extract with 20 mL dichloromethane. Dry the extract over $MgSO_4$ and concentrated to 300 mg oil (1.3 mmol, 100%).

- By the method of Example 436 the following compounds were prepared: a) 6-Ethoxytryptamine: ISMS 305 (M+1); Analysis for $C_{17}H_{24}N_2O_3$: calcd: C, 67.08; H, 7.95; N, 9.20; found: C, 66.85; H, 7.79; N, 9.14

Example 437

N-t-Butoxycarbonyl-2-(6-phenylsulfonate-1H-indol-3-yl)ethylamine


- Combine N-t-butoxycarbonyl-2-(6-hydroxy-1H-indol-3-yl)ethylamine (750 mg, 2.7 mmol) and pyridine (430 mg, 5.4 mmol) in dichloromethane (30 mL) and cool to 0°C and treat with benzene sulfonyl chloride (480 mg, 2.7 mmol). Allow the reaction to warm to room temperature and stir overnight. Concentrate to dryness

-117-

the mixture under vacuum, mix with dichloromethane and chromatograph on silica gel eluting with 30% EtOAc in hexanes to give N-t-butoxycarbonyl-2-(6-phenylsulfonate-1H-indol-3-yl)ethylamine as an oil: ISMS 415 (M-1); ¹H NMR (CDCl₃) 8.14 (bs, 1H), 7.66-7.62 (m, 2H), 7.51-7.47 (m, 1H), 7.40-7.38 (m, 2H), 7.10 (s, 1H), 7.04-7.03 (m, 2H), 6.59-6.57 (m, 1H), 4.57 (bs, 1H), 3.40-3.80 (m, 2H), 2.89-2.86 (m, 2H), 1.41 (s, 9H).

Place N-t-butoxycarbonyl-2-(6-phenylsulfonate-1H-indol-3-yl)ethylamine (0.5 g, 1.2 mmol) in a flask with a stream of N₂ passing through it and heat to 200 °C overnight and cool to room temperature. Dissolve the residue in dichloromethane and chromatography on silica gel eluting with 2% MeOH in CHCl₃-NH₄OH to give the title compound as an oil.

By the method of Example 425 the following compounds were prepared and isolated as the hydrochloride except where noted:




No.	Z'	R ₄	Data
438	5-propoxy	phenyl	ISMS 401 (M+1); Analysis for C ₂₀ H ₁₈ N ₂ O ₃ 0.1H ₂ O: calcd: C, 71.17; H, 6.71; N, 6.38; found: C, 71.02; H, 6.54; N, 6.33; ¹ H NMR (Free base-CDCl ₃) 7.93 (bs, 1H), 7.34-7.30 (m, 2H), 7.35-7.28 (m, 2H), 7.12-7.07 (m, 1H), 7.06-6.96 (m, 6H), 6.89-6.84 (m, 2H), 3.97-3.94 (m, 2H), 3.79 (s, 2H), 2.97-2.94 (m, 4H), 1.89-1.7 (m, 2H), 1.51 (bs, 1H), 1.07-1.04 (t, 3H)

15

Example 440

N-(2-(5-Propoxy-1H-indol-3-yl)ethyl)-3-(2,2,3,3-tetrafluoropropoxy)benzylamine




Combine 2-(5-propoxy-1-triisopropylsilyl)tryptamine (138 mg, 0.37 mmol), 3-(2,2,3,3-tetrafluoropropoxy)benzaldehyde (87 mg, 1.8 mmol) and 1 g 3 Å molecular sieves in 25 mL EtOH and reflux overnight. Decant the liquid into a separate flask, cool to 0°C and treat with 42 mg (1.1 mmol) NaBH₄. Stir the reaction at ambient temperature for 1 hour, treat with 0.74 mmol of tetrabutylammonium fluoride and stir for an additional

20

-118-

hour. Concentrate under vacuum to give a residue. Chromatograph the residue on silica gel eluting with 10% MeOH in CHCl₃ to give the title compound. Treat the title compound with 10 mL EtOH with 0.25 mL 5 N HCl and 40 mL toluene then concentrating gives the hydrochloride of the title compound: ISMS 439 (M+1); ¹H NMR (Free base-CDCl₃) 7.89 (bs, 1H), 7.23-7.21 (m, 1H), 7.03-7.02 (d, 1H), 6.99-6.98 (d, 1H), 6.94-6.92 (m, 1H), 6.89-6.83 (m, 2H), 6.78-6.75 (m, 1H), 6.18-5.90 (m, 1H), 4.29-4.23 (m, 2H), 3.95-3.91 (m, 2H), 3.78 (s, 2H), 2.95 (s, 4H), 1.85-1.75 (m, 2H), 1.51 (bs, 1H), 1.06-1.03 (t, 3H).

By the method of Example 440 the following compounds were prepared and isolated as the hydrochloride except where noted:




No.	Z'	R ₄	Data
441	5-n-propyl amido	2,2,2-trifluoro ethyl	ISMS 434 (M+1); C ₂₃ H ₂₆ F ₃ N ₃ O ₂ •HCl•0.8H ₂ O•0.1C ₇ H ₈ : calcd: C, 57.67; H, 6.00; N, 8.51; found: C, 57.55; H, 5.77; N, 8.43
442	5-ethoxy carbonyl	2,2,3,3-tetrafluoro propyl	ISMS 453 (M+1); C ₂₃ H ₂₇ ClF ₃ N ₃ O ₂ : calcd: C, 56.50; H, 5.15; N, 5.73; found: C, 56.26; H, 5.04; N, 5.76
443	5-ethoxy carbonyl	phenyl	ISMS 415 (M+1); C ₂₆ H ₂₇ ClN ₂ O ₃ •0.1H ₂ O: calcd: C, 68.97; H, 6.06; N, 6.19; found: C, 68.78; H, 5.87; N, 6.19
445	5-phenoxy	2,2,3,3-tetrafluoro propyl	ISMS 473 (M+1); C ₂₆ H ₂₅ ClF ₄ N ₂ O ₂ •0.5H ₂ O: calcd: C, 60.29; H, 5.06; N, 5.41; found: C, 60.27; H, 4.81; N, 5.33 (isolated as the base)
456	H	2,2,2-trifluoro ethyl	ISMS 349 (M+1); C ₁₉ H ₂₀ ClF ₃ N ₂ O•0.2H ₂ O: calcd: C, 58.75; H, 5.29; N, 7.21; found: C, 58.62; H, 5.04; N, 7.08
457	H	2,2,3,3,3-pentafluoro propyl	ISMS 385 (M+1); C ₁₉ H ₁₈ ClF ₅ N ₂ O•0.2H ₂ O: calcd: C, 53.77; H, 4.37; N, 6.60; found: C, 53.81; H, 4.19; N, 6.59
458	5-phenyl	phenyl	Analysis for C ₂₉ H ₂₆ N ₂ O•HCl•0.2H ₂ O: calcd: C, 75.95; H, 6.02; N, 6.11; found: C, 76.01; H, 5.92; N, 5.97 ISMS 419 (M+1)

No.	Z'	R ₄	Data
459	5-(4-fluorophenyl)	phenyl	Analysis for C ₂₉ H ₂₅ FN ₂ O•HCl•0.2H ₂ O: calcd: C, 73.08; H, 5.58; N, 5.88; found: C, 72.99; H, 5.38; N, 5.83 ISMS 437 (M+1)
460	5-(N-butylamido)	2,2,2-trifluoro ethyl	Analysis for C ₂₄ H ₂₈ F ₃ N ₃ O ₂ •HCl•0.7H ₂ O: calcd: C, 58.05; H, 6.17; N, 8.46; found: C, 57.86; H, 5.98; N, 8.39 ISMS 448 (M+1)
461	5-hydroxy	2,2,2-trifluoro ethyl	ISMS 365 (M+1) ¹ H NMR (DMSO-d ₆) 10.6 (bs, 1H), 9.4 (bs, 2H), 8.75 (s, 1H), 7.45-6.6 (m, 7H), 4.9-4.7 (m, 2H), 4.2 (bs, 2H), 3.2-2.9 (m, 4H);
462	5-benzyloxy	2,2,3,3,3-pentafluoro propyl	Analysis for C ₂₇ H ₂₅ F ₅ N ₂ O ₂ •HCl: calcd: C, 58.87; H, 4.95; N, 5.09; found: C, 59.02; H, 4.76; N, 5.1 ISMS 505 (M+1) ⁴
463	6-benzyloxy	2,2,2-trifluoro ethyl	ISMS 455 (M+1) ¹ H NMR (CDCl ₃ -freebase) 7.88 (bs, 1H), 7.48-7.45 (m, 3H), 7.41-7.34 (m, 2H), 7.35-7.30 (m, 1H), 7.25-7.23 (m, 1H), 6.95-6.93 (m, 1H), 6.91-6.90 (m, 2H), 6.88-6.86 (m, 2H), 6.81-6.79 (m, 1H), 5.10 (s, 2H), 4.31-4.25 (m, 2H), 3.79 (s, 2H), 2.96 (s, 4H), 1.65 (bs, 1H)
464	6-benzyloxy	2,2,3,3-tetrafluoro propyl	ISMS 487 (M+1) Analysis for C ₂₇ H ₂₇ F ₄ N ₂ O ₂ •HCl: calcd: C, 62.01; H, 5.20; N, 5.36; found: C, 61.69; H, 5.07; N, 5.33
465	6-butyloxy	2,2,2-trifluoro ethyl	¹ H NMR (CDCl ₃ -freebase) 7.86 (bs, 1H), 7.45-7.43 (m, 1H), 7.23-7.19 (m, 1H), 6.94-6.92 (m, 1H), 6.89-6.88 (m, 1H), 6.84-6.75 (m, 4H), 4.29-4.23 (m, 2H), 3.99-3.96 (m, 2H), 3.78 (s, 2H), 2.94 (s, 4H), 1.81-1.74 (m, 2H), 1.55-1.45 (m, 3H), 0.99-0.95 (m, 3H); Analysis for C ₂₃ H ₂₇ F ₃ N ₂ O ₂ HCl: calcd: C, 60.46; H, 6.18; N, 6.13; found: C, 60.23; H, 5.99; N, 6.01
466	5-butyloxy	2,2,3,3-tetrafluoro propyl	Analysis for C ₂₄ H ₂₈ F ₄ N ₂ O ₂ •HCl: calcd: C, 58.96; H, 5.98; N, 5.73; found: C, 58.62; H, 5.96; N, 5.77 ISMS 453 (M+1)

-120-


No.	Z'	R ₄	Data
467	6-ethoxy	2,2,2-trifluoroethyl	ISMS 393 (M+1); Analysis for C ₂₁ H ₂₃ F ₃ N ₂ O ₂ HCl: calcd: C, 58.81; H, 5.64; N, 6.53; found: C, 58.94; H, 5.58; N, 6.55
468	6-phenyl sulfonate	2,2,2-trifluoroethyl	ISMS 505 (M+1); Analysis for C ₂₅ H ₂₃ F ₃ N ₂ O ₄ S HCl: calcd: C, 55.51; H, 4.47; N, 5.18; found: C, 55.27; H, 4.41; N, 5.15
469	6-phenyl sulfonate	2,2,3,3-tetrafluoropropyl	ISMS 536 (M+1); Analysis for C ₂₆ H ₂₄ F ₄ N ₂ O ₄ S HCl: calcd: C, 54.50; H, 4.40; N, 4.89; found: C, 54.63; H, 4.41; N, 4.86
470	6-phenyl	phenyl	ISMS 419 (M+1); Analysis for C ₂₆ H ₂₄ F ₄ N ₂ O ₄ S HCl 0.3H ₂ O: calcd: C, 75.65; H, 6.04; N, 6.08; found: C, 75.63; H, 5.89; N, 6.07
470 A	6-butyloxy	2,2,3,3-tetrafluoropropyl	Analysis for C ₂₃ H ₂₇ F ₃ N ₂ O ₂ .HCl.HCl: Calcd: C, 58.52; H, 6.02; N, 5.69; found: C, 58.15; H, 5.64; N, 5.58.

Example 471N-(2-(5-Hydroxy-1H-indol-3-yl)ethyl)-3-(2,2,2-trifluoroethoxy)benzylamine

- 5 Combine N-(2-(5-benzyloxy-1H-indol-3-yl)ethyl)-3-(2,2,2-trifluoroethoxy)benzylamine hydrochloride (295 mg, 0.6 mmol) and 25 mL EtOH and treat with 0.3 mL 5 N HCl and 300 mg 5% Pd/C and hydrogenate at atmospheric pressure overnight. Filter the reaction through a pad of celite and concentrate to dryness then chromatograph on silica gel to give the title compound:
- 10 ISMS 365 (M+1); ¹H NMR (DMSO-d₆) 10.6 (bs, 1H), 9.4 (bs, 2H), 8.75 (s, 1H), 7.45-6.6 (m, 7H), 4.9-4.7 (m, 2H), 4.2 (bs, 2H), 3.2-2.9 (m, 4H).

-121-

By the method of Example 471 the following compounds were prepared and isolated as the hydrochloride except where noted:




No.	Z'	R ₄	Data
472	5-hydroxy	2,2,3,3-tetrafluoro propyl	ISMS 397 (M+1); Analysis for C ₂₀ H ₂₀ F ₄ N ₂ O ₂ HCl H ₂ O: calcd: C, 53.28; H, 5.14; N, 6.21; found: C, 53.31; H, 4.91; N, 6.33
473	6-hydroxy	2,2,2-trifluoro ethyl	ISMS 487 (M+1); ¹ H NMR (DMSO-d ₆) δ10.52 (bs, 1H), 9.36 (s, 2H), 8.91 (s, 1H), 7.38-7.33 (m, 2H), 7.28-7.26 (m, 1H), 7.20-7.18 (m, 1H), 7.09-7.07 (m, 1H), 6.94-6.93 (m, 1H), 6.68-6.67 (m, 1H), 6.50-6.47 (m, 1H), 4.79-4.72 (m, 2H), 4.13 (s, 2H), 3.05-3.02 (m, 4H)
474	6-hydroxy	2,2,3,3-tetrafluoro propyl	ISMS 397 (M+1); Analysis for C ₂₀ H ₂₀ F ₄ N ₂ O ₂ HCl H ₂ O: calcd: C, 53.28; H, 5.14; N, 6.21; found: C, 53.33; H, 4.76; N, 6.12

5

Example 475

N-(2-(5-Carboxy-1H-indol-3-yl)ethyl)-3-(2,2,2-trifluoroethoxy)benzylamine



- Combine N-(2-(5-methoxycarbonyl-1H-indol-3-yl)ethyl)-3-(2,2,2-trifluoroethoxy)benzylamine (200 mg, 0.5 mmol) in 50 mL THF and 1 mL 3 N NaOH. Reflux the mixture overnight, treat with 0.7 mL 5 N HCl and concentrate to dryness. Chromatograph to give the title compound: ISMS 393 (M+1); Analysis for C₂₀H₁₉F₃N₂O₃ CF₃COOH 1.2C₇H₈ 2.1H₂O: calcd: C, 55.76; H, 5.20; N, 4.28; found: C, 55.51; H, 5.47; N, 4.50.

-122-

Example 4803-(3-Fluoropropoxy)benzaldehyde

- Combine 1-bromo-3-fluoropropane (10.0 g, 77.1 mmol) and 3-hydroxybenzaldehyde (10.4 g, 92.5 mmol) in dimethylformamide (220 mL) and stir at room temperature. Treat with potassium carbonate in portions (21.3 g, 144.2 mmol). Heat the reaction mixture at 100°C for 36 hours, then pour into a 1:1 mixture of ice water and dichloromethane. Separate the phases and extract the aqueous layer with additional dichloromethane. Wash the combined organic extracts sequentially with 1.0 N sodium hydroxide, saturated sodium bicarbonate, brine, and then dry over sodium sulfate.
- Filtration and removal of the solvent *in vacuo* provides a residue. Chromatograph the residue on silica gel eluting with 40% ethyl acetate in hexanes to give the title compound as a yellow oil: ¹H NMR (400 MHz, CDCl₃) 9.98 (s, 1H), 7.50-7.42 (m, 2H), 7.42-7.38 (m, 1H), 7.22-7.16 (m, 1H), 4.66 (dt, 2H, J = 46.8, 5.8 Hz), 4.17 (t, 2H, J = 6.0 Hz), 2.19 (d quintuplets, 2H, J = 26.0, 6.0 Hz); MS (APCI): m/e 183.1 (M+1).


Example 4812,2-Difluoroethyltosylate

- Combine *p*-toluenesulfonyl chloride (12.9 g, 67.4 mmol) in pyridine (15 mL) at room temperature and treat dropwise with 2,2-difluoroethanol (5.0 g, 60.9 mmol) via syringe. Stir the reaction mixture under nitrogen for 72 hours, partition between water (20 mL) and dichloromethane (20 mL). Separate the aqueous phase and extract with additional dichloromethane (2x 40 mL). Combine the organic extracts and wash sequentially with 1 N hydrochloric acid (2x 50 mL), sodium bicarbonate (2x 50 mL), and brine (2x 50 mL). Dry the organic layer over sodium sulfate and concentrate *in vacuo* to give the title compound as a yellowish oil: ¹H NMR (300 MHz, CDCl₃): 7.82 (d, 2H, J = 9.0 Hz), 7.40 (d, 2H, J = 9.0 Hz), 5.92 (tt, 1H, J = 55.0, 0.4 Hz), 4.19 (td, 2H, J = 12.6, 4.0 Hz), 2.48 (s, 3H).

Example 4823-(2,2-Difluoroethoxy)benzaldehyde


- The method of Example 480 gives the title compound as a yellow oil. ¹H NMR (400 MHz, CDCl₃): 9.97 (s, 1H), 7.56-7.44 (m, 2H), 7.41-7.37 (m, 1H), 7.21 (ddd, 1H, J = 8.0, 2.8, 1.2 Hz), 6.11 (tt, 1H, J = 55.0, 4.0 Hz), 4.24 (td, 2H, J = 12.6, 4.0 Hz).

-123-

Example 483N-(2-(6-Chloro-1H-indol-3-yl)ethyl)-3-(3-fluoropropoxy)benzylamine

Combine 6-chlorotryptamine (1.4 g, 7.2 mmol), 3-(3-fluoropropoxy)benzaldehyde (1.3 g, 7.2 mmol), and molecular sieves in ethanol (150 mL), and heat at 78°C overnight. Filter the reaction mixture through a plug of celite, and treat the resulting filtrate with sodium borohydride (817 mg, 21.6 mmol) and stir overnight at room temperature. Evaporate the solvent *in vacuo* to give a residue. Chromatograph the residue on silica gel eluting with 9:1 mixture of dichloromethane and 1N ammonia in methanol to give a residue. Chromatograph that residue on 10 g SCX column (wash column with methanol then elute with 1 N ammonia in methanol) and concentrate *in vacuo* to give a light yellow oil. Dissolve the oil in methanol and treated with a methanolic solution of ammonium chloride (112 mg, 2.1 mmol). Sonicate the resulting mixture for 10 minutes, remove the solvent *in vacuo*, and triturate the resulting residue with ether containing a few drops of acetonitrile to give a solid. Collect the solid by filtration to give the title compound as the hydrochloride: mp 177.8-178.9°C; ¹H NMR (400 MHz, dmso-d₆): 11.15 (br s, 1H), 9.41 (br s, 2H), 7.57 (d, 1H, *J* = 8.0 Hz), 7.39 (d, 1H, *J* = 2.0 Hz), 7.32 (t, 1H, *J* = 7.8 Hz), 7.26 (d, 1H, *J* = 2.4 Hz), 7.25-7.21 (m, 1H), 7.11 (d, 1H, *J* = 8.0 Hz), 7.01 (dd, 1H, *J* = 8.8, 2.0 Hz), 6.97 (dd, 1H, *J* = 8.0, 2.0 Hz), 4.60 (dt, 2H, *J* = 47.6, 6.0 Hz), 4.13 (br s, 2H), 4.08 (t, 2H, *J* = 6.4 Hz), 3.10 (br s, 4H), 2.11 (d quintuplets, 2H, *J* = 26.0, 6.0 Hz); MS (ES⁺): *m/e* 361.3 (M+1); CHN (for C₂₀H₂₂ClFN₂O•HCl) calcd: C 60.46, H 5.83, N 7.05; found: C 60.48, H 5.86, N 7.16.

By the method of Example 483 the following compounds were prepared and isolated as the hydrochloride except where noted:




No.	Z'	R ₄	Data
484	6-fluoro	3-fluoro propyl	mp: 174.8-176.0°C; ¹ H NMR (400 MHz, dmso-d ₆): 11.03 (br s, 1H), 9.35 (br s, 2H), 7.52 (dd, 1H, <i>J</i> = 8.8, 5.2 Hz), 7.30 (t, 1H, <i>J</i> = 7.8 Hz), 7.22-7.17 (m, 2H), 7.13-7.06 (m, 2H), 6.95 (dd, 1H, <i>J</i> = 7.8, 2.2 Hz), 6.83 (ddd, 1H, <i>J</i> = 9.6, 8.8, 2.4 Hz), 4.58 (dt, 2H, <i>J</i> = 47.2, 5.8 Hz), 4.11 (s, 2H), 4.06 (t, 2H, <i>J</i> = 6.2 Hz), 3.08 (br s, 4H), 2.08 (d quintuplets, 2H, <i>J</i> = 26.0, 6.0 Hz); MS (ES ⁺): <i>m/e</i> 345.3 (M+1); CHN (for C ₂₀ H ₂₂ F ₂ N ₂ O HCl) calcd: C 63.07, H 6.09, N 7.36; found: C 62.82, H 6.13, N 7.57
485	6-fluoro	2,2-difluoro ethyl	mp 165.0-166.5°C. ¹ H NMR (400 MHz, dmso-d ₆): 11.08 (br s, 1H), 7.56 (dd, 1H, <i>J</i> = 8.7, 5.2 Hz), 7.39-7.31 (m, 2H), 7.21 (d, 1H, <i>J</i> = 2.0 Hz), 7.18 (d, 1H, <i>J</i> = 6.9 Hz), 7.12 (dd, 1H, <i>J</i> = 10.4, 1.7 Hz), 7.04 (dd, 1H, <i>J</i> = 8.7, 1.7 Hz), 6.89-6.81 (m, 1H), 6.42 (tt, 1H, <i>J</i> = 53.9, 3.5 Hz), 4.32 (td, 2H, <i>J</i> = 11.3, 3.2 Hz), 4.14 (s, 2H), 3.20-3.00 (m, 4H); MS: (ES ⁺): <i>m/e</i> 349.0 (M+1)
486	6-chloro	2,2-difluoro ethyl	mp 131.6-133°C: ¹ H NMR (400 MHz, dmso-d ₆): 11.15 (br s, 1H), 9.50 (br s, 2H), 7.57 (d, 1H, <i>J</i> = 8.8 Hz), 7.39 (d, 1H, <i>J</i> = 2.0 Hz), 7.36 (t, 1H, <i>J</i> = 8.2 Hz), 7.32 (br s, 1H), 7.26 (d, 1H, <i>J</i> = 2.0 Hz), 7.17 (d, 1H, <i>J</i> = 7.6 Hz), 7.04 (dd, 1H, <i>J</i> = 7.8, 2.2 Hz), 7.01 (dd, 1H, <i>J</i> = 8.4, 2.0 Hz), 6.41 (tt, 1H, <i>J</i> = 54.4, 3.4 Hz), 4.32 (td, 2H, <i>J</i> = 14.8, 3.6 Hz), 4.14 (br s, 2H), 3.11 (br s, 4H); MS (ES ⁺): <i>m/e</i> 365.3 (M+1); CHN (for C ₁₉ H ₁₉ F ₂ ClN ₂ O•HCl•0.3 H ₂ O) calcd: C 56.11; H 5.11; N 6.89; found: C 56.03; H 4.95; N 7.18
487	6-chloro	2,2,3,3,3-pentafluoro propyl	mp 199.8-201.1°C; ¹ H NMR (400 MHz, dmso-d ₆): 11.15 (br s, 1H), 9.35 (br s, 2H), 7.57 (d, 1H, <i>J</i> = 8.4 Hz), 7.44-7.32 (m, 3H), 7.26 (d, 1H, <i>J</i> = 2.0 Hz), 7.22 (d, 1H, <i>J</i> = 8.0 Hz), 7.10 (dd, 1H, <i>J</i> = 8.4, 2.0 Hz), 7.00 (dd, 1H, <i>J</i> = 8.6, 1.8 Hz), 4.85 (t, 2H, <i>J</i> = 13.2 Hz), 4.13 (s, 2H), 3.10 (br s, 4H); MS (ES ⁺): <i>m/e</i> 433.0 (M+1); CHN (for C ₂₀ H ₁₈ ClF ₅ N ₂ O•0.97HCl) calcd: C 51.31, H 4.08, N 5.98; found: C 51.61, H 4.07, N 6.00
488	5-isopropyl	2,2,3,3,3-pentafluoro propyl	mp 168.5-171.0°C; MS (ES ⁺): <i>m/e</i> 441.1 (M+1); CHN (for C ₂₃ H ₂₅ F ₅ N ₂ O•HCl•0.3H ₂ O) calcd: C 57.28, H 5.56, N 5.81; found: C 57.10, H 5.21, N 6.03

-125-

489	5-isopropyl	2,2,3,3-tetrafluoro propyl	mp 167.0-168.2 °C; ¹ H NMR (400 MHz, dmsd-d ₆): 10.72 (br s, 1H), 7.44 (t, 1H, <i>J</i> = 7.8 Hz), 7.34 (br s, 1H), 7.22-7.15 (m, 2H), 7.14 (br s, 1H Hz), 7.06 (d, 1H, <i>J</i> = 7.6 Hz), 7.01 (dd, 1H, <i>J</i> = 8.4, 1.6 Hz), 6.69 (tt, 1H, <i>J</i> = 51.6, 5.6 Hz), 5.86 (s, 1H), 4.70-4.50 (m, 2H), 3.50-3.25 (m, 4H, overlapping with H ₂ O), 3.17-3.05 (m, 1H), 3.05-2.91 (m, 2H), 1.24 (d, 6H, <i>J</i> = 6.8 Hz); MS (ES ⁺): <i>m/e</i> 422.1 (M+1)
-----	-------------	----------------------------	---

Example 490

N-(2-(6-Chloro-1H-indol-3-yl)ethyl)-N-methyl-3-(2,2-difluoroethoxy)benzylamine




5

Combine N-(2-(6-chloro-1H-indol-3-yl)ethyl)-3-(2,2-difluoroethoxy)benzylamine (276 mg, 0.76 mmol) and formaldehyde (55.5 μ L of a 38% aqueous solution, 0.76 mmol) in dichloroethane (15 mL) and stir at room temperature for 10 minutes; then add in two portions over 10 minutes sodium triacetoxyborohydride (321 mg, 1.51 mmol). Stir the reaction mixture at room temperature overnight and dilute with methanol (10 mL) and quench with one drop of glacial acetic acid. Remove the solvent *in vacuo*, to give a residue, redissolve the crude residue in methanol and directly load onto a 10 g SCX column. After washing the column thoroughly with methanol, elute with 2 N ammonia in methanol. Concentrate *in vacuo* to give the title compound as an oil. Dissolve the oil (239 mg, 0.64 mmol) in methanol (20 mL) and treat with a solution of ammonium chloride (36 mg, 0.67 mmol) in methanol (5 mL). Sonicate the mixture for 10 minutes before removal of the solvent *in vacuo* to give the title compound as the hydrochloride salt. Dissolve the salt in 10 mL of 1:1 acetonitrile-water and lyophilize overnight, providing a fluffy white solid. Triturate the solid with diethyl ether (10 mL) and acetonitrile (2 drops), filter, and dry to give the title compound as the hydrochloride salt: mp: 63.8-65.8°C; ¹H NMR (400 MHz, dmsd-d₆): 11.10 (br s, 1H), 7.52 (d, 1H, *J* = 8.4 Hz) 7.36 (d, 1H, *J* = 2.0 Hz), 7.40-7.26 (m, 2H), 7.22 (d, 1H, *J* = 2.4 Hz), 7.20-7.11 (m, 1H), 7.04 (br d, 1H, *J* = 7.6 Hz), 6.96 (dd, 1H, *J* = 8.6, 1.4 Hz), 6.38 (tt, 1H, *J* = 54.4, 3.6, Hz), 4.50-4.02 (br m, 2H), 4.30 (td, 2H, *J* = 14.4, 3.2 Hz), 3.15 (br s, 4H), 2.68 (br s, 3H); MS (ES⁺): *m/e* 378.9 (M+1).

20

-126-

By the method of Example 490 the following compounds were prepared and isolated as the hydrochloride except where noted:




No.	Z'	R ₄	Data
491	6-fluoro	2,2-difluoro ethyl	mp: 70.8-73.0°C; ¹ H NMR (400 MHz, CDCl ₃): 9.01 (br s, 1H), 7.40-7.35 (m, 1H), 7.35 (dd, 1H, J = 8.8, 5.6 Hz), 7.31-7.25 (m, 1H), 7.10-7.02 (m, 2H), 6.97-6.91 (m, 2H), 6.77 (td, 1H, J = 9.2, 2.0 Hz), 6.05 (tt, 1H, J = 54.8, 4.0 Hz), 4.21 (td, 2H, J = 13.0, 4.0 Hz), 4.08 (br s, 2H), 3.30-3.18 (m, 2H), 3.18-3.05 (m, 2H), 2.66 (s, 3H); MS (APCI): m/e 363.1 (M+1)
492	6-fluoro	3-fluoro propyl	mp: 66.4-69.3 °C; ¹ H NMR (300 MHz, dmsod ₆): 11.04 (s, 1H), 11.20-10.70 (br s, 1H), 7.52 (dd, 1H, J = 8.8, 5.5 Hz), 7.34 (t, 1H, J = 7.9 Hz), 7.30-7.20 (m, 1H), 7.20 (d, 1H, J = 2.2 Hz), 7.12 (AB _q , 2H, J _{AB} = 2.4 Hz, ΔJ _{AB} = 9.8 Hz), 7.00 (br d, 1H, J = 8.4 Hz), 6.84 (ddd, 1H, J = 9.9, 8.8, 2.2 Hz), 4.61 (dt, 2H, J = 47.2, 5.9 Hz), 4.44-4.03 (br m, 2H), 4.08 (t, 2H, J = 6.4 Hz), 3.17 (br s, 4H), 2.68 (br s, 3H), 2.11 (dqintuplets, 2H, J = 25.6, 6.1 Hz); MS (ES ⁺): m/e 358.9 (M+1)
493	6-chloro	3-fluoro propyl	mp: 61.4-63.4°C; ¹ H NMR (400 MHz, DMSO-d ₆): 11.14 (s, 1H), 7.54 (d, 1H, J = 8.4 Hz), 7.40 (d, 1H, J = 2.4 Hz), 7.35 (t, 1H, J = 8.0 Hz), 7.32-7.23 (m, 2H), 7.13 (br d, 1H, J = 7.2 Hz), 7.07-7.00 (m, 1H), 6.99 (dd, 1H, J = 8.6, 1.8 Hz), 4.60 (dt, 2H, J = 46.8, 5.8 Hz), 4.50-4.15 (br m, 2H), 4.08 (t, 2H, J = 6.4 Hz), 3.18 (br s, 4H), 2.72 (br s, 3H), 2.11 (dqintuplets, 2H, J = 26.0, 6.4 Hz); MS (APCI): m/e 375.1 (M+1)
494	6-chloro	2,2,3,3,3-pentafluoro propyl	mp 206.6-207.5 °C; ¹ H NMR (400 MHz, methanol-d ₄): 7.97 (d, 1H, J = 8.0 Hz), 7.93-7.85 (m, 2H), 7.75-7.68 (m, 2H), 7.65 (br d, 1H, J = 7.2 Hz), 7.58 (br d, 1H, J = 8.0 Hz), 7.47 (br d, 1H, J = 9.2 Hz), 5.21 (t, 2H, J = 13.0 Hz), 4.60 (br s, 2H), 3.61 (br s, 4H), 3.14 (br s, 3H); MS (ES ⁺): m/e 447.1 (M+1); CHN (for C ₂₁ H ₂₀ ClF ₅ N ₂ O·HCl) calcd: C 52.19; H 4.38; N 5.80; found: C 52.16; H 4.29; N 5.82


-127-

Example 495

N-(2-(6-Chloro-1H-indol-3-yl)ethyl)-N-isopropyl-3-(2,2,3,3,3-pentafluoropropoxy)benzylamine



- 5 Combine N-(2-(6-Chloro-1H-indol-3-yl)ethyl)-3-(2,2,3,3,3-pentafluoropropoxy)benzylamine (254 mg, 0.59 mmol) in 20 mL of 95:5 methanol-acetic acid, treat with acetone (441 μ L, 5.9 mmol) followed by sodium cyanoborohydride in portions (148 mg, 2.3 mmol). Stir the reaction mixture at 50°C overnight; then at room temperature for an additional 2 days. Remove the solvent *in vacuo* to give a residue.
- 10 Chromatograph the residue on silica gel eluting with 4% methanol in dichloromethane to give the title compound as a colorless oil. Dissolve the oil (237 mg, 0.49 mmol) in methanol (15 mL) and treat with a solution of ammonium chloride (27 mg, 0.49 mmol) in methanol (5 mL). Sonicate the mixture for 10 minutes before concentrating it to a tacky white solid. Dissolve the tacky solid in 10 mL of 1:1 acetonitrile-water and lyophilize to
- 15 give 241 mg (96%) of the title compound as the hydrochloride: mp: 77.0-80.2°C; ^1H NMR (400 MHz, methanol- d_4): 7.31 (br t, 1H, $J = 7.8$ Hz), 7.26-7.21 (m, 1H), 7.16 (br d, 1H, $J = 8.4$ Hz), 7.15-7.07 (m, 2H), 7.05-6.95 (m, 2H), 6.83 (dd, 1H, $J = 8.0, 2.0$ Hz), 4.52 (t, 2H, $J = 12.8$ Hz), 4.12 (br s, 2H), 3.53 (br s, 1H), 3.11 (br s, 2H), 2.89 (br s, 2H), 1.27 (br s, 6H); MS (APCI): m/e 475.1 ($M+1$).
- 20 By the method of Example 495 the following compounds were prepared and isolated as the maleate:




No.	Z'	R	Data
496	6-chloro	propyl	mp 92.4-94.6 °C. Mass (ES ⁺): m/z 475.0 ($M+1$). Elemental Analysis Calculated for $\text{C}_{23}\text{H}_{24}\text{ClF}_5\text{N}_2\text{O} \cdot 1.0\text{C}_4\text{H}_4\text{O}_4 \cdot 0.5\text{H}_2\text{O}$: C, 53.30; H, 4.93; N, 4.57. Found: C, 53.00; H, 4.55; N, 4.86.
497	6-chloro	ethyl	mp 101.0-1-104.0 °C. Mass (ES ⁺): m/z 461.0 ($M+1$).

-128-


Example 500

N-(2-(6-Chloro-5-methoxy-1H-indol-3-yl)ethyl)-3-(2,2,3,3-tetrafluoropropoxybenzylamine




- 5 Combine 5-methoxy-6-chlorotryptamine (0.2 mmol) in dichloromethane (1 mL) and 3-(2,2,3,3-tetrafluoropropoxy)benzaldehyde (0.32 mmol) in dichloromethane (1 mL) and rotate. After 2h, add sodium borohydride (37.83 mg, 1.0 mmol) as a stock solution in dichloromethane (1 mL). After overnight rotation, dilute the reaction mixture with 1 mL of methanol, and apply the resulting solution directly to a 2 g SCX column.
- 10 Thoroughly wash the column with methanol, elute with 2 M ammonia-methanol and concentrate to a residue. If a TLC of eluent, indicates that the reaction was not complete. Dilute the residue with dichloromethane (1 mL) and add a second stock solution of sodium borohydride (37.83 mg, 1.0 mmol) in 1-methyl-2-pyrrolidinone (1mL). After rotation for 2 h, dilute the reaction mixture with 1 mL of methanol, and directly apply the
- 15 resulting solution to a 2 g SCX column. Thoroughly wash the column with methanol, elute with 2 M ammonia-methanol and concentrate to a residue. Further purification on a SI column. Elution with straight ethyl acetate. Compound was characterized using LC method 1 or 2. LCMS R_t 2.749 min at 254 nm, 2.800 min at 220 nm; m/e 445 (M+1).
General LC Methods:
- 20 Method 1: (Shimadzu QP8000) 10 – 90 in 4.5 min. Solvent A: water 0.1% trifluoroacetic acid, Solvent B: acetonitrile 0.1% trifluoroacetic acid. Column: C18 Metachem, monochrom 3 micron, 2.5x25.
Method 2: (Shimadzu) 10 – 80 in 9 min. Solvent A: water 0.1% trifluoroacetic acid, Solvent B: acetonitrile 0.08% trifluoroacetic acid. Column: C18 Metachem, monochrom 5
- 25 micron, 4.6x50.

The following compounds were prepared in a manner similar to Example 500 and isolate as the base unless otherwise indicated:




-129-

No.	Z'	R ₄	Data
501	3-CF ₃	2,2,2-trifluoroethyl	Method 2: LC Rf 3.90 min at 220nm, 3.908 min at 264 nm.
502	3,5-dimethoxy	2,2,2-trifluoroethyl	Method 2: LC Rf 3.620 min at 254 nm, 3.62 min at 220 nm, m/e 367 (M+1).
503	3-chloro	2,2,3,3-tetrafluoropropyl	Method 1: LCMS Rf 2.800 min at 220 nm, m/e 376 (M+1).
504	3-CF ₃	2,2,3,3-tetrafluoropropyl	Method 1: LCMS Rf 2.885 min at 254 nm, m/e 410 (M+1).
506	3-chloro	2-fluoroethyl	Method 2: LC Rf 3.420 min at 254 nm, 3.42 min at 220 nm.
507	3-trifluoromethyl	2-fluoroethyl	Method 2: LC Rf 3.580 min at 254 nm, 3.58 min at 220 nm.
508	3,5-dimethoxy	2-fluoroethyl	Method 2: LC Rf 3.212 min at 254 nm, 3.22 min at 220 nm.
509	3-trifluoromethyl	propyl	Method 2: LC Rf 3.892 min at 254 nm, 3.89 min at 220 nm.
510	2-chloro	phenyl	Method 1: LCMS Rf 2.479 min at 2854 nm, m/e 338 (M+1).
511	3-trifluoromethyl	phenyl	Method 1: LCMS Rf 2.969 min at 254 nm, m/e 372 (M+1).



No.	Z'	R ₄	Data
512	5-methoxy 6-chloro	2,2,2-trifluoroethyl	Method 1: LCMS Rf 2.651 min at 220 nm, m/e 413 (M+1).
513	6-fluoro	2,2,2-trifluoroethyl	Method 1: LCMS Rf 2.618 min at 254 nm, 2.700 min at 220 nm, m/e 367 (M+1).
514	4-chloro 5-methoxy	2,2,2-trifluoroethyl	Method 1: LCMS Rf 2.683 min at 254 nm, 2.661 min at 220 nm, m/e 399 (M+1).
515	5-methoxy 6-chloro	2,2,3,3-tetrafluoropropyl	Method 1: LCMS Rf 2.749 min at 254 nm, 2.800 min at 220 nm, m/e 445 (M+1).
516	6-fluoro	2,2,3,3-tetrafluoropropyl	Method 1: LCMS Rf 2.683 min at 254 nm, 2.661 min at 220 nm, m/e 399 (M+1).
517	4-chloro 5-methoxy	2,2,3,3-tetrafluoropropyl	Method 1: LCMS Rf 2.682 min at 254 nm, 2.663 min at 220 nm, m/e 445 (M+1).
522	5-methoxy	2-fluoroethyl	Method 2: LC Rf 3.19 min at 220 nm.

Example 523


N-(2-(6-Trifluoromethyl-1H-indol-3-yl)ethyl)-3-phenoxybenzylamine

- 5 Combine 5-trifluoromethyltryptamine (0.1 mmol) in methanol (1 mL) and 3-phenoxybenzaldehyde (0.2 mmol) in methanol (1 mL) and rotate. After 3 h, add sodium borohydride (18 mg, 0.5 mmol) as a stock solution in 1-methyl-2-pyrrolidinone (0.5 mL). After overnight rotation, dilute the reaction mixture with 1 mL of methanol, directly apply
- 10 and elute with 2 M ammonia-methanol and concentrate the eluent. Further purification on SI column eluting with ethyl acetate affords the desired compound. Characterization of the compound is achieved by using method 1. LCMS R_f 2.954 min at 254 nm, 2.954 min at 220 nm, m/e 411 ($M+1$).

LC method:

- 15 Method 1: (Shimadzu QP8000) 10 – 90 in 4.5 min. Solvent A: water 0.1% trifluoroacetic acid, Solvent B: acetonitrile 0.1% trifluoroacetic acid. Column: C18 Metachem, monochrom 3 micron, 2.5x25.

The following compounds were prepared following a similar procedure as in Example 523 and isolated as the base unless otherwise indicated:




20

No.	Z'	R ₄	Data
524	6-trifluoro methyl	phenyl	Method 1: LCMS R_f 2.954 min at 254 nm, 2.954 min at 220 nm, m/e 411 ($M+1$).
525	6-fluoro	phenyl	Method 1: LCMS R_f 2.712 min at 254 nm, 2.712 min at 220 nm, m/e 361 ($M+1$).
526	5-methoxy 6-chloro	phenyl	Method 1: LCMS R_f 2.757 min at 254 nm, 2.757 min at 220 nm, m/e 407 ($M+1$).
527	4-chloro 5-methoxy	propyl	Method 1: LCMS R_f 2.578 min at 254 nm, 2.577 min at 220 nm, m/e 373 ($M+1$).
528	6-trifluoro methyl	propyl	Method 1: LCMS R_f 2.850 min at 254 nm, 2.849 min at 220 nm, m/e 377 ($M+1$).
529	6-fluoro	propyl	Method 1: LCMS R_f 2.576 min at 254 nm, 2.576 min at 220 nm, m/e 327 ($M+1$).


-131-

530	5-methoxy 6-chloro	propyl	Method 1: LCMS R _f 2.637 min at 220 nm, m/e 373 (M+1).
-----	-----------------------	--------	---

Example 531N-(2-(4-Sulfonamidophenyl)ethyl)-3-(2,2,3,3,3-pentafluoropropoxy)benzylamine

- 5 Combine 4-sulfonamidophenylethylamine (0.2 mmol) in methanol (1 mL) and 3-(2,2,3,3,3-pentapropylfluororpropoxy)benzaldehyde (0.32 mmol) in methanol (1 mL) and rotate. After 1 hour add sodium borohydride (18 mg, 1.0 mmol) as a stock solution in 1-methyl-2-pyrrolidinone (1 mL). After overnight rotation, dilute the reaction mixture with 1 mL of methanol, and directly apply the resulting solution to a 2 g SCX column. After
- 10 thoroughly washing with methanol, elute the column with 2 M ammonia-methanol and concentrate the eluent to a residue. Further purification by Gilson UV prep system afforded the desired compound and the compound was characterized using method 1. LCMS R_f 2.345 min at 254 nm, 2.347 min at 220 nm, m/e 439 (M+1) 461 (M+22). LC method:
- 15 Method 1: (Shimadzu QP8000) 10 – 90 in 4.5 min. Solvent A: water 0.1% trifluoroacetic acid, Solvent B: acetonitrile 0.1% trifluoroacetic acid. Column: C18 Metachem, monochrom 3 micron, 2.5x25.


- The following compounds were prepared using a similar procedure as in Example
- 20 531 and isolated as the base unless otherwise indicated:



No.	Z'	R ₄	Data
532	2,5-dimethoxy	2,2,3,3,3-pentafluoropropyl	Method 1: LCMS R _f 2.816 min at 254 nm, 2.815 min at 220 nm, m/e 420 (M+1).
533	3,4-dimethoxy	2,2,3,3,3-pentafluoropropyl	Method 1: LCMS R _f 2.634 min at 254 nm, 2.637 min at 220 nm, m/e 420 (M+1).
534	4-sulfonamide	2,2,3,3-tetrafluoropropyl	Method 1: LCMS R _f 2.155 min at 254 nm, 2.156 min at 220 nm, m/e 421 (M+1).

-132-


535	4-sulfonamide	3-fluoropropyl	Method 1: LCMS Rf 1.816 min at 254 nm, 1.818 min at 220 nm, m/e 367 (M+1), 389 (M+22).
537	4-sulfonamide	2-fluoroethyl	Method 1: LCMS Rf 1.606 min at 254 nm, 1.606 min at 220 nm, m/e 375 (M+22).
538	3,4-dimethoxy	phenyl	Method 1: LCMS Rf 2.511 min at 254 nm, 2.511 min at 220 nm, m/e 364 (M+1).
539	4-sulfonamide	2,2-difluoroethyl	Method 1: LCMS Rf 1.782 min at 254 nm, 1.782 min at 220 nm, m/e 371 (M+1), 393 (M+22).
540	2,5-dimethoxy	2,2-difluoroethyl	Method 1: LCMS Rf 2.359 min at 254 nm, m/e 352 (M+1).
541	3,4-dimethoxy	2,2-difluoroethyl	Method 1: LCMS Rf 2.085 min at 254 nm, 2.070 min at 220 nm, m/e 335 (M+1), 352 (M+22).
542	4-sulfonamide	2,2-difluoroethyl	Method 1: LCMS Rf 1.816 min at 254 nm, 1.818 min at 220 nm, m/e 367 (M+1), 389 (M+22).
543	2,5-dimethoxy	3-fluoropropyl	Method 1: LCMS Rf 2.387 min at 254 nm, 2.381 min at 220 nm, m/e 348 (M+1).

Example 545N-(2-(6-Methoxy-1H-indol-3-yl)ethyl)-3-(2,2,3,3,3-pentafluoropropoxy)benzylamine

- 5 Combine amine (0.2 mmol) in 1-methyl-2-pyrrolidinone (1 mL) and aldehyde (0.32 mmol) in dichloromethane (1 mL) and rotate. After 1 h, add sodium borohydride (18 mg, 1.0 mmol) as a stock solution in 1-methyl-2-pyrrolidinone (1 mL). After rotation overnight, dilute the reaction mixture with 1 mL of 10% acetic acid/methanol, and
- 10 directly apply the resulting solution to a 2 g SCX column. Thoroughly wash with methanol, elute the column with 2 M ammonia-methanol and concentrate the eluent to a residue, which was further purified by Gilson UV prep system. Characterize the compound using method 1. LCMS Rf 3.752 min at 254 nm, 3.753 min at 220 nm, m/e 429 (M+1).
- LC method:
- 15 Method 1: (Shimadzu QP8000) 10 – 90 in 4.5 min. Solvent A: water 0.1% trifluoroacetic acid, Solvent B: acetonitrile 0.1% trifluoroacetic acid. Column: C18 Metachem, monochrom 3 micron, 2.5x25.

-133-

The following compounds were prepared by following a similar procedure to Example 545:




No.	Z'	R ₄	Data
546	4-chloro	2,2,3,3,3-pentafluoro propyl	Method 1: LCMS Rf 3.873 min at 254 nm, 3.877 min at 220 nm, m/e 433 (M+1).
547	4-methoxy	2,2,3,3,3-pentafluoro propyl	Method 1: LCMS Rf 3.828 min at 254 nm, 3.833 min at 220 nm, m/e 429 (M+1).
548	5-methoxy 2-methyl	2,2,3,3,3-pentafluoro propyl	Method 1: LCMS Rf 3.802 min at 254 nm, 3.805 min at 220 nm, m/e 433 (M+1).
549	7-methoxy	2,2,3,3,3-pentafluoro propyl	Method 1: LCMS Rf 3.800 min at 254 nm, 3.806 min at 220 nm, m/e 429 (M+1).
550	6-chloro	2,2,3,3,3-pentafluoro propyl	Method 1: LCMS Rf 3.947 min at 254 nm, 3.952 min at 220 nm, m/e 433 (M+1).
551	4-methoxy	2,2,3,3-tetrafluoro propyl	Method 1: LCMS Rf 3.695 min at 254 nm, 3.695 min at 220 nm, m/e 411 (M+1).
552	5-methoxy 2-methyl	2,2,3,3-tetrafluoro propyl	Method 1: LCMS Rf 3.654 min at 254 nm, 3.654 min at 220 nm, m/e 425 (M+1).
553	7-methoxy	2,2,3,3-tetrafluoro propyl	Method 1: LCMS Rf 3.659 min at 254 nm, 3.661 min at 220 nm, m/e 411 (M+1).
554	6-chloro	2,2,3,3-tetrafluoro propyl	Method 1: LCMS Rf 3.821 min at 254 nm, 3.821 min at 220 nm, m/e 415 (M+1).
555	6-methoxy	2-fluoroethyl	Method 1: LCMS Rf 3.169 min at 254 nm, 3.169 min at 220 nm, m/e 345 (M+1).
556	4-chloro	2-fluoroethyl	Method 1: LCMS Rf 3.411 min at 254 nm, 3.412 min at 220 nm, m/e 347 (M+1).
557	4-methoxy	2-fluoroethyl	Method 1: LCMS Rf 3.303 min at 254 nm, 3.304 min at 220 nm, m/e 343 (M+1).
558	5-methoxy 2-methyl	2-fluoroethyl	Method 1: LCMS Rf 3.236 min at 254 nm, 3.236 min at 220 nm, m/e 357 (M+1).
559	7-methoxy	2-fluoroethyl	Method 1: LCMS Rf 3.263 min at 254 nm, 3.264 min at 220 nm, m/e 343 (M+1).

560	6-chloro	2-fluoroethyl	Method 1: LCMS Rf 3.465 min at 254 nm, 3.466 min at 220 nm, m/e 347 (M+1).
561	6-methoxy	2,2-difluoroethyl	Method 1: LCMS Rf 3.190 min at 254 nm, 3.190 min at 220 nm.
562	6-chloro	phenyl	Method 1: LCMS Rf 3.795 min at 254 nm, 3.795 min at 220 nm, m/e 377 (M+1).
563	6-fluoro	2-fluoroethyl	Method 1: LCMS Rf 3.305 min at 254 nm, 3.306 min at 220 nm, m/e 331 (M+1).
571	4-chloro	propyl	Method 1: LCMS Rf 3.668 min at 254 nm, 3.668 min at 220 nm, m/e 343 (M+1).
572	4-methoxy	propyl	Method 1: LCMS Rf 3.581 min at 254 nm, 3.582 min at 220 nm, m/e 339 (M+1).
573	5-methoxy 2-methyl	propyl	Method 1: LCMS Rf 3.524 min at 254 nm, 3.524 min at 220 nm, m/e 353 (M+1).
574	7-methoxy	propyl	Method 1: LCMS Rf 3.553 min at 254 nm, 3.554 min at 220 nm, m/e 339 (M+1).
575	6-chloro	propyl	Method 1: LCMS Rf 3.736 min at 254 nm, 3.736 min at 220 nm, m/e 343 (M+1).
576	4,6-difluoro 5-methoxy	phenyl	Method 1: LCMS Rf 3.830 min at 254 nm, 3.832 min at 220 nm, m/e 423 (M+1).
577	6-methoxy	phenyl	Method 1: LCMS Rf 3.527 min at 254 nm, 3.531 min at 220 nm, m/e 373 (M+1).
578	4-chloro	phenyl	Method 1: LCMS Rf 3.749 min at 254 nm, 3.749 min at 220 nm, m/e 377 (M+1).
579	4-methoxy	phenyl	Method 1: LCMS Rf 3.657 min at 254 nm, 3.658 min at 220 nm, m/e 373 (M+1).
580	5-methoxy -2-methyl	phenyl	Method 1: LCMS Rf 3.609 min at 254 nm, 3.609 min at 220 nm, m/e 3387 (M+1).
581	7-methoxy	phenyl	Method 1: LCMS Rf 3.622 min at 254 nm, 3.622 min at 220 nm, m/e 373 (M+1).
582	6-chloro	phenyl	Method 1: LCMS Rf 3.795 min at 254 nm, 3.795 min at 220 nm, m/e 377 (M+1).
583	4,6-difluoro 5-methoxy	2,2-difluoro ethyl	Method 1: LCMS Rf 3.514 min at 254 nm, 3.519 min at 220 nm, m/e 411 (M+1).
585	4-chloro	2,2-difluoro ethyl	Method 1: LCMS Rf 3.418 min at 254 nm, 3.419 min at 220 nm, m/e 365 (M+1).
586	4-methoxy	2,2-difluoro ethyl	Method 1: LCMS Rf 3.301 min at 254 nm, 3.305 min at 220 nm, m/e 361 (M+1).
587	5-methoxy -2-methyl	2,2-difluoro ethyl	Method 1: LCMS Rf 3.269 min at 254 nm, 3.269 min at 220 nm, m/e 375 (M+1).
588	7-methoxy	2,2-difluoro ethyl	Method 1: LCMS Rf 3.265 min at 254 nm, 3.271 min at 220 nm, m/e 361 (M+1).
589	6-chloro	2,2-difluoro ethyl	Method 1: LCMS Rf 3.476 min at 254 nm, 3.476 min at 220 nm, m/e 365 (M+1).

-135-

590	6-fluoro	2,2-difluoro ethyl	Method 1: LCMS Rf 3.326 min at 254 nm, 3.326 min at 220 nm, m/e 349 (M+1).
592	6-methoxy	3-fluoropropyl	Method 1: LCMS Rf 3.170 min at 254 nm, 3.176 min at 220 nm, m/e 357 (M+1).
593	4-chloro	3-fluoropropyl	Method 1: LCMS Rf 3.400 min at 254 nm, 3.407 min at 220 nm, m/e 361 (M+1).
594	4-methoxy	3-fluoropropyl	Method 1: LCMS Rf 3.326 min at 254 nm, 3.327 min at 220 nm, m/e 357 (M+1).
595	5-methoxy -2-methyl	3-fluoropropyl	Method 1: LCMS Rf 3.277 min at 254 nm, 3.277 min at 220 nm, m/e 371 (M+1).
596	7-methoxy	3-fluoropropyl	Method 1: LCMS Rf 3.290 min at 254 nm, 3.291 min at 220 nm, m/e 357 (M+1).
597	6-chloro	3-fluoropropyl	Method 1: LCMS Rf 3.498 min at 254 nm, 3.499 min at 220 nm, m/e 361 (M+1).
598	6-fluoro	3-fluoropropyl	Method 1: LCMS Rf 3.329 min at 254 nm, 3.330 min at 220 nm, m/e 345 (M+1).
600	6-methoxy	2,2,2-trifluoroethyl	Method 1: LCMS Rf 3.288 min at 254 nm, 3.228 min at 220 nm, m/e 379 (M+1).
601	4-chloro	2,2,2-trifluoroethyl	Method 1: LCMS Rf 3.518 min at 254 nm, 3.518 min at 220 nm, m/e 383 (M+1).
602	4-methoxy	2,2,2-trifluoroethyl	Method 1: LCMS Rf 3.427 min at 254 nm, 3.428 min at 220 nm, m/e 379 (M+1).
603	5-methoxy -2-methyl	2,2,2-trifluoroethyl	Method 1: LCMS Rf 3.378 min at 254 nm, 3.378 min at 220 nm, m/e 393 (M+1).
604	7-methoxy	2,2,2-trifluoroethyl	Method 1: LCMS Rf 3.234 min at 254 nm, 3.255 min at 220 nm, m/e 379 (M+1).
605	6-chloro	2,2,2-trifluoroethyl	Method 1: LCMS Rf 3.587 min at 254 nm, 3.587 min at 220 nm, m/e 383 (M+1).


The following compounds were prepared by following a similar procedure to Example 545:



No.	Z'	R ₄	Data
606	6-methoxy	2,2-difluoroethyl	Method 1: LCMS Rf 3.190 min at 254 nm, 3.190 min at 220 nm.
607	4-fluoro 5-methoxy 6-fluoro	3-fluoropropyl	Method 1: LCMS Rf 3.390 min at 254 nm, 3.395 min at 220 nm, m/e 401 (M+1).

-136-

608	4-fluoro 6-fluoro 5-methoxy	2,2,2- trifluoroethyl	Method 1: LCMS R _f 3.442 min at 254 nm, 3.453 min at 220 nm, m/e 429 (M+1).
-----	-----------------------------------	--------------------------	---


Example 620N-(2-(5-Methoxy-1H-indol-3-yl)ethyl)-3-(2,2,3,3,3-pentafluoropropoxy)benzylamine

- 5 Combine amine (0.2 mmol) in dichloromethane (0.5 mL) and aldehyde (0.4 mmol) in dichloromethane (1 mL) and rotate. After 1h, add sodium triacetoxyborohydride (82 mg, 0.8 mmol) as a stock solution in 1-methyl-2-pyrrolidinone (1 mL) and rotate. After overnight rotation, dilute the reaction mixture with 1 mL of methanol and directly apply to a 2 g SCX column. After thoroughly washing with methanol, elute the column with 2
- 10 M ammonia-methanol and concentrate the eluent to a residue, which was further purified by Gilson UV prep system. Compound was characterized using method 3. LCMS R_f 4.823 min at 254 nm, 4.823 min at 220 nm, m/e 443 (M+1).

LC Method:


- Method 3: (Shimadzu QP8000) 5 – 90 in 4.5 min. Solvent A: water 0.1% trifluoroacetic acid, Solvent B: acetonitrile 0.1% trifluoroacetic acid. Column: C18 Metachem,
- 15 monochrom 3 micron, 2.5x25.

The following compounds were prepared using a similar procedure as in Example 620:



No.	R ⁴	Data
622	2,2,3,3-tetrafluoropropyl	Method 3: LCMS R _f 4.681 min at 254 nm, 4.692 min at 220 nm, m/e 425 (M+1).
623	2,2,2-trifluoroethyl	Method 3: LCMS R _f 4.639 min at 254 nm, 4.643 min at 220 nm, m/e 393 (M+1).

-137-


Example 624N-(2-(6-Fluoro-1-methyl-1H-indol-3-yl)ethyl)-N-methyl-3-propoxybenzylamine

- Combine N-methyl-N-(2-(6-fluoro-1-methyl-1H-indol-3-yl)ethylamine (0.2 mmol) in 1-methyl-2-pyrrolidinone (0.5 mL) and 3-propyloxybenzaldehyde (0.32 mmol) in dichloromethane (1 mL) and rotate. After overnight rotation, add sodium borohydride (1.0 mmol) as a stock solution in 1-methyl-2-pyrrolidinone (0.5 mL) and rotate. After rotation for 3 h, dilute the reaction mixture with 1 mL of 10% acetic acid/methanol, and directly apply the resulting solution to a 2 g SCX column. After thoroughly washing with methanol, elute the column with 2 M ammonia-methanol and concentrate the eluent to a residue, which was further purified by Gilson UV prep system.

LC method:

- Method 1: (Shimadzu QP8000) 10 – 90 in 4.5 min. Solvent A: water 0.1% trifluoroacetic acid, Solvent B: acetonitrile 0.1% trifluoroacetic acid. Column: C18 Metachem, monochrom 3 micron, 2.5x25.


The following compounds were prepared following a similar procedure as found in Example 624:



	Z'	R ₃	R ₄	Data
625	3-trifluoromethyl 4-fluoro	4-CH ₃	propyl	Method 1: LCMS Rf 3.214 min at 254 nm, 3.213 min at 220 nm, m/e 371 (M+1).
626	3-trifluoromethyl 4-fluoro	H	3,3,3-trifluoropropyl	Method 1: LCMS Rf 3.042 min at 254 nm, 3.042 min at 220 nm, m/e 410 (M+1).
627	3-trifluoromethyl 4-fluoro	H	2,2-difluoro ethyl	Method 1: LCMS Rf 2.828 min at 254 nm, 2.828 min at 220 nm, m/e 378 (M+1).
628	3-trifluoromethyl 4-fluoro	H	2,2,3,3,3-penta fluoro propyl	Method 1: LCMS Rf 3.196 min at 254 nm, 3.196 min at 220 nm, m/e 446 (M+1).

-138-


629	3-trifluoromethyl 4-fluoro	H	2,2,2- trifluoroethyl	Method 1: LCMS Rf 2.984 min at 254 nm, 2.984 min at 220 nm, m/e 396 (M+1).
630	3-trifluoromethyl 4-fluoro	H	3-fluoro propyl	Method 1: LCMS Rf 2.855 min at 254 nm, 2.855 min at 220 nm, m/e 374 (M+1).



No.	Z'	R ₃	R ₄	Data
632	5-fluoro 6-chloro	4- methyl	propyl	Method 1: LCMS Rf 3.141 min at 254 nm, 3.140 min at 220 nm, m/e 375 (M+1).
633	6- trifluoro methyl	H	3,3,3-trifluoro propyl	Method 1: LCMS Rf 3.065 min at 254 nm, 3.066 min at 220 nm, m/e 431 (M+1).
634	5-fluoro 6-chloro	H	3,3,3-trifluoro propyl	Method 1: LCMS Rf 2.977 min at 254 nm, 2.977 min at 220 nm, m/e 415 (M+1).
635	5,6- difluoro	H	3,3,3-trifluoro propyl	Method 1: LCMS Rf 2.871 min at 254 nm, 2.872 min at 220 nm, m/e 399 (M+1).
636	6- trifluoro methyl	H	3,3,3-trifluoro propyl	Method 1: LCMS Rf 3.065 min at 254 nm, 3.066 min at 220 nm, m/e 431 (M+1).
637	5-fluoro 6-chloro	H	2,2-difluoro ethyl	Method 1: LCMS Rf 2.782 min at 254 nm, 2.782 min at 220 nm, m/e 383 (M+1).
638	5,6- difluoro	H	2,2-difluoro ethyl	Method 1: LCMS Rf 2.655 min at 254 nm, 2.655 min at 220 nm, m/e 367 (M+1).
639	6- trifluoro methyl	H	2,2-difluoro ethyl	Method 1: LCMS Rf 2.876 min at 254 nm, 2.875 min at 220 nm, m/e 399 (M+1).
640	6- trifluoro methyl	H	2,2,2 trifluoro ethyl	Method 1: LCMS Rf 3.009 min at 254 nm, 3.009 min at 220 nm, m/e 417 (M+1).
641	5-fluoro 6-chloro	H	2,2,3,3,3- pentafluoro propyl	Method 1: LCMS Rf 3.135 min at 254 nm, 3.135 min at 220 nm, m/e 451 (M+1).
642	5,6- difluoro	H	2,2,3,3,3- pentafluoro propyl	Method 1: LCMS Rf 3.027 min at 254 nm, 3.027 min at 220 nm, m/e 435 (M+1).
643	6- trifluoro methyl	H	2,2,3,3,3- pentafluoro propyl	Method 1: LCMS Rf 3.202 min at 254 nm, 3.202 min at 220 nm, m/e 467 (M+1).
645	5,6- difluoro	H	2,2,2-trifluoro ethyl	Method 1: LCMS Rf 2.982 min at 254 nm, 2.982 min at 220 nm, m/e 396 (M+1).

-139-


646	6-trifluoro methyl	H	2,2,2-trifluoroethyl	Method 1: LCMS Rf 3.009 min at 254 nm, 3.009 min at 220 nm, m/e 417 (M+1).
647	5-fluoro 6-chloro	H	3-fluoro propyl	Method 1: LCMS Rf 2.796 min at 254 nm, 2.796 min at 220 nm, m/e 379 (M+1).
648	5,6-difluoro	H	3-fluoro propyl	Method 1: LCMS Rf 2.644 min at 254 nm, 2.646 min at 220 nm, m/e 363 (M+1).
649 A	6-trifluoro methyl	H	3-fluoro propyl	Method 1: LCMS Rf 2.900 min at 254 nm, 2.900 min at 220 nm, m/e 395 (M+1).



No.	Z'	R ₃	R ₄	Data
631	6-fluoro	4-methyl	propyl	Method 1: LCMS Rf 3.152 min at 220 nm, m/e 355 (M+1).
633 A	6-fluoro	H	3,3,3-trifluoro propyl	Method 1: LCMS Rf 2.949 min at 254 nm, 2.953 min at 220 nm, m/e 395 (M+1).
640 A	6-fluoro	H	2,2,3,3,3-pentafluoro propyl	Method 1: LCMS Rf 3.112 min at 254 nm, 3.117 min at 220 nm, m/e 431 (M+1).
649	6-fluoro	H	2,2,2-trifluoro ethyl	Method 1: LCMS Rf 2.895 min at 254 nm, 2.898 min at 220 nm, m/e 381 (M+1).

Example 650

5 N-2-(3-chlorophenyl)ethyl-3-hydroxybenzylamine



Combine 2-(3-chlorophenyl)ethylamine (1.866 gm; 15.28 mmol) and 3-hydroxybenzaldehyde (1.567 gm; 10.07 mmol) in 40 mL of methanol and stir at room temperature for 20 min and treat with sodium borohydride (0.950 gm; 25.1 mmol) in one portion. Stir the mixture at room temperature. After 15 h, add water (10 mL), and remove

-140-

the methanol by rotary evaporation. Add to this slurry water (25 mL) and dichloromethane (50 mL), separate the layers and extract the aqueous layer with dichloromethane (50 mL). Wash the combined organic layers with saturated brine (3x), dry over MgSO_4 , and concentrate to give the title compound.

5

Example 650AN-t-Butoxycarbonyl-N-2-(3-chlorophenyl)ethyl-3-hydroxybenzylamine


Combine N-2-(3-chlorophenyl)ethyl-3-hydroxybenzylamine, dichloromethane (40 mL), and di-*tert*-butyl dicarbonate (1.556 gm; 7.131 mmol) and triethylamine (1.0 mL; 7.2 mmol). After 18 hours, pour into water (50 mL), separate the layers, and extract the aqueous layer with dichloromethane. Wash combined organic layers with water, dry over MgSO_4 , and concentrate. Chromatograph on silica gel eluting with 5% ethyl acetate in hexanes to give the title compound.

15

Example 651N-t-Butoxycarbonyl-N-2-(3-chlorophenyl)ethyl-3-propoxybenzylamine

Add a solution of 50% sodium hydroxide in water (0.8 mL) to a solution of N-t-butoxycarbonyl-N-2-(3-chlorophenyl)ethyl-3-hydroxybenzylamine (46.7 mg, 0.129 mmol), n-propyl iodide (0.17 gm, 1.00 mmol), and tetrabutylammonium bromide (18 mg, 0.057 mmol) in toluene (1 mL). Stir the mixture at 1200 rpm and heat at 50-54 °C. After 64.5 hour pour the mixture into 5 mL of water, separate the phases and extract the aqueous phase twice with dichloromethane. Combine the organic phases and wash with saturated sodium bicarbonate solution, and then saturated brine, dry (MgSO_4), and concentrate to give a residue. Chromatograph the residue on silica gel to give the title compound: MS (ES⁺): m/e (M+1) 404. TLC (20% EtOAc in hexanes, R_f 0.54).

25

Example 652N-(2-(3-Chlorophenyl)ethyl)-3-propoxybenzylamine

30


Add methanesulfonic acid (70 uL) to a solution of N-t-Butoxycarbonyl-N-2-(3-chlorophenyl)ethyl-3-propoxybenzylamine in dichloromethane (4 mL) and agitate the mixture for 3 hours at room temperature. Add 10% aqueous Na_2CO_3 (2 mL), separate

-141-

the layers, and concentrate the organic layer in a nitrogen stream to give a residue.

Dissolve the residue in 4 mL of 5% acetic acid in methanol and pass through a 1 gm SCX column, eluting with 1 M ammonia in methanol to give the title compound: MS (ES+): m/e (M+1). HPLC (10-90% water/acetonitrile over 7.5 min, Tr = 4.490 min).

- 5 The following compounds were prepared by a similar procedure to Examples 651 and 652:




No.	R ₄	Data
654	ethyl	Method 1: LCMS Rf 4.223 min at 254/220 nm; m/e 298.9 (M+1)
655	butyl	Method 1: LCMS Rf 4.715 min at 254/220 nm; m/e 317.9 (M+1)
656	hexyl	Method 1: LCMS Rf 5.137 min at 254/220 nm; m/e 345.9 (M+1)
658	allyl	Method 1: LCMS Rf 4.373 min at 254/220 nm; m/e 301.9 (M+1)
660	pyridin-2-ylmethyl	Method 1: LCMS Rf 3.547 min at 254/220 nm; m/e 352.9 (M+1)
661	pyridin-3-ylmethyl	Method 1: LCMS Rf 3.487 min at 254/220 nm; m/e 352.9 (M+1)
662	pyridin-4-ylmethyl	Method 1: LCMS Rf 3.455 min at 254/220 nm; m/e 352.9 (M+1)

10

Example 665

N-(2-(5-Methoxy-1-ethyl-1H-indol-3-yl)ethyl)-N-ethyl-3-phenyloxybenzylamine




- 15 Add acetaldehyde (0.080 mL; 0.77 mmol) to a solution of N-(2-(5-methoxy-1H-indol-3-yl)ethyl)-3-phenyloxybenzylamine (free base, 55.5 mg, 0.149 mmol) in dichloromethane (1 mL) followed by a suspension of sodium triacetoxymethylborohydride (64 mg; 0.30 mmol) in dichloromethane (1 mL). After 44 hours, quench by the addition of methanol (0.5 mL) and concentrate in a stream of nitrogen to give a residue. Dissolve the residue in 4 mL of 5% acetic acid in methanol and partially purify by passage through a 1 gm SCX column, eluting with
- 20 1 M ammonia in methanol to give a residue. Chromatograph the residue by preparative HPLC (C-18 column, flow rate of 20 ml/min, 5-90 % water/acetonitrile

-142-


over 12 min) to give the title compound: MS (ES⁺): m/e (M+1); HPLC: (10-90% water/acetonitrile over 10 min, Tr = 5.25 min).

The following compounds were prepared following a similar procedure in Example

5 665:



No.	R ₂	Data
666	methyl	LC Method 2: R _f 5.12 min at 254/220 nm; m/e 351.9 (M+1)
667	ethyl	LC Method 2: R _f 5.25 min at 254/220 nm; m/e 365.9 (M+1)



No.	R ₂	Data
668	ethyl	LC Method 2: R _f 4.98 min at 254/220 nm; m/e 401.09 (M+1)

10

Example 670

3-Propoxybenzaldehyde

Combine 3-hydroxybenzaldehyde (7.50 gm; 61.4 mmol), n-propyl iodide (17.3 gm; 102 mmol), and potassium carbonate (16.90 gm; 122 mmol) in 2-butanone (100 mL) and reflux. After 17 h, allow the mixture to cool to room temperature, decant the solution and concentrate by rotary evaporation. Partition the residue between diethyl ether (150 mL) and water (150 mL), separate the layers and extract the aqueous layer with diethyl ether (2 x 100 mL). Combine organic layers and wash with water, 1 N NaOH, and then water, dry over MgSO₄, and concentrate to give a residue. Distill the residue to give the title compound: bp: 122-125 °C (15 mm); TLC (10% Et₂O/hexanes; R_f 0.35).

20

Example 671

3-(3,3,3-Trifluoropropoxy)benzaldehyde

Cool a mixture of toluenesulfonyl chloride (7.43 gm; 39.0 mmol) and pyridine (50 mL) to 0 °C, add 3,3,3-trifluoropropanol (2.23 gm; 19.5 mmol) and store the mixture at 3 °C. After 48 hour pour the reaction mixture into 350 mL of ice water and

25

-143-

extract with diethyl ether (3 x 125 mL). Combine the organic layers and wash with 5 N HCl, water, saturated sodium bicarbonate solution, and brine, dry over MgSO₄, and concentrate to give 3,3,3-trifluoropropyl tosylate. The material was carried into the next step without purification.

- 5 Combine 3,3,3-trifluoropropyl tosylate (4.057 gm; 15.12 mmol), 3-hydroxybenzaldehyde (1.85 gm; 15.12 mmol), and K₂CO₃ (4.15 gm; 30.0 mmol) in DMF (80 mL) and heat at 100 °C. After 18 hours, cool to room temperature, dilute with water (200 mL) and extract with dichloromethane (2 x 200 mL). Combine organic extracts and wash sequentially with water (100 mL), 0.1 M NaOH (2 x 100 mL), saturated sodium
10 bicarbonate (100 mL) and saturated brine (100 mL), dry (MgSO₄), and concentrate. Chromatography on silica gel (0-20% ethyl acetate in hexane) to give the title product.


Example 672

3-(2-Fluoroethoxy)benzaldehyde

- 15 Combine 1-bromo-2-fluoroethane (4.575 g; 36.0 mmol), 3-hydroxybenzaldehyde (4.103 gm; 33.60 mmol), and K₂CO₃ (7.05 gm; 51.0 mmol) in 2-butanone (100 mL) and reflux. After 18 hour cool the mixture to ambient temperature, concentrate, and partition between 100 mL of water and 100 mL of dichloromethane. Separate the layers and extract the aqueous layer with dichloromethane (2 x 75 mL). Combine the organic layers and
20 wash sequentially with brine (2 x 150 mL), 1 M NaOH (2 x 100 mL), NaHCO₃ (saturated, 100 mL), and brine (150 mL), dried (MgSO₄), concentrate, and chromatograph on silica gel (0-25% diethyl ether in hexanes) to give the title compound.

Example 673

- 25 N-(2-(5-Fluoro-1H-indol-3-yl)ethyl)-3-propoxybenzylamine



- Combine 3-propoxybenzaldehyde (29.6 mg; 0.18 mmol) and 5-fluorotryptamine (14.2 mg; 0.080 mmol) in methanol (2 mL). Add a solution of sodium borohydride in diglyme (1 ml of a 0.5 M solution; 0.50 mmol) and agitate. After 63 h at room
30 temperature concentrate in a stream of nitrogen. Dissolve the residue in methanol and add to a 1 gm SCX column previously rinsed with 5% acetic acid in methanol. Elute the product from the SCX column with 1 M ammonia in methanol to give the title compound: MS (ES⁺): m/e (M+1); HPLC (10-90% water/acetonitrile over 10 min, Tr = 4.08 min.

-144-

General LC Methods:


Method 1: (Shimadzu Class VP HPLC and Micromass Platform LC with HP1100 LC system) 10 – 90 in 7.5 min. Solvent A: water 0.1% trifluoroacetic acid, Solvent B: acetonitrile 0.1% trifluoroacetic acid. Column: C18 Metachem, monochrom 3 micron, 2.5x25.

Method 2: (Shimadzu Class VP HPLC and Micromass Platform LC with HP1100 LC system) 10 – 90 in 10 min. Solvent A: water 0.1% trifluoroacetic acid, Solvent B: acetonitrile 0.1% trifluoroacetic acid. Column: C18 Metachem, monochrom 3 micron, 2.5x25.

Method 3: (Waters Millennium HPLC and Micromass Platform LC with HP1100 LC system) 10-100 in 10 min. Solvent A: 0.1% trifluoroacetic acid, Solvent B: acetonitrile 0.08% trifluoroacetic acid. Column: YMC, 5 micron, 2.5x25.

Method 4: (Shimadzu QP8000) 10 – 90 in 4.5 min. Solvent A: water 0.1% trifluoroacetic acid, Solvent B: acetonitrile 0.1% trifluoroacetic acid. Column: C18 Metachem, monochrom 3 micron, 2.5x25.

The following compounds were prepared following a procedure following Example 673:




No.	Z'	Data
675	2-fluoro	LC Method 3: Rf 4.18 min at 254/220 nm; m/e 322.0 (M+1)
676	3-fluoro	LC Method 3: Rf 4.23 min at 254/220 nm; m/e 322.0 (M+1)
677	4-chloro	LC Method 3: Rf 4.48 min at 254/220 nm; m/e 337.9 (M+1)
678	4-hydroxy	LC Method 3: Rf 3.62 min at 254/220 nm; m/e 320.0 (M+1)
679	2-methoxy	LC Method 3: Rf 4.30 min at 254/220 nm; m/e 334.0 (M+1)
680	4-bromo 3-methoxy	LC Method 3: Rf 4.50 min at 254/220 nm; m/e 411.9 (M+1)
681	4-fluoro	LC Method 3: Rf 4.22 min at 254/220 nm; m/e 322.0 (M+1)
682	2-chloro	LC Method 3: Rf 4.36 min at 254/220 nm; m/e 338.0 (M+1)
683	4-bromo	LC Method 3: Rf 4.55 min at 254/220 nm; m/e 383.91 (M+1)
684	4-methyl	LC Method 3: Rf 4.42 min at 254/220 nm; m/e 318.0 (M+1)
685	3-methoxy	LC Method 3: Rf 4.19 min at 254/220 nm; m/e 334.0 (M+1)
686	4-methoxy	LC Method 3: Rf 4.15 min at 254/220 nm; m/e 334.0 (M+1)
687	2-ethoxy	LC Method 3: Rf 4.55 min at 254/220 nm; m/e 348.0 (M+1)
688	4-ethoxy	LC Method 3: Rf 4.43 min at 254/220 nm; m/e 348.0 (M+1)
689	4-phenoxy	LC Method 3: Rf 5.00 min at 254/220 nm; m/e 396.0 (M+1)

-145-


690	4-sulfonamide	LC Method 3: Rf 3.46 min at 254/220 nm; m/e 383.0 (M+1)
691	3,4-dichloro	LC Method 3: Rf 4.74 min at 254/220 nm; m/e 372.0 (M+1)
692	2,5-dichloro	LC Method 3: Rf 4.74 min at 254/220 nm; m/e 372.0 (M+1)
693	2,6-dichloro	LC Method 3: Rf 4.51 min at 254/220 nm; m/e 372.0 (M+1)
694	2,5-dimethoxy	LC Method 3: Rf 4.31 min at 254/220 nm; m/e 364.0 (M+1)
695	2,3-dimethoxy	LC Method 3: Rf 4.24 min at 254/220 nm; m/e 364.0 (M+1)
696	3,5-dimethoxy	LC Method 3: Rf 4.26 min at 254/220 nm; m/e 364.0 (M+1)
697	3-ethoxy-4-methoxy	LC Method 3: Rf 4.14 min at 254/220 nm; m/e 378.0 (M+1)

The following compounds were prepared following a procedure following Example 673:



5

No.	Z'	Data
698	5-methyl	LC Method 4: Rf 2.852 min at 254/220 nm; m/e 357 (M+1)
699	5-chloro	LC Method 4: Rf 2.893 min at 254/220 nm; m/e 377 (M+1)




No.	Z'	Data
700	2-fluoro	LC Method 3: Rf 3.90 min at 254/220 nm; m/e 288.0 (M+1)
701	3-fluoro	LC Method 3: Rf 3.95 min at 254/220 nm; m/e 288.0 (M+1)
702	4-fluoro	LC Method 3: Rf 3.96 min at 254/220 nm; m/e 288.0 (M+1)
703	2-chloro	LC Method 3: Rf 4.23 min at 254/220 nm; m/e 303.9 (M+1)
704	4-chloro	LC Method 3: Rf 4.12 min at 254/220 nm; m/e 303.9 (M+1)
705	4-bromo	LC Method 3: Rf 4.33 min at 254/220 nm; m/e 347.9 (M+1)
706	4-methyl	LC Method 3: Rf 4.17 min at 254/220 nm; m/e 284.0 (M+1)
707	4-hydroxy	LC Method 3: Rf 3.26 min at 254/220 nm; m/e 286.0 (M+1)
708	2-methoxy	LC Method 3: Rf 4.03 min at 254/220 nm; m/e 300.0 (M+1)
709	3-methoxy	LC Method 3: Rf 3.91 min at 254/220 nm; m/e 300.0 (M+1)
710	4-methoxy	LC Method 3: Rf 3.91 min at 254/220 nm; m/e 300.0 (M+1)
711	3-ethoxy	LC Method 3: Rf 4.31 min at 254/220 nm; m/e 314.0 (M+1)

-146-

712	4-ethoxy	LC Method 3: Rf 4.14 min at 254/220 nm; m/e 314.0 (M+1)
713	4-phenoxy	LC Method 3: Rf 4.77 min at 254/220 nm; m/e 362.0 (M+1)
714	4-sulfonamide	LC Method 3: Rf 3.06 min at 254/220 nm; m/e 349.0 (M+1)
715	3,4-dichloro	LC Method 3: Rf 4.52 min at 254/220 nm; m/e 337.9 (M+1)
716	2,5-dichloro	LC Method 3: Rf 4.51 min at 254/220 nm; m/e 337.9 (M+1)
717	2,6-dichloro	LC Method 3: Rf 4.28 min at 254/220 nm; m/e 337.9 (M+1)
718	3,4-dimethoxy	LC Method 3: Rf 3.59 min at 254/220 nm; m/e 330.0 (M+1)
719	2,5-dimethoxy	LC Method 3: Rf 4.04 min at 254/220 nm; m/e 330.0 (M+1)
720	2,3-dimethoxy	LC Method 3: Rf 3.96 min at 254/220 nm; m/e 330.0 (M+1)
721	3,5-dimethoxy	LC Method 3: Rf 3.99 min at 254/220 nm; m/e 330.0 (M+1)
722	3-bromo 4-methoxy	LC Method 3: Rf 4.22 min at 254/220 nm; m/e 379.9 (M+1)
723	4-ethoxy-3- methoxy	LC Method 3: Rf 3.88 min at 254/220 nm; m/e 344.0 (M+1)
724	3-ethoxy-4- methoxy	LC Method 3: Rf 3.84 min at 254/220 nm; m/e 344.0 (M+1)


The following compounds were prepared following a procedure following Example 673:



No.	R ₁	Data
725	pyridine-2-yl	LC Method 3: Rf 2.38 min at 254/220 nm; m/e 271.0 (M+1)
726	pyridin-3-yl	LC Method 3: Rf 2.25 min at 254/220 nm; m/e 271.0 (M+1)
727	pyridin-4-yl	LC Method 3: Rf 2.21 min at 254/220 nm; m/e 271.0 (M+1)
729	7-methyl-1H-indol-3-yl	LC Method 3: Rf 4.19 min at 254/220 nm; m/e 323.0 (M+1)
730	6-methoxy-1H-indol-3-yl	LC Method 3: Rf 3.90 min at 254/220 nm; m/e 339.0 (M+1)
731	thiophen-3-yl	LC Method 3: Rf 3.70 min at 254/220 nm; m/e 275.9 (M+1)
732	5-methyl-1H-indol-3-yl	LC Method 4: Rf 2.680 min at 254/220 nm; m/e 323 (M+1)
733	5-chloro-1H-indol-3-yl	LC Method 4: Rf 4.019 min at 254/220 nm; m/e 344 (M+1)

5

The following compounds were prepared following a procedure following Example 673:




-147-

No.	Z'	R ₄	Data
734	5-methyl	2-fluoroethyl	LC Method 4: Rf 2.381 min at 254/220 nm; m/e 327 (M+1)
735	5-fluoro	2-fluoroethyl	LC Method 4: Rf 2.300 min at 254/220 nm; m/e 331 (M+1)
736	5-methyl	2,2-difluoroethyl	LC Method 4: Rf 2.520 min at 254/220 nm; m/e 345 (M+1)
737	5-fluoro	2,2difluoroethyl	LC Method 4: Rf 2.445 min at 254/220 nm; m/e 349 (M+1)
738	5-chloro	2,2-difluoroethyl	LC Method 4: Rf 2.598 min at 254/220 nm; m/e 365 (M+1)
739	5-fluoro	4,4,4-trifluorobutyl	LC Method 4: Rf 3.017 min at 254/220 nm; m/e 395 (M+1)
740	5-fluoro	2,2,2-trifluoroethyl	LC Method 4: Rf 2.787 min at 254/220 nm; m/e 367 (M+1)
741	5-methoxy	2,2,2-trifluoroethyl	LC Method 4: Rf 2.681 min at 254/220 nm; m/e 379 (M+1)
742	5-chloro	4,4,4-trifluorobutyl	LC Method 4: Rf 3.151 min at 254/220 nm; m/e 411 (M+1)
743	5-fluoro	3-fluoropropyl	LC Method 4: Rf 2.475 min at 254/220 nm; m/e 345 (M+1)
744	5-methoxy	3,3,3-trifluoropropyl	LC Method 4: Rf 2.889 min at 254/220 nm; m/e 393 (M+1)
745	5-chloro	3-fluoropropyl	LC Method 4: Rf 2.628 min at 254/220 nm; m/e 361 (M+1)
746	5-fluoro	2,2,3,3-tetrafluoropropyl	LC Method 4: Rf 2.680 min at 254/220 nm; m/e 399 (M+1)
747	5-methyl	2,2,3,3-tetrafluoropropyl	LC Method 4: Rf 2.756 min at 254/220 nm; m/e 397 (M+1)
748	5-chloro	2,2,3,3-tetrafluoropropyl	LC Method 4: Rf 2.820 min at 254/220 nm; m/e 417 (M+1)
750	5-fluoro	2,2,3,3,3-pentafluoropropyl	LC Method 4: Rf 2.833 min at 254/220 nm; m/e 417 (M+1)
751	5-methyl	2,2,3,3,3-pentafluoropropyl	LC Method 4: Rf 2.908 min at 254/220 nm; m/e 415 (M+1)
752	5-chloro	2,2,3,3,3-pentafluoropropyl	LC Method 4: Rf 2.784 min at 254/220 nm; m/e 433 (M+1)
754	5-methyl	3-fluoropropyl	LC Method 4: Rf 2.457 min at 254/220 nm; m/e 341 (M+1)
755	5-methoxy	4,4,4-trifluorobutyl	LC Method 4: Rf 2.931 min at 254/220 nm; m/e 406 (M+1)
756	5-methoxy	2,2,3,3-tetrafluoropropyl	LC Method 4: Rf 2.795 min at 254/220 nm; m/e 411 (M+1)
757	5-chloro	2-fluoroethyl	LC Method 4: Rf 2.477 min at 254/220 nm; m/e 347 (M+1)

-148-

The following compounds were prepared following a procedure following Example 673:



No.	Z'	R ₄	Data
758	3-trifluoromethyl	2,2,3,3-tetrafluoropropyl	LC Method 4: R _f 2.650 min at 254/220 nm; m/e 410 (M+1)
759	3-trifluoromethyl	4,4,4-trifluorobutyl	LC Method 4: R _f 2.761 min at 254/220 nm; m/e 406 (M+1)

5

Example 760


3-Trifluoromethoxyphenethylamine

Combine nitromethane (1.8 g, 30 mmol), ethanol (4 mL) and 10 N NaOH (0.1 mL). Add 3-trifluoromethoxybenzaldehyde (5.0 g, 28.6 mmol) and stir. After 20 hours, pour into ethyl acetate, wash with water, dried over Na₂SO₄, filter, and concentrate to give a residue. Chromatograph the residue on silica gel to give 2-nitro-1-(3-trifluoroethoxyphenyl)ethanol: MS (M-1) 250; ¹H NMR (CDCl₃) 7.45 (1 H, t, J = 8.4 Hz), 7.36-7.30 (2 h, m), 7.24-7.20 (1 h, m), 5.51 (1 h, dt, J = 8.8 and 4.0 Hz), 4.62-4.51 (2 H, m).

Combine 2-nitro-1-(3-trifluoroethoxyphenyl)ethanol (6.1 g, 24.2 mmol) and methanesulfonyl chloride (2.02 mL) in dichloromethane (50 mL) and cool in an ice-bath. Add dropwise, triethylamine (7.28 mL) while maintaining the temperature near 0 °C. After 2 hours, pour into ethyl acetate, wash with water, dry with Na₂SO₄, filter, and then concentrate to residue. Chromatograph the residue on silica gel to give 3-(2-nitrovinyl)-1-trifluoroethoxybenzene: MS (MH⁺) 234; ¹H NMR (CDCl₃) 7.97 (1 H, d, J = 13.6 Hz), 7.57 (1 H, d, J = 13.6 Hz), 7.53-7.48 (2 H, m), 7.40-7.35 (2 H, m).


Combine 3-(2-nitrovinyl)-1-trifluoroethoxybenzene (3.0 g, 12.88 mmol) and methanol (50 mL) and concentrated HCl (5 mL) and hydrogenate at ambient temperature and 50 psi (340 kPa) in the presence of PtO₂ (0.6 g). After 5 hours, filter to the catalyst, dilute the filtrate with 1N HCl (50 mL) and wash with ethyl acetate. Separate the aqueous layer, neutralize with 2N NaOH (100 mL), extract with ether, dry with Na₂SO₄, filter and then concentrated to give the title compound which can be used without further purification. MS (MH⁺) 206; ¹H NMR (CDCl₃) 7.32 (1 H, t, J = 7.6 Hz), 7.18-7.06 (3 H, m), 2.98 (2 H, t, J = 7.2 Hz), 2.77 (2 H, t, J = 7.2 Hz).

-149-

Example 761N-(2-(3-Trifluoromethoxyphenyl)ethyl)-3-(2,2,2-trifluoroethyl)benzylamine


- 5 Combine trifluoromethoxyphenethylamine (400 mg, 1.95 mmol), and 3-(2-trifluoroethoxy)benzaldehyde (596 mg, 2.92 mmol), and 4Å molecular sieve (4.0 g) in ethanol (30 mL) and reflux. after 4.5 hours, decant and treat of NaBH₄ (221 mg, 5.85 mmol). After 1 hour, evaporate and partition between 5N NaOH and dichloromethane. Separate the organic layer, dry over Na₂SO₄, filter, and concentrate to give a residue.
- 10 Chromatograph the residue by HPLC to give the title compound. The HCl salt of the title compound gives a white solid: MS (MH⁺) 394; ¹H (DMSO-d₆) 9.48 (2 H, br s), 7.48 (1 H, t, J = 7.6 Hz), 7.40 (1 H, t, J = 8.0 Hz), 7.34 (1 H, s), 7.32-7.21 (4 H, m), 7.11 (1 H, dd, J = 8.4 and 2.8 Hz), 4.79 (2 H, q, J = 8.8 Hz), 4.15 (2 H, s), 3.22-3.12 (2 H, m), 3.11-3.04 (2 H, m).

15

Example 762N-(2-(3-Trifluoromethoxyphenyl)ethyl)-3-(2,2,3,3-tetrafluoropropyl)benzylamine


- 20 The method of Example 761 gives the title compound. The HCl salt of the title compound gives a white solid: MS (MH⁺) 426; ¹H (DMSO-d₆) 9.42 (2 H, br s), 7.48 (1 H, t, J = 7.6 Hz), 7.40 (1 H, t, J = 7.6 Hz), 7.32-7.26 (3 H, m), 7.20 (1 H, d, J = 7.2 Hz), 7.11 (1 H, dd, J = 8.4 and 2.8 Hz), 6.70 (1 H, tt, J = 5.2 and 5.2 Hz), 4.62 (2 H, t, J = 13.6 Hz), 4.15 (2 H, s), 3.22-3.12 (2 H, m), 3.10-3.02 (2 H, M).

-150-

Example 763N-(2-(4,7-Difluoro-1H-indol-3-yl)ethyl)-3-(2,2,2-trifluoroethoxy)benzylamine

- Combine 2-(4,7-difluoro-1H-indol-3-yl)ethylamine (483 mg, 2.46 mmol) and ethanol (45mL) and stir. After 10 minutes treat with 3-(2,2,2-trifluoroethoxy)benzaldehyde (502 mg, 2.46 mmol) and anhydrous sodium sulfate (3.5 g) and stir under nitrogen and heat at 70° C. After 2h., cool the reaction vacuum filter to remove the sodium sulfate and treat with sodium borohydride (279 mg, 7.38 mmol) in a 500 mL round bottom flask equipped with magnetic stirring. Allow the solution to stir for 2 hours at room temperature and then carefully treat with three drops of glacial acetic acid to quench the excess hydride. Remove the solvent *in vacuo* and re-dissolve the crude material in methanol. Purify by a 10 g SCX column by washing thoroughly with methanol, eluting with 2N ammonia in methanol, and concentrating *in vacuo* to give the title compound as a straw colored oil. Prepare the hydrochloride salt by dissolving the free base (800 mg, 2.08 mmol) in methanol (15 mL) and treating with a solution of ammonium chloride (111 mg, 2.08 mmol) in methanol (5 mL). Sonicate the mixture for 10 minutes before concentrating *in vacuo* to give a white solid. Recrystallize from ethyl acetate to obtain the hydrochloride salt of the title compound: mp 208.5-210.0 °C; ¹H NMR (400 MHz, dms_o-d₆): 11.79 (br s, 1H), 9.21 (br s, 2H), 7.39 (t, 1H, *J* = 7.8 Hz), 7.32 (d, 1H, *J* = 2.0 Hz), 7.30 (s, 1H), 7.18 (d, 1H, *J* = 8.0 Hz), 7.11 (dd, 1H, *J* = 2.6, 8.2 Hz), 6.85-6.91 (m, 1H), 6.67-6.73 (m, 1H), 4.77 (q, 2H, *J* = 8.8 Hz), 4.16 (s, 4H), 3.12-3.16 (m, 4H); MS (APCI): *m/e* 385.1 (*M*+1); CHN (for C₁₉H₁₇F₅N₂O•HCl) calcd: C 54.23, H 4.31, N 6.66; found: C 54.20, H 4.30, N 6.66.

25

Example 764N-(2-(4,5,6,7-Tetrafluoro-1H-indol-3-yl)ethyl)-3-(2,2,2-trifluoroethoxy)benzylamine

- By a method similar to Example 763, using 2-(4,5,6,7-tetrafluoro-1H-indol-3-yl)ethylamine (484 mg, 2.08 mmol), ethanol (45 mL), 3-(2,2,2-trifluoroethoxy)benzaldehyde (425 mg, 2.08 mmol), anhydrous sodium sulfate (3.5 g)

30

-151-

sodium borohydride (236 mg, 6.24 mmol) to give the free base of the title compound as a straw colored solid. Recrystallize from methylene chloride to obtain the title compound: mp 107.2-108.2 °C. ¹H NMR (400 MHz, dms_o-d₆): 11.92 (br s, 1H), 7.32 (s, 2H), 6.95-6.99 (m, 2H), 6.87 (dd, 1H, *J* = 2.4, 8.0 Hz) 4.68 (q, 2H, *J* = 8.8 Hz), 3.70 (s, 2H), 2.88 (t, 2H, *J* = 7.2 Hz) 2.75 (t, 2H, *J* = 7.2 Hz). MS (ES⁺): *m/e* 421.1 (*M*+1). CHN (for C₁₉H₁₅F₇N₂O•1HCl•0.20 H₂O) calcd: C 53.83, H 3.66, N 6.61; found: C 53.75, H 3.33, N 6.

Example 765

10 5-Trifluoromethyltryptamine

Combine 4-trifluoromethylaniline (32.2 g, 199.8 mmol) and dichloromethane (600 ml) in a 2L round bottomed flask under nitrogen and cool to -70°C. Add tert-butylhypochlorite (protected from the light) (22.8 g, 210 mmol) in dichloromethane (150 ml) stir for a total of 45 min at about -65 to -70°C. At 35 min, add a solution of methylthioacetaldehyde dimethylacetal (30 g, 220.2 mmol) in dichloromethane (150 ml) is added. At 45 min at -70°C, add a solution of triethylamine (31.2 ml, 22.78 g, 225.1 mmol) in dichloromethane (80 ml). Bring the reaction mixture to room temperature. Wash with water and evaporate to dryness to give 72 g an oil.

Dissolve the oil in toluene (600 ml) and add triethylamine (60 ml). Heat to reflux. After 24 hours, evaporate the solvent and dry the residue under vacuum to yield 7a residue. Combine the residue, diethyl ether (600 ml), and 2N HCl (500 ml) and stir 24 hours at room temperature. Separate the aqueous layer and wash the organic layer successively with water and saturated NaHCO₃, dry over MgSO₄, filter, and evaporate to give a residue. Chromatograph on silica gel by eluting with cyclohexane-ethyl acetate (8/2, v/v), and pool the fractions containing the expected and evaporate to give 33.8g of 2-methylthio-5-trifluoromethyl-1H-indole.

Combine moist Raney nickel (330 g), 2-methylthio-5-trifluoromethyl-1H-indole (33.8 g, 146.2 mmol) and absolute ethanol (850 ml) and stir. After 1.5 hours, filter the mixture through celite and wash the celite with ethanol (500 ml). Evaporate the filtrate to dryness, add toluene (20 ml) and evaporate and dry to give 5-trifluoromethylindole: mp=55—60°C

Dissolve 5-trifluoromethylindole (24 g, 130 mmol) in anhydrous diethyl ether (288 ml) and cool to 10°C and add dropwise oxalyl chloride (12 ml) over 10 min. (exothermic reaction) and stir at room temperature for 4 h. Add and additional amount of oxalyl chloride (3 ml) and stir overnight at room temperature to give a solid. Collect the solid, wash with anhydrous diethyl ether (20 ml), and dry to give 2-(5-(trifluoromethyl-1H-indol-3-yl)-2-oxo-acetyl chloride.

-152-

Combine 2-(5-(trifluoromethyl)-1H-indol-3-yl)-2-oxo-acetyl chloride and NH_4OH 1N (700 ml) and stir the suspension intensely. After 3 hours, collect the 2-(5-(trifluoromethyl)-1H-indol-3-yl)-2-oxo-acetamide.

- 5 Add LiAlH_4 (37.95 g, 1.00 mol) to THF (650 mL) under ice-bath cooling. Prepare a solution of AlCl_3 (50 g, 375 mmol) in THF (600 ml) and add dropwise to the LiAlH_4 solution over 45 min at 5-10°C. While maintaining the temperature at about 5°C, add a solution of 2-(5-(trifluoromethyl)-1H-indol-3-yl)-2-oxo-acetamide (21.4 g, 83.5 mmol) in THF (600 ml) and stir overnight with warming to ambient temperature. Cool the mixture with ice water and treat with 30% NaOH (100 ml) while maintaining the temperature at less than about 30°C. After stirring for about 30 minutes, filter, wash with THF (2 L), and evaporate the filtrate to give the title compound. Form of the HCl salt by dissolving the title compound in diethyl ether and adding of a solution of HCl in diethyl ether (until acidic). Collect the solid by filtration, wash with diethyl ether, and dry under reduced pressure to give the hydrochloride salt of the title compound.
- 10
- 15 The title compound can be further purified by basic extraction of the hydrochloride salt into ethyl acetate, drying over MgSO_4 , filter, and evaporate to dryness followed by hydrochloride salt formation in diethyl ether.

Example 766

20 3-Propoxybenzaldehyde

- Combine 3-hydroxybenzaldehyde (790 g), K_2CO_3 , (1627 g) and DMF (8 L). Add 1-iodopropane (1000 g) and heat to 105°C and stir for 4h. Cool to about 50°C and add water (15 L), continue cooling to about room temperature and add toluene (10 L). Separate the organic layer and extract the aqueous phase with toluene (2 x 10 L), combine organic phases and wash with NaOH 1N (2 x 5.8 L), concentrate the combined organic layers *in vacuo* to afford the title compound.
- 25

Example 777

N-(2-(5-Methoxy-1H-indol-3-yl)ethyl)-3-propoxybenzylamine

- 30 Combine 3-propoxybenzaldehyde (14.05 g, 0.0856 mole) and 5-methoxytryptamine (13.64 g, 0.0717 mole) in 390 mL absolute EtOH. Add molecular sieves (19.2 g) and heat the suspension to reflux. After 4 hours, cool to room temperature and add NaBH_4 (37.32 g, 0.2146 mole) in 3 portions. Stir the mixture for 1 hour at room temperature, filter, evaporate the filtrate to a mass of about 100g, add water and dichloromethane. After separation, wash the aqueous phase with dichloromethane, combine the organic layers, dry over MgSO_4 , filter, evaporate the solvent *in vacuo* to afford the title compound.
- 35

-153-

Combine the title compound and isopropanol (250 mL) and slowly add a solution of HCl in EtOH (33 ml, 2.5N). Heat to reflux and stir for 30 min. Cool to room temperature and stir for 2h to give a solid. collect the solid by filtration, wash isopropanol, and dry to give the hydrochloride salt of the title compound.

5

Example 7782,2,3,3,3-Pentafluoropropyl tosylate

Combine 2,2,3,3,3-pentafluoropropan-1-ol (9.7 ml) and pyridine. Cool to between 0°C and 10 °C and add portion-wise the p-toluenesulfonylchloride (6.2g) and stir with warming to room temperature. After 3 hours at room temperature, pour the reaction mixture into ice water and stir for 30 min to give a solid. Filter the solid, wash with water, and dry to give of the title compound.

10

Example 7793,3,3-Trifluoro-propyl tosylate

Add 3,3,3-trifluoropropan-1-ol (61.8 ml) and pyridine (224 ml). Cool to between 0°C and 10°C and add portion-wise p-toluenesulfonylchloride (147 g). Allow to warm to ambient temperature and stir overnight. Add HCl 0.5N (1.6L), extract with ethyl acetate, combine the organic layers, dry over MgSO₄, filter, and evaporate to give the title compound.

20

Example 7806-Fluorotryptamine

Add dropwise 422 mL of glacial acetic acid to 40% aqueous dimethylamine (408 mL) over 40 minutes while maintaining the temperature below about 15°C. Cool to 0°C. After stirring for 20 minutes at 0°C, slowly add 37% aqueous formaldehyde (289 mL, 1.3 eq.) over about 15 minutes. Add 6-fluoroindole (400 g, 2.96 mol, 1eq.) in four portions over about 15 minutes. After 30 minutes, divide the reaction mixture into two portions. To one portion, slowly 1149g (75% of total mass) over 30 minutes to 3 L of 10% NaOH and stir at room temperature. After 18 hours, collect the solid that forms, wash three times with 200 mL of water, dry by suction to give wet 3-(N,N-dimethylaminomethyl)-6-fluoroindole.

25

30

Dilute another portion of the reaction mixture (383g, 25% of total mass) with aqueous NaOH till pH 12-13 to give a solid. After 30 minutes, collect the solid by filtration, wash with water, dry at 50°C overnight to give 3-(N,N-dimethylaminomethyl)-6-fluoroindole.

35

Combine KCN (50.8 g, 0.78 mol), 3-(N,N-dimethylaminomethyl)-6-fluoroindole (100 g, 0.52 mol), DMF (400 mL) and water (200 mL). Heat to reflux. Evolution of gas


-154-

begins at about 70°C. Maintain the reflux for 4 hours. Cool the reaction mixture to room temperature, dilute with water and toluene and stir for 10 minutes. Decant the organic layer and wash successively with of saturated aqueous sodium bicarbonate and of 2M aqueous hydrochloric acid. Concentrate to dryness the organic layer to give 2-(6-fluoro-1H-indol-3-yl)acetonitrile.

Combine 2-(6-fluoro-1H-indol-3-yl)acetonitrile (165 g, 0.925 mol) and THF (1.32 L). Slowly add 1M solution of BH₃ (2.042 L, 1,832 Kg, 0.131 mol) in THF over about 40 minutes. When the addition is complete, heat to reflux within 1 hour. After 1 hour at reflux, cool to room temperature and the reaction mixture, over about 25 minutes, to a well-stirred 15% aqueous solution of NaOH (1.9 L, 9.5 mol). After addition, slowly and gradually heat to 50°C. After 1 hour, heat 60°C. After 30 minutes, heat to reflux for 1 hour. Cool to room temperature and stir overnight, decant the alkaline aqueous layer and replace by water. Heat to 30°C under a pressure of 200 mbars in order to distill the THF until about 2.5 kg of distillate is removed. Extract the mixture with dichloromethane. Slowly add to the combined organic layers over a 25 minutes a mixture of 37% aqueous HCl (143 g) and water (220g) and stir to give a solid. After 1 hour, collect the solid by filtration, wash with of dichloromethane, and dry overnight to afford of the hydrochloride salt of the title compound.

Combine 6-fluorotryptamine hydrochloride (100 g, 0.437 mol), 2% w/w NaOH (2.5 kg), and dichloromethane (1.5 L) and stir. After 15 minutes, decant the organic layer, extract the aqueous layer with dichloromethane, combine organic layers, and concentrate to give a residue. Combine the residue and isopropanol and evaporate *in vacuo* to give the title compound.

25

Example 782N-(2-(6-Fluoro-1H-indol-3-yl)ethyl)-3-(2,2,3,3-tetrafluoropropyl)benzylamine

Combine isopropanol (500 g), 2,2,3,3-tetrafluoropropylbenzaldehyde (116.8 g), and 6-fluorotryptamine (1.15 equiv.). Heat to reflux over about 1.5 hour. After 30 minutes at the reflux, distill and over 30 minutes collect about 380g of distillate. Cool the reaction mixture to 50°C and add NaBH₄ (19.71 g) in one portion. After 1 hour at 50°C, slowly add water over 15 minutes and allow the resulting solution to cool to room temperature overnight. Distill the isopropanol under reduced pressure to give a residue and extract with dichloromethane, combine organic layers, and treat with 1N aqueous HCl (650 mL)

3-(2,2,2-Trifluoroethoxy)benzaldehyde

Combine 3-hydroxybenzaldehyde (134.3 g), potassium carbonate (304.0 g), 2,2,2-trifluoroethyl *p*-toluenesulfonate (293.6 g) and dimethylformamide (2 L). Heat the mixture at 90°C. After 15 hours, cool to room temperature, pour on ice-water, and extract with dichloromethane. Combine the organic layers, wash with 1N sodium hydroxide, and then with water. Dry the organic phase over magnesium sulfate, filter, and concentrate to a residue. Dissolve residue in toluene (200 ml), Chromatograph on silica gel eluting sequentially with toluene and then ethyl acetate to give a residue. Distill the residue under reduced pressure using a Claisen flask equipped with a Vigreux column gives title compound: bp 0.8 mm Hg, 84-85°C. Redistilling some fractions using a Claisen flask equipped with a Vigreux column and subsequently an adiabatic column filled with Rasching rings gives title compound: bp 0.9-1.0 mm Hg, 74-76°C.

Combine 3-(2,2,3,3-tetrafluoropropoxy)tosylate (200 g, 0.664 mol), 3-hydroxybenzaldehyde (101.7 g, 0.833 mol), dimethylformamide (1.5 L) and powdered potassium carbonate (192 g). Heat under stirring at 92°C for about 22 h. Cool the reaction mixture to 40°C, pour over ice-water and extract with ethyl acetate. Combine organic phases, wash with 1 N sodium hydroxide (1 L and 0.5 L) and then a solution of saturated sodium hydrogen carbonate, dry over magnesium sulfate, filter, evaporate to dryness to provide a oily residue. Distill the oily residue under reduced pressure in a Claisen flask to give a first fraction of the title compound: bp 108-110°C under 0.4-0.5 mmHg and a second fraction at 110-111°C under 0.4-0.5 mmHg.

Combine 3-(2,2,3,3-tetrafluoropropoxy)tosylate (5.72 g, 17.2 mmol), 3-hydroxybenzaldehyde (2.44 g, 20.0 mmol), dimethylformamide (36 ml) and powdered potassium carbonate (3.03 g) and heat at 110°C for 10 h. Cool to 20°C. Pass through a bed of aluminium-oxide-90 (57.2 g, 70-230 mesh, grade II-III, Brockmann: Merck # 1.01097) and elute with toluene (120 ml). Wash the eluted organic phase with 1N HCl

-156-

(36 ml) and then water. Evaporate the organic layer under reduced pressure to give the title compound.

Example 785

5 2-(5-Chloro-1*H*-Indol-3-yl)-2-oxo-acetyl chloride

Combine 5-chloroindole (20 g, 0.13 mole) and dibutyl ether (230 mL) and cool to 5°C and slowly add the oxalyl chloride (20.08 g, 0.16 mole) over 15 min while maintaining the temperature between 5°C and 10°C. Warm to room temperature and stir for 1 hour to give a solid. Cool to 5°C and stir for 15 minutes, collect the solid by
10 filtration, wash with dibutyl ether, and dry under vacuum to give the title compound.

Example 786

2-(5-Chloro-1*H*-Indol-3-yl)-2-oxoacetamide

Combine 2-(5-chloro-1*H*-indol-3-yl)-oxo-acetyl chloride (28.9 g, 0.12 mole) and
15 NH₄OH 1N solution (720 ml) to give a suspension. After 18 hours, filter, wash with water, and dry under vacuum to give the title compound.


Example 787

5-Chlorotryptamine

20 Cool to 5°C, a suspension of LiAlH₄ (40.97 g) in THF (700 ml). Add a solution of AlCl₃ (53.9 g, 0.40 mole) to THF (645 ml) over about 30 minutes while maintaining the temperature at about 5°C and 10°C. Add 2-(5-chloro-1*H*-Indol-3-yl)-2-oxo-acetamide (20 g, 0.09 mole) in THF (900 ml) while maintaining the temperature at between 5°C and 7.5°C. When the addition is complete warm to room temperature. Stir overnight and
25 then cool to 7°C and slowly add a solution of NaOH 50% (342 g, 4.28 mol). After stirring for about 1 hour, add anhydrous Na₂SO₄ (30 g) and filter the suspension on a celite bed. Evaporate the filtrate to dryness to give an oil. Combine the and Et₂O (500 mL) and add a solution of Et₂O/HCl 4.5N (15 mL) at room temperature to give a solid. Stir the suspension at room temperature for 1 hour, filter, and wash with 50 mL Et₂O, dry under
30 vacuum at 50°C to give the hydrochloride of the title compound.

Add 5-chlorotryptamine hydrochloride (15 g, 0.06 mole) water (150 ml), NaOH 1N (75 ml), and dichloromethane (350 mL). Stir the mixture at room temperature for 30 minutes, and separate the phases. Wash the aqueous phase with dichloromethane, combine the organic phases, dry over MgSO₄, filter, and evaporate to dryness under
35 vacuum to give the title compound.

-157-

Example 789N-(5-Chloro-1H-indol-3-yl)ethyl)-3-(2,2,3,3-tetrafluoro-propoxy)benzylamine

Combine 5-chlorotryptamine (12.1 g, 0.0621 mol) and 3-(2,2,3,3-
 5 tetrafluoropropoxybenzaldehyde (17.6 g, 0.0621 mole) in EtOH (340 mL). Add molecular
 sieves and heat to reflux and stir for 4h. Cool the mixture to room temperature and add
 NaBH₄ (7g, 0.1876 mol) in 3 portions. Stir the for 1h at room temperature. Filter the solid
 and evaporate the filtrate to a weight of about 90 g, add water, and extract with
 dichloromethane. Dry the combined organic layers over MgSO₄, filter, and remove the
 10 solvent under reduced pressure to afford the title compound.

Combine the title compound (27.6 g) and isopropanol (300 mL). Add a solution
 of oxalic acid (6 g) in isopropanol (60 mL) to give a suspension. Heat the suspension to
 reflux and stir for 30 min and then cool to room temperature. Stir for 1 hour at room
 temperature. collect the solid by filtration, wash with isopropanol, and dry under vacuum
 15 to afford the oxalate of the title compound.


Example 790N-2-(5-Chloro-1H-indol-3-yl)ethyl)-3-(2,2,3,3-tetrafluoro-propoxy)benzylamine L-
tartaric salt

Combine N-2-(5-chloro-1H-indol-3-yl)ethyl)-3-(2,2,3,3-
 20 tetrafluoropropoxy)benzylamine oxalic acid salt and dichloromethane (700 mL) and add
 NaOH 1N(150 mL), water (450 mL) and MeOH (190 mL). Stir the mixture for 1 h at
 room temperature. Separate the layers. Add water (200 mL) to the aqueous phase and
 extract with dichloromethane, combine the organic layers, dry over MgSO₄, filter, and
 25 evaporate under vacuum to afford 19.4g of N-2-(5-chloro-1H-indol-3-yl)ethyl)-3-(2,2,3,3-
 tetrafluoropropoxy)benzylamine.

Combine N-2-(5-chloro-1H-indol-3-yl)ethyl)-3-(2,2,3,3-
 tetrafluoropropoxy)benzylamine (19.4g) in isopropanol (125 mL) and warm to dissolve.
 Add a solution of L-tartaric acid (7.02 g) in isopropanol (70 mL). Add seeding crystals
 30 and stir to give a solid. After 2.5 hours, collect the solid by filtration, wash with
 isopropanol, and dry under vacuum at 45°C to afford the title compound.

-158-

By the method of Example 221 the following compounds were prepared, isolated as the maleate except where noted:



No.	Z'	R ₄	Data
791	6-bromo	2,2,3,3-tetrafluoro propyl	mp 162-164 °C. Analysis for C ₂₄ H ₂₃ BrF ₄ N ₂ O ₅ : Calcd: C, 50.10; H, 4.03; N, 4.87; found: C, 50.24; H, 4.02; N, 4.87.
792	6-bromo	2,2,2-trifluoro ethyl	mp 168-171 °C. Analysis for C ₂₃ H ₂₂ BrF ₃ N ₂ O ₅ : Calcd: C, 50.84; H, 4.08; N, 5.16; found: C, 51.02; H, 4.13; N, 5.21.
793	6- methane sulfonyl	2,2,3,3-tetrafluoro propyl	mp 233-235 °C. MS (ACPI): m/e 459.1 (M+1). Analysis for C ₂₁ H ₂₃ ClF ₄ N ₂ O ₃ S: Calcd: C, 50.96; H, 4.68; N, 5.66; found: C, 50.87; H, 4.65; N, 5.64. (isolated as the hydrochloride)
794	6- methane sulfonyl	2,2,2-trifluoro ethyl	mp 234-236 °C. MS (ACPI): m/e 427.0 (M+1). Analysis for C ₂₀ H ₂₂ ClF ₃ N ₂ O ₃ S: Calcd: C, 51.89; H, 4.79; N, 6.05; found: C, 51.84; H, 4.79; N, 6.10. (isolated as the hydrochloride)
795	6-benzene sulfonyl	2,2,3,3-tetrafluoro propyl	mp 213-215 °C. MS (ACPI): m/e 521.0 (M+1). Analysis for C ₂₆ H ₂₅ ClF ₄ N ₂ O ₃ S: Calcd: C, 56.07; H, 4.52; N, 5.03; found: C, 55.81; H, 4.66; N, 4.96. (isolated as the hydrochloride)
796	6-benzene sulfonyl	2,2,2-trifluoro ethyl	mp 231-233.5 °C. MS (ACPI): m/e 489.0 (M+1). Analysis for C ₂₅ H ₂₄ ClF ₃ N ₂ O ₃ S: Calcd: C, 57.20; H, 4.61; N, 5.34; found: C, 56.98; H, 4.63; N, 5.21. (isolated as the hydrochloride)

5

Example 7996-Methanesulfonyl-1H-indole

Dissolve 6-Methanesulfonyl-indol-1-ol (5.0 g, 23.7 mmol) in triethyl phosphite (35 ml) and heat at 160° for 5 hours. Cool the solution to ambient temperature and dilute with diethyl ether. Wash the ether solution with brine and water followed by drying (sodium sulfate) and reducing to residue. Crystallize the residue from warm ethyl acetate to give the title compound as colorless cubes: mp 149-152 °C. MS (ACPI): m/e 196.0 (M+1). Analysis for C₉H₉NO₂S: Calcd: C, 55.37; H, 4.65; N, 7.17; found: C, 55.14; H, 4.71; N, 7.20.

-159-


Example 8006-Benzenesulfonyl-1H-indole

Dissolve 6-Bromoindole (6.0 g, 30.6 mmol) in THF (100 ml) and cool the mixture to -10° . Slowly add 60% NaH in mineral oil (3.67 g). After 1 hour, add triisopropylsilyltrifluoromethane sulfonate (9.9 ml, 36.7 mmol) slowly, remove the cooling bath, and stir for 24 hours. Quench excess NaH with ice and remove the THF under vacuum. Dilute the remaining residue with water and extract with dichloromethane. Combine, wash (brine), dry (sodium sulfate), and reduce the extracts to residue. Purify the residue on silica gel using 60% hexanes/dichloromethane to give a yellow oil.

Cool a solution of 6-bromo-1-triisopropylsilyl-1H-indole (5.5 g, 15.7 mmol) in 100 mL anhydrous THF to -78°C under nitrogen and treat with 1.7 M t-butyl lithium (20.5 mL, 34.5 mmol) while keeping the temperature at -78°C . After the addition, slowly add phenylsulfonyl fluoride (2.1 ml, 17.3 mmol) and stir for 30 minutes at -78°C . Warm the mixture to ambient temperature and stir for 1 hour. Quench excess t-butyl lithium with ice and dilute the mixture with water followed by extraction with ethyl acetate. Combine, wash (brine), dry (sodium sulfate), and reduce the extracts to residue. Purify the residue on silica gel using 50% hexanes/dichloromethane to give the product as a white solid.

Dissolve the resulting white solid in THF (50 ml) and treat the solution with 1 M tetrabutylammonium fluoride (18.1 mL) and 1 M boric acid (18.1 ml). After stirring for 1.5 hours at ambient temperature, dilute the mixture with water and extract with ethyl acetate. Combine, wash (brine), dry (sodium sulfate), and reduce the extracts to residue. Purify the residue on silica gel using 1% methanol/dichloromethane to give the title compound as a white solid: mp $141-144^{\circ}\text{C}$. MS (ACPI): m/e 258.0 (M+1). Analysis for $\text{C}_{14}\text{H}_{11}\text{NO}_2\text{S}$: Calcd: C, 65.35; H, 4.31; N, 5.44; found: C, 64.99; H, 4.31; N, 5.39.

By the method of Example 440 the following compounds were prepared and isolated as the hydrochloride except where noted:




30

-160-

No.	Z'	R ₄	Data
802	7-Chloro	2,2,3,3-tetrafluoro propyl	ISMS 415 (M+1); ¹ H NMR (DMSO-d ₆ -HCl salt) 11.3 (bs, 1H), 9.4 (bs, 2H), 7.6-7.5 (m, 1H), 7.45-7.3 (m, 3H), 7.25-6.95 (m, 4H), 6.9-6.5 (m, 1H), 4.7-4.5 (m, 2H), 4.2 (bs, 2H), 3.25 (bs, 4H)
803	6-Methoxy	2,2,3,3-tetrafluoro propyl	¹ H NMR (CDCl ₃ -freebase) 7.99 (bs, 1H), 7.47-7.44 (d, 1H), 7.23-7.19 (m, 1H), 6.94-6.92 (d, 1H), 6.89-6.88 (m, 1H), 6.83-6.82 (m, 2H), 6.79-6.75 (m, 2H), 6.19-5.90 (m, 1H), 4.29-4.22 (m, 2H), 3.82 (s, 3H), 3.78 (m, 2H), 2.95 (s, 4H), no N-H observed

By the method of Example 270 the following compounds were prepared, isolated as the maleate except where noted:




5

809	5-(4-fluoro phenyl)	phenyl	ISMS 437 (M+1); C ₂₉ H ₂₆ FCIN ₂ O·0.2 H ₂ O: calcd: C, 73.08; H, 5.58; N, 5.88; found: C, 72.99; H, 5.38; N, 5.83
-----	---------------------	--------	--

Example 811

N-(2-(5-Methoxy-1H-indol-3-yl)-ethyl)-(3-(3,3,3-trifluoropropoxy)benzyl)amine




- 10 Combine 350 mg (1.8 mmol) 5-methoxytryptamine, 401 mg 3-trifluoropropoxybenzaldehyde (1.8 mmol) and 4g 4A molecular sieves in 35 mL EtOH and reflux overnight. Decant the liquid into a separate flask and treat with 209 mg (5.5 mmol) NaBH₄. Stir the reaction at ambient temperature for 1 hour. Concentrate under vacuum, and partition between 50 mL 1 N NaOH and 25 mL
- 15 dichloromethane. Extract the aqueous layer with 25 ml dichloromethane and combine the organic layers and concentrate to dryness. Purify the resulting oil by radial chromatography (SiO₂; 1% MeOH in CHCl₃ mixed with conc. NH₄OH) to afford 705 mg (1.8 mmol; 100%) of the desired compound as an oil. Conversion

-161-

to its HCl salt by stirring a solution of the compound in 50 mL 50/50 THF/EtOH with 1 g polyvinyl pyridine hydrochloride overnight, filtering and concentrating to a solid. Recrystallize the product from EtOAc: Analysis for $C_{21}H_{23}F_3N_2O_2 \cdot HCl$: calcd: C, 58.81; H, 5.64; N, 6.53; found: C, 58.42; H, 5.44; N, 6.51; ISMS 393

5 (M+1).


By the method of Example 811 the following compounds were prepared, isolated as the hydrochloride except where noted:



No.	Z'	R ₄	Data
812	5-fluoro	3,3,3-trifluoro propyl	Analysis for $C_{20}H_{20}F_4N_2O \cdot HCl$: calcd: C, 57.63; H, 5.08; N, 6.72; found: C, 57.49; H, 5.04; N, 6.76; ISMS 381 (M+1)
814	5-bromo	phenyl	Analysis for $C_{23}H_{21}BrN_2O \cdot HCl \cdot 0.5H_2O$: calcd: C, 59.18; H, 4.97; N, 6.00; found: C, 59.18; H, 4.80; N, 5.92 ISMS 422 (M+1)
815	5-bromo	2,2,3,3-tetrafluoro propyl	Analysis for $C_{20}H_{19}BrF_4N_2O \cdot HCl$: calcd: C, 48.46; H, 4.07; N, 5.65; found: C, 48.39; H, 3.95; N, 5.55; ISMS 459 (M+)
816	5-bromo	2,2,3,3,3-pentafluoro propyl	Analysis for $C_{20}H_{18}BrF_5N_2O \cdot HCl$: calcd: C, 46.76; H, 3.73; N, 5.45; found: C, 46.47; H, 3.67; N, 5.46; ISMS 478 (M+1)
817	5-SO ₂ CH ₃	phenyl	Analysis for $C_{20}H_{18}BrF_5N_2O \cdot HCl \cdot 0.5H_2O \cdot 0.4C_7H_8$: calcd: C, 64.01; H, 5.85; N, 5.57; found: C, 64.09; H, 5.64; N, 5.48 ISMS 421 (M+1)
818	5-cyano	phenyl	Analysis for $C_{24}H_{21}N_3O \cdot HCl \cdot 0.3H_2O$: calcd: C, 70.42; H, 5.57; N, 10.27; found: C, 70.55; H, 5.41; N, 10.25 ISMS 368 (M+1)
819	5-carboxylic acid methyl ester	phenyl	Analysis for $C_{25}H_{24}N_2O_3 \cdot HCl \cdot 0.3H_2O$: calcd: C, 68.04; H, 5.62; N, 6.35; found: C, 68.06; H, 5.64; N, 6.43 ISMS 401 (M+1)
820	5-carboxylic acid methyl ester	2,2,2-trifluoroethyl	Analysis for $C_{21}H_{21}F_3N_2O_3 \cdot HCl \cdot 0.1H_2O$: calcd: C, 56.72; H, 5.03; N, 6.30; found: C, 56.46; H, 4.77; N, 6.04 ISMS 407 (M+1)


-162-

821	5-carboxylic acid amide	phenyl	ISMS 385 (M ⁺); Analysis for C ₂₄ H ₂₃ N ₃ O ₂ •HCl•0.9H ₂ O•0.1C ₇ H ₈ : calcd: C, 66.32; H, 5.99; N, 9.39; found: C, 66.07; H, 5.68; N, 9.01; ¹ H NMR (Free base CDCl ₃) δ 8.56 (s, 1H), 8.13 (s, 1H), 7.64-7.62 (m, 1H), 7.33-7.22 (m, 4H), 7.10-6.94 (m, 6H), 6.87-6.84 (m, 1H), 6.2 (bs, 1H), 5.8 (bs, 1H), 3.77 (s, 2H), 2.99-2.94 (m, 4H), 1.7 (bs, 1H)
-----	-------------------------	--------	--

Example 825N-2-(5-Nitro-1H-indol-3-yl)-ethyl)-3-phenoxybenzylamine

- 5 Combine 5-nitrotryptamine(500 mg, 2.4 mmol), 3-phenoxybenzaldehyde (480 mg, 2.4 mmol) and 4g 4A molecular sieves in 30 mL EtOH and reflux overnight. Decant the liquid into a separate flask and treat with NaBH₄ (280 mg, 7.2 mmol) at ambient temperature. After 1 hour concentrate under vacuum and partition the residue between 25 mL 1 N NaOH and 25 mL dichloromethane.
- 10 Extract the aqueous layer with 25 ml dichloromethane and dry the combined organic layers over MgSO₄ and concentrate to dryness. Purify the resulting oil by radial chromatography (SiO₂; 2% MeOH in CHCl₃) to afford the desired compound as an oil. Convert to the HCl salt by treating a solution of the compound in 10 mL EtOH with 0.25 mL 5 N HCl and 40 mL toluene then concentrating to a solid.
- 15 Analysis for C₂₃H₂₁N₃O₃•HCl•0.2 EtOH: calcd: C, 64.62; H, 5.17; N, 9.75; found: C, 64.89; H, 5.40; N, 9.75; ISMS 388 (M+1).

By the method of Example 825 the following compounds were prepared, isolated as the hydrochloride except where noted:



No.	Z'	R ₄	Data
826	5-butoxy	phenyl	Analysis for C ₂₇ H ₃₀ N ₂ O ₂ •HCl•0.4H ₂ O: calcd: C, 70.77; H, 7.00; N, 6.11; found: C, 70.87; H, 6.84; N, 6.14; ISMS 415 (M+1)
827	5-benzamide	phenyl	Analysis for C ₃₀ H ₂₇ N ₃ O ₂ •HCl•0.2H ₂ O: calcd: C, 71.83; H, 5.71; N, 8.38; found: C, 71.63; H, 5.35; N, 8.09; ISMS 462 (M+1)
828	5-benzamide	2,2,2-trifluoro ethyl	Analysis for C ₂₆ H ₂₄ F ₃ N ₃ O ₂ •HCl: calcd: C, 61.97; H, 5.00; N, 8.33; found: C, 61.78; H, 5.16; N, 7.97; ISMS 468 (M+1)
829	5-benzamide	2,2,3,3- tetrafluoro propyl	Analysis for C ₂₇ H ₂₅ F ₄ N ₃ O ₂ •HCl: calcd: C, 60.51; H, 4.89; N, 7.84; found: C, 60.47; H, 4.95; N, 7.49; ISMS 500 (M+1)
830	5-methane sulfonamide	phenyl	Analysis for C ₂₄ H ₂₅ N ₃ O ₃ S • HCl• 0.5H ₂ O• 0.5 C ₇ H ₈ : calcd: C, 63.53; H, 5.86; N, 8.08; found: C, 63.57; H, 5.77; N, 7.81; ISMS 436 (M+1)
831	5-methane sulfonamide	2,2,2-trifluoro ethyl	Analysis for C ₂₀ H ₂₂ F ₃ N ₃ O ₃ S • HCl• 0.1H ₂ O• 0.5 C ₇ H ₈ : calcd: C, 53.68; H, 5.21; N, 7.99; found: C, 53.48; H, 5.19; N, 7.72; ISMS 442 (M+1)
832	5-methane sulfonamide	2,2,3,3- tetrafluoro propyl	Analysis for C ₂₁ H ₂₃ F ₄ N ₃ O ₃ S• HCl•0.1EtOH• 0.8C ₇ H ₈ : calcd: C, 54.72; H, 5.31; N, 7.14; found: C, 54.63; H, 5.25; N, 6.99; ISMS 474 (M+1)
833	5-isopropoxy	phenyl	Analysis for C ₂₆ H ₂₈ N ₂ O ₂ •1.1HCl • 0.1H ₂ O: calcd: C, 70.58; H, 6.68; N, 6.33; found: C, 70.37; H, 6.31; N, 6.35; ISMS 401 (M+1)
834	5-isopropoxy	2,2,2-trifluoro ethyl	Analysis for C ₂₂ H ₂₅ F ₃ N ₂ O ₂ •HCl• 0.3H ₂ O: calcd: C, 58.94; H, 5.98; N, 6.25; found: C, 59.08; H, 5.78; N, 6.25; ISMS 407 (M+1)
835	5-isopropoxy	2,2,3,3- tetrafluoro propyl	Analysis for C ₂₃ H ₂₆ F ₄ N ₂ O ₂ •HCl• 0.3H ₂ O: calcd: C, 57.51; H, 5.79; N, 5.83; found: C, 57.66; H, 5.55; N, 5.80; ISMS 439 (M+1)
836	5-ethoxy	phenyl	Analysis for C ₂₅ H ₂₆ N ₂ O ₂ •HCl• 0.2H ₂ O: calcd: C, 70.39; H, 6.47; N, 6.57; found: C, 70.40; H, 6.32; N, 6.68; ISMS 387 (M+1)

-164-

837	5-ethoxy	2,2,2-tri fluoroethyl	Analysis for $C_{21}H_{23}F_3N_2O_2 \cdot HCl$: calcd: C, 58.81; H, 5.64; N, 6.53; found: C, 58.61; H, 5.61; N, 6.52; ISMS 393 (M+1)
838	5-ethoxy	2,2,3,3-tetra fluoropropyl	Analysis for $C_{22}H_{24}F_4N_2O_2 \cdot HCl$: calcd: C, 57.33; H, 5.47; N, 6.08; found: C, 57.01; H, 5.35; N, 6.03; ISMS 425 (M+1)
839	2,2,2- trifluoro- ethoxy	phenyl	Analysis for $C_{25}H_{23}F_3N_2O_2 \cdot HCl$: calcd: C, 62.96; H, 5.07; N, 5.87; found: C, 62.76; H, 4.93; N, 5.88; ISMS 441 (M+1)
840	2,2,2- trifluoro- ethoxy	2,2,2-trifluoro ethyl	Analysis for $C_{21}H_{20}F_6N_2O_2 \cdot HCl$: calcd: C, 52.24; H, 4.38; N, 5.80; found: C, 52.21; H, 4.28; N, 6.18; ISMS 447 (M+1)
841	2,2,2- trifluoro- ethoxy	2,2,3,3-tetra fluoropropyl	Analysis for $C_{22}H_{21}F_7N_2O_2$ $\cdot HCl \cdot 0.2H_2O \cdot 0.2C_7H_8$: calcd: C, 52.35; H, 4.51; N, 5.22; found: C, 52.15; H, 4.30; N, 5.58; ISMS 479 (M+1)
842	5-butyloxy	pyridin-2-yl	Analysis for $C_{26}H_{29}N_3O_2 \cdot 2HCl$ $\cdot 0.5EtOH \cdot 0.3C_7H_8$: calcd: C, 64.83; H, 6.81; N, 7.79; found: C, 64.99; H, 6.48; N, 7.47; ISMS 416 (M+1)
843	5-isopropyl	2,2,2-tri fluoroethyl	Analysis for $C_{22}H_{25}F_3N_2O \cdot HCl$: calcd: C, 61.90; H, 6.14; N, 6.56; found: C, 61.72; H, 6.14; N, 6.42; ISMS 391 (M+1)
844	5-isopropyl	phenyl	Analysis for $C_{26}H_{28}N_2O \cdot HCl$: calcd: C, 74.18; H, 6.94; N, 6.65; found: C, 73.82; H, 6.79; N, 6.65; ISMS 385 (M+1)
845	5-benzene sulfonyl	phenyl	Analysis for $C_{29}H_{26}N_2O_3S \cdot 2HCl$: calcd: C, 67.11; H, 5.24; N, 5.40; found: C, 67.46; H, 5.37; N, 5.09; ISMS 483 (M+1)
846	5-benzene sulfonyl	2,2,3,3-tetra fluoropropyl	Analysis for $C_{26}H_{24}F_4N_2O_3S \cdot HCl$ $\cdot 0.3EtOH \cdot 0.2C_7H_8$: calcd: C, 57.07; H, 4.86; N, 4.75; found: C, 56.95; H, 4.68; N, 4.77; ISMS 521 (M+1)
847	5-benzene sulfonyl	2,2,2-tri fluoroethyl	Analysis for $C_{26}H_{29}N_3O_2$ $\cdot HCl \cdot 0.6H_2O$: calcd: C, 56.04; H, 4.74; N, 5.23; found: C, 56.05; H, 4.71; N, 5.12; ISMS 489 (M+1)

-165-

848	5-carboxylic acid ethyl ester	2,2,2-tri fluoroethyl	Analysis for $C_{22}H_{23}F_3N_2O_3 \cdot HCl$: calcd: C, 57.84; H, 5.30; N, 6.13; found: C, 57.85; H, 5.17; N, 6.09; ISMS 421 (M+1)
849	5-carboxylic acid propylamide	2,2,3,3-tetra fluoropropyl	Analysis for $C_{23}H_{26}F_3N_3O_2 \cdot HCl \cdot 0.6H_2O \cdot 0.1C_7H_8$: calcd: C, 56.84; H, 5.79; N, 8.05; found: C, 56.65; H, 5.63; N, 7.71; ISMS 466 (M+1)
850	5-carboxylic acid propylamide	phenyl	Analysis for $C_{27}H_{29}N_3O_2 \cdot HCl \cdot 0.4H_2O \cdot 0.2C_7H_8$: calcd: C, 69.66; H, 6.67; N, 8.58; found: C, 69.75; H, 6.57; N, 8.38; ISMS 428 (M+1)
851	5-carboxylic acid propylamide	2,2,2-tri fluoroethyl	Analysis for $C_{23}H_{26}F_3N_3O_2 \cdot HCl \cdot 0.8H_2O \cdot 0.1C_7H_8$: calcd: C, 57.67; H, 6.00; N, 8.51; found: C, 57.55; H, 5.77; N, 8.43; ISMS 434 (M+1)
852	5-carboxylic acid butylamide	phenyl	Analysis for $C_{28}H_{31}N_3O_2 \cdot HCl \cdot 0.7H_2O$: calcd: C, 68.54; H, 6.86; N, 8.56; found: C, 68.41; H, 6.60; N, 8.37; ISMS 442 (M+1)
853	5-carboxylic acid butylamide	2,2,3,3-tetra fluoropropyl	Analysis for $C_{25}H_{29}F_4N_3O_2 \cdot HCl \cdot H_2O$: calcd: C, 56.23; H, 6.04; N, 7.87; found: C, 56.23; H, 5.79; N, 7.84; ISMS 480 (M+1)
854	H	2,2,3,3-tetra fluoro propyl	Analysis for $C_{20}H_{20}F_4N_2O \cdot HCl \cdot 0.5H_2O$: calcd: C, 56.41; H, 5.21; N, 6.58; found: C, 56.98; H, 4.93; N, 6.53; ISMS 381 (M+1)
855	5-benzyloxy	2,2,2-tri fluoroethyl	Analysis for $C_{26}H_{25}F_3N_2O_2 \cdot HCl$: calcd: C, 63.61; H, 5.34; N, 5.71; found: C, 63.46; H, 5.53; N, 5.72; ISMS 455 (M+1)
856	5-benzyloxy	2,2,3,3 tetra fluoropropyl	Analysis for $C_{27}H_{26}F_4N_2O_2 \cdot HCl$: calcd: C, 62.01; H, 5.20; N, 5.36; found: C, 62.04; H, 5.16; N, 5.36; ISMS 487 (M+1)
857	6-phenoxy	phenoxy	ISMS 435 (M+1); $C_{29}H_{27}ClN_2O_2 \cdot 0.1 H_2O$: calcd: C, 73.67; H, 5.80; N, 5.93; found: C, 73.49; H, 5.49; N, 5.82
858	6-Phenoxy	2,2,3,3-tetra fluoropropyl	ISMS 473 (M+1); $C_{26}H_{25}F_4ClN_2O_2$: calcd: C, 61.36; H, 4.95; N, 5.50; found: C, 61.02; H, 4.67; N, 5.42
859	6-Phenoxy	2,2,2-tri fluoroethyl	ISMS 441 (M+1); $C_{26}H_{25}F_4ClN_2O_2 \cdot 0.2 H_2O$: calcd: C, 62.49; H, 5.12; N, 5.83; found: C, 62.27; H, 4.78; N, 5.74

-166-

860	5(3-pyridyloxy)	2,2,3,3-tetrafluoropropyl	ISMS 474 (M+1); C ₂₅ H ₂₅ F ₄ Cl ₂ N ₃ O ₂ ·0.5 H ₂ O: calcd: C, 54.06; H, 4.72; N, 7.57; found: C, 53.97; H, 4.76; N, 7.29
861	5-(Pyridinyl-3-oxy)	2,2,2-trifluoroethyl	ISMS 442 (M+1); ¹ H NMR (CDCl ₃) 8.37-8.36 (m, 1H), 8.27-8.26 (m, 1H), 8.01 (bs, 1H), 7.35-7.32 (m, 1H), 7.26-7.24 (m, 3H), 7.22-7.18 (m, 2H), 7.08-7.07 (m, 1H), 6.93-6.91 (m, 2H), 6.9-6.86 (m, 1H), 6.79-6.76 (m, 1H), 4.31-4.25 (m, 2H), 3.77 (s, 2H), 3.77 (s, 4H). Isolated as dihydrochloride salt

Example 8636-Phenoxytryptamine

By using a method similar to Example 422, the title compound was prepared:

- 5 ISMS 253 (M+1); ¹H NMR (CDCl₃) 8.1 (bs, 1H), 7.56-7.54 (m, 1H), 7.32-7.28 (m, 3H), 7.07-6.98 (m, 4H), 6.89-6.86 (m, 1H), 3.06-3.02 (m, 2H), 2.92-2.88 (m, 2H), 1.68 (bs, 2H).

Example 864

- 10 2-(5-(Pyridin-3-yloxy)-1H-indol-3-yl)-ethylamine

By using a method similar to Example 422, the title compound was prepared:

ISMS 254 (M+1); C₁₅H₁₅N₃O·1.1 C₂H₂O₄·0.2 H₂O: calcd: C, 58.04; H, 4.98; N, 11.81; found: C, 58.17; H, 4.62; N, 11.45.

15

Example 8656-Phenoxy-1H-indole-3-carbaldehyde

By using a method similar to Example 414, the title compound was prepared: ISMS 238 (M+1); ¹H NMR (CDCl₃) 10.78 (bs, 1H), 9.95 (s, 1H), 8.20-8.18 (m, 1H), 7.76-7.75 (m, 1H), 7.30-7.26 (m, 2H), 7.06-7.02 (m, 2H), 7.00-6.95 (m, 3H).

20

Example 8665-(Pyridin-3-yloxy)-1H-indole-3-carbaldehyde

By using a method similar to Example 414, the title compound was prepared: ISMS 239 (M+1); C₁₄H₁₀N₂O₂·0.3 H₂O: calcd: C, 69.01; H, 4.39; N, 11.50; found: C, 68.91; H, 4.16; N, 11.39.

25

-167-

Example 8673-(3-Methyl-4-nitrophenoxy)pyridine

Rinse 35% oil dispersion of KH (12g, 11 mmol) with 100 mL hexanes twice and dry under vacuum before cooling in an ice bath. Add 100 mL dry DMF then a
5 solution of 3-hydroxypyridine (10 g, 105 mmol) in 100 mL DMF dropwise. Treat with a solution of 5-fluoro-2-nitrotoluene (16.3 g, 105 mmol) in 50 mL DMF to obtain a dark solution. Stir at ambient temperature for 1 hour, pour the mixture into 1 liter of brine and extract twice with 200 mL of EtOAc. Combine the extracts and wash twice with 500 mL brine, dry over MgSO₄ and concentrate to 24 g of a dark
10 oil. Purification by chromatography 20% EtOAc in hexanes give the title compound as an oil: ISMS 231 (M+1); C₁₂H₁₀N₂O₃: calcd: C, 62.61; H, 4.38; N, 12.17; found: C, 62.63; H, 4.58; N, 12.06.

Example 8693-Ethoxybenzaldehyde

Combine 5.6 g of 3-hydroxybenzaldehyde (46 mmol) and 10.7 g of 1-iodoethane (69 mmol) in DMSO (25 mL) and warm to 80°C and treat with 22.4 g of cesium carbonate (69 mmol) portionwise and stir. After 1 hour, pour into 200 mL brine and extract twice with 150 mL diethyl ether. Combine the extracts and
20 wash twice with 200 mL brine, dry over MgSO₄ and concentrate under vacuum to give an oil. Purification by chromatography (SiO₂; 2.5% EtOAc in Hexanes) affords 5.73 g (38 mmol; 83%) of the desired compound as an oil: ¹H NMR (CDCl₃) 9.94 (s, 1H), 7.42-7.41 (m, 2H), 7.36-7.35 (m, 1H), 7.16-7.13 (m, 1H), 4.10-4.04 (q, 2H), 1.64-1.40 (t, 3H).

25

Example 8703-Propoxybenzaldehyde

By using a method similar to Example 869, the title compound was prepared: ¹H NMR (CDCl₃) 9.95 (s, 1H), 7.43-7.41 (m, 2H), 7.37-7.36 (m, 1H), 7.17-7.14 (m, 1H), 9.98-3.95
30 (t, 2H), 1.84-1.79 (m, 2H), 1.05-1.02 (t, 3H).

Example 8724-Phenoxy-1-methyl-2-nitrobenzene

Combine phenyl boronic acid (7.32 g, 60 mmol), 4-methyl-3-nitrophenol (4.5 g, 30
35 mmol), and Cu(oAC)₂·H₂O (6 g, 30 mmol) in 30 mL CH₂Cl₂ and treat with 6 g 4A molecular sieves powder. Add Et₃N (15.18 g, 150 mmol) dropwise, and stir the reaction at ambient temperature for 8 days. Dilute with 100 mL CH₂Cl₂ and filter through celite and

-168-

concentrate to dryness. Purification by chromatography using 2% EtOAc in hexanes gave the desired product as a yellow oil.

Example 873

5 6-Phenoxy-1H-indole

Combine 4-phenoxy-1-methyl-2-nitro-benzene (6 g, 26.2 mmol) and DMF dimethylacetal (15.6 g, 131 mmol) in 60 mL dry DMF and heat at 170 °C for 16 hours. Cool to room temperature and concentrate to dryness. Dissolve residue in 50 mL EtOAc and hydrogenate with 2 g 5% Pd/C and hydrogen for 3 hours at atmospheric pressure.

- 10 Filter through celite and concentrate to an oil. Purify by chromatography using Hex/EtOAc to obtain a tan solid: ISMS 210 (M+1)
¹H NMR (CDCl₃) 8.08 (bs, 1H), 7.61-7.59 (m, 1H), 7.34-7.29 (m, 2H), 7.18-7.17 (m, 1H), 7.18-7.0 (m, 4H), 6.92-6.89 (m, 1H), 6.56-6.54 (m, 1H).

15 Example 874

5-Pyridin-3-yl-1-methyl-2-nitro-benzene

By a method similar to Example 872, the title compound was prepared.


Example 875

20 5-(Pyridin-3-yloxy)-1H-indole

By a method similar to Example 873, the title compound was prepared: ISMS 211 (M+1); C₁₃H₁₀N₂O·0.1 H₂O: calcd: C, 73.64; H, 4.85; N, 13.21; found: C, 73.76; H, 4.80; N, 13.09.

25 Example 877

N-2-(5-Phenoxy-1H-indol-3-yl)-ethyl-3-phenoxybenzylamine



- Combine 2-(5-phenoxy-1H-indol-3-yl)ethylamine (0.400 g, 1.59 mmol), 3-phenoxybenzaldehyde (0.377 g, 1.90 mmol) and molecular sieves 4A (0.40 g) and stir in
30 methanol (15 mL). After 4h, filter the molecular sieves and wash several times with MeOH. To this MeOH solution, add portionwise NaBH₄ (61.5 mg, 1.59 mmol), stir the

-169-


- resulting mixture at room temperature for 1h. Remove MeOH under vacuum, dilute the residue with CH₂Cl₂ / water, extract with CH₂Cl₂, the combine the organic layers and dry over Na₂SO₄. Concentrate *in vacuo* the solvent, purification on silica gel (CH₂Cl₂ / MeOH) to give the free base of the title compound. React free base with oxalic acid to form the salt: m.p. 196-198 °C; ¹H NMR (300 MHz, DMSO-d₆) 2.95-3.15 (m, 4H), 4.15 (s, 2H), 6.85-7.46 (m, 18H), 11.06 (br, 1H); MS (ELECTROSPRAY) m/e: 435.3 (M+1); HRMS (ES+) calcd for C₂₉H₂₇N₂O₂ (M+H) 435.2084 found 435.2073.

Example 878

- (3-Phenoxybenzyl)-(2-(5-phenoxy-1H-indol-3-yl)-ethyl)-carbamic acid tert butyl ester
- Combine (3-phenoxy-benzyl)-(2-(5-phenoxy-1H-indol-3-yl)-ethyl)-amine (0.96 g, 2.2 mmol) and NaOH (87.7 mg, 2.2 mmol) and dissolve in THF (10 mL), stir at room temperature for 15 min. Add di-*tert*-butyl dicarbonate (0.58 g, 2.64 mmol) in THF (10 mL) and stir. After 2h, dilute the reaction with water, extract with EtOAc (3 x 15 mL), dry over Na₂SO₄. Concentrate the organic solvent on *vacuum* to give the title compound as an oil: ¹H NMR (300 MHz, CDCl₃) 1.36 (s, 9H), 2.85-2.91 (m, 2H), 3.89-3.65 (m, 2H), 4.26 (s, 1H), 4.39 (s, 1H) 6.83-7.13 (m, 10H), 7.21-7.33 (m, 7H), 8.00 (s, 1H); MS (ELECTROSPRAY) m/e 534.9 (M+1).

Example 879

- N-Methyl-N-2-(5-Phenoxy-1H-indol-3-yl)-ethyl)-3-phenoxybenzylamine




- Add slowly 1.0 M solution of LiAlH₄-THF (5.5 mL, 5.5 mmol) to a solution of (3-phenoxy-benzyl)-(2-(5-phenoxy-1H-indol-3-yl)-ethyl)-carbamic acid tert butyl ester (0.60g, 1.12 mmol) in 10 mL dry THF. After addition, heat to reflux the reaction mixture. After 2h, cool to room temperature, quench the reaction by adding water 1.5 mL cautiously, followed by 2N NaOH (1.0 mL). Filter the suspension and wash repeatedly with ether, dry the organic solution over Na₂SO₄ and concentrate *in vacuo*. Purification on silica gel using CH₂Cl₂ / MeOH as eluent gives the free base of the title compound and further reaction with oxalic acid to form the salt: m.p. 174-175 °C; ¹H NMR (250 MHz,

-170-

DMSO-d₆) 2.51(s, 3H), 3.00-3.13 (m, 4H), 4.15 (s, 2H), 6.81-7.03 (m, 7H), 7.11-7.42 (m, 11H), 11.05 (br, 1H); MS (ELECTROSPRAY) m/e: 449.1 (M+1-C₂H₂O₄).

Example 880


5 N-(2-(6-Chloro-1H-indol-3-yl)ethyl)-(3-(2,2-difluoroethoxy)benzyl)amine



Combine 2-(6-chloro-1H-indol-3-yl)ethylamine hydrochloride(1.0 g, 4.3 mmol) and ethyldiisopropylamine (900 μ L, 5.2 mmol) in ethanol (150 mL) and stir at room
 10 temperature and treat with 3-(2,2-difluoroethoxy)benzaldehyde (856 mg, 4.6 mmol) and anhydrous sodium sulfate (12 g) and heat at 78°C overnight. Cool to room temperature and filter. Treat the resulting filtrate with sodium borohydride (488 mg, 12.9 mmol) and stir the milky-white mixture at room temperature overnight. Remove *in vacuo* the solvent, and purification of the crude on silica gel eluting with 10% methanol in dichloromethane
 15 gives the free base of the title compound as a light yellow oil. Dissolve a portion of the oil (651 mg, 1.78 mmol) in methanol (15 mL) and treat with a homogenous solution of ammonium chloride (95 mg, 1.78 mmol) in methanol (3 mL). Sonicate the resulting solution for 10 minutes before removal of the solvent *in vacuo* to provide an off-white solid. Triturate with diethyl ether containing a few drops of acetonitrile. Filtration and
 20 drying of the precipitate afforded the title hydrochloride as a white solid: mp 131.6-133°C; ¹H NMR (400 MHz, dmso-d₆): 11.15 (br s, 1H), 9.50 (br s, 2H), 7.57 (d, 1H, *J* = 8.8 Hz), 7.39 (d, 1H, *J* = 2.0 Hz), 7.36 (t, 1H, *J* = 8.2 Hz), 7.32 (br s, 1H), 7.26 (d, 1H, *J* = 2.0 Hz), 7.17 (d, 1H, *J* = 7.6 Hz), 7.04 (dd, 1H, *J* = 7.8, 2.2 Hz), 7.01 (dd, 1H, *J* = 8.4, 2.0 Hz), 6.41 (tt, 1H, *J* = 54.4, 3.4 Hz), 4.32 (td, 2H, *J* = 14.8, 3.6 Hz), 4.14 (br s, 2H), 3.11
 25 (br s, 4H); MS (ES+): m/e 365.3 (M+1); CHN (for C₁₉H₁₉F₂ClN₂O•HCl•0.3 H₂O) calcd: C 56.11; H 5.11; N 6.89; found: C 56.03; H 4.95; N 7.18.

Example 881

N-Methyl-N-(2-(6-Chloro-1H-indol-3-yl)ethyl)-3-(2,2-difluoroethoxy)benzylamine




30

-171-

- Combine (2-(6-chloro-1*H*-indol-3-yl)ethyl)-(3-(2,2-difluoroethoxy)benzyl)amine (276 mg, 0.76 mmol) and formaldehyde (55.5 μ L of a 38% aqueous solution, 0.76 mmol) in dichloroethane (15 mL) and stir at room temperature for 10 minutes; add in two portions sodium triacetoxymethylborohydride (321 mg, 1.51 mmol) over 10 minutes and stir at room temperature overnight before diluting with methanol (10 mL) and quenching with one drop of glacial acetic acid. Remove *in vacuo* the solvent, and redissolve the crude residue in methanol and directly load onto a 10 g SCX column. After washing thoroughly with methanol, elute the column with 2 N ammonia in methanol. Concentrate *in vacuo* the eluant to give the free base of the title compound as a straw-colored oil. Dissolve the free base (239 mg, 0.64 mmol) in methanol (20 mL) and treat with a solution of ammonium chloride (36 mg, 0.67 mmol) in methanol (5 mL). Sonicate the mixture for 10 minutes before removal of the solvent *in vacuo* to give the hydrochloride salt as a tacky yellow oil. Dissolve the oil in 10 mL of 1:1 acetonitrile-water and lyophilize overnight, providing a fluffy white solid and triturate with diethyl ether (10 mL) and acetonitrile (2 drops). Filtration and drying of the resulting precipitate afforded the desired hydrochloride as a white amorphous solid: mp: 63.8-65.8°C; ^1H NMR (400 MHz, $\text{dms}\text{-d}_6$): 11.10 (br s, 1H), 7.52 (d, 1H, $J = 8.4$ Hz) 7.36 (d, 1H, $J = 2.0$ Hz), 7.40-7.26 (m, 2H), 7.22 (d, 1H, $J = 2.4$ Hz), 7.20-7.11 (m, 1H), 7.04 (br d, 1H, $J = 7.6$ Hz), 6.96 (dd, 1H, $J = 8.6, 1.4$ Hz), 6.38 (tt, 1H, $J = 54.4, 3.6$, Hz), 4.50-4.02 (br m, 2H), 4.30 (td, 2H, $J = 14.4, 3.2$ Hz), 3.15 (br s, 4H), 2.68 (br s, 3H); MS (ES $^+$): m/e 378.9 ($M+1$); CHN (for $\text{C}_{20}\text{H}_{21}\text{ClF}_2\text{N}_2\text{O}\cdot\text{HCl}\cdot 0.7\text{H}_2\text{O}$) calcd: C 56.14, H 5.51, N 6.55; found: C 55.72, H 5.32, N 7.07.

- By the method of Example 319 the following compounds were prepared, isolated as the oxalate except where noted:



No.	Z'	R ₄	Data
883	4,7-difluoro	2,2,2-trifluoro ethyl	mp 208.5-210.0 °C; ¹ H NMR (400 MHz, dmso-d ₆): 11.79 (br s, 1H), 9.21 (br s, 2H), 7.39 (t, 1H, <i>J</i> = 7.8 Hz), 7.32 (d, 1H, <i>J</i> = 2.0 Hz) 7.30 (s, 1H), 7.18 (d, 1H, <i>J</i> = 8.0 Hz), 7.11 (dd, 1H, <i>J</i> = 2.6, 8.2 Hz), 6.85-6.91 (m, 1H), 6.67-6.73 (m, 1H), 4.77 (q, 2H, <i>J</i> = 8.8 Hz), 4.16 (s, 4H), 3.12-3.16 (m, 4H). MS (APCI): <i>m/e</i> 385.1 (M+1). CHN (for C ₁₉ H ₁₇ F ₅ N ₂ O•1HCl) calcd: C 54.23, H 4.31, N 6.66; found: C 54.20, H 4.30, N 6.66.
884	4,5,6,7-tetrafluoro	2,2,2-trifluoro ethyl	mp 107.2-108.2 °C; ¹ H NMR (400 MHz, dmso-d ₆): 11.92 (br s, 1H), 7.32 (s, 2H), 6.95-6.99 (m, 2H), 6.87 (dd, 1H, <i>J</i> = 2.4, 8.0 Hz) 4.68 (q, 2H, <i>J</i> = 8.8 Hz), 3.70 (s, 2H), 2.88 (t, 2H, <i>J</i> = 7.2 Hz) 2.75 (t, 2H, <i>J</i> = 7.2 Hz). MS (ES+): <i>m/e</i> 421.1 (M+1). CHN (for C ₁₉ H ₁₅ F ₇ N ₂ O•1HCl•0.20 H ₂ O) calcd: C 53.83, H 3.66, N 6.61; found: C 53.75, H 3.33, N 6.54.
885	4,7-difluoro	2,2,3,3-tetrafluoro propyl	mp 171.8-173.0 °C; ¹ H NMR (400 MHz, dmso-d ₆): 11.80 (br s, 1H), 9.21 (s, 2H), 7.39 (t, 1H, <i>J</i> = 8.0 Hz), 7.30-7.33 (m, 2H), 7.18 (d, 1H, <i>J</i> = 7.6 Hz), 7.10 (dd, 1H, <i>J</i> = 2.4, 8.0 Hz), 6.85-6.91 (m, 1H), 6.54-6.83 (m, 2H), 4.60 (t, 2H, <i>J</i> = 13.6 Hz), 4.16 (s, 2H), 3.16 (s, 4H). MS (APCI): <i>m/e</i> 417.1 (M+1). CHN (for C ₂₀ H ₁₈ F ₆ N ₂ O•1HCl•0.25 H ₂ O) calcd: C 52.53, H 4.30, N 6.13; found: C 52.75, H 4.24, N 5.76.
886	4,5,6,7-tetrafluoro	2,2,3,3-tetra fluoropropyl	mp 262.5-263.8 °C; ¹ H NMR (400 MHz, dmso-d ₆): 12.16 (br s, 1H), 9.43 (s, 2H), 7.44 (d, 1H, <i>J</i> = 2.0 Hz), 7.34-7.40 (m, 2H) 7.19-7.21 (d, 1H, <i>J</i> = 3.6 Hz), 7.08-7.10 (dd, 1H, <i>J</i> = 2.0, 8.0 Hz), 6.69 (tt, 1H, <i>J</i> = 5.2, 52.0 Hz) 4.59 (t, 2H, <i>J</i> = 13.4 Hz), 4.15 (s, 2H), 3.16 (s, 4H). MS (APCI): <i>m/e</i> 453.1 (M+1). CHN (for C ₂₀ H ₁₆ F ₈ N ₂ O•1HCl•0.10 H ₂ O) calcd: C 48.96, H 3.53, N 5.71; found: C 48.74, H 3.33, N 5.61.
887	7-trifluoro methyl	2,2,2-trifluoroethyl	mp 173.8-175.6 °C. ¹ H NMR (400 MHz, dmso-d ₆): 11.36 (br s, 1H), 9.07 (br s, 1H), 7.87 (d, 1H, <i>J</i> = 7.6 Hz), 7.45 (d, 1H, <i>J</i> = 7.6 Hz) 7.38-7.42 (m, 1H), 7.36 (d, 1H, <i>J</i> = 2.4 Hz), 7.28-7.29 (m, 2H), 7.16-7.29 (m, 2H) 7.11 (dd, 1H, <i>J</i> = 2.0, 8.0 Hz), 4.77 (q, 2H, <i>J</i> = 8.8 Hz), 4.15 (s, 2H), 3.12-3.16 (m, 4H). MS (APCI): <i>m/e</i> 417.1 (M+1). CHN (for C ₂₀ H ₁₈ F ₆ N ₂ O•1HCl•0.20 H ₂ O) calcd: C 52.63, H 4.28, N 6.14; found: C 52.56, H 4.05, N 5.79.

-173-

888	7-trifluoro methyl	2,2,3,3-tetra fluoropropyl	mp 154.0-155.8 °C; ¹ H NMR (400 MHz, dmso-d ₆): 11.35 (br s, 1H), 9.51 (br s, 2H), 7.91 (d, 1H, <i>J</i> = 8.0 Hz), 7.36-7.45 (m, 4H) 7.22 (d, 1H, <i>J</i> = 8.0 Hz) 7.17 (t, 1H, <i>J</i> = 7.6 Hz), 7.09 (dd, 1H, <i>J</i> = 2.2, 8.0 Hz), 6.69 (tt, 1H, <i>J</i> = 5.2, 52.0 Hz), 4.60 (t, 2H, <i>J</i> = 13.6 Hz), 4.15 (s, 2H), 3.13-3.20 (m, 4H). MS (ES ⁺): <i>m/e</i> 449.0 (M+1). CHN (for C ₂₁ H ₁₉ F ₇ N ₂ O • 1HCl • 0.10 H ₂ O) calcd: C 51.83, H 4.18, N 5.76; found: C 51.54, H 3.97, N 5.68.
889	7-nitro	2,2,2-trifluoroethyl	mp 133.0-134.8 °C; ¹ H NMR (400 MHz, dmso-d ₆): 11.81 (s, 1H), 9.46 (br s, 2H) 8.14 (d, 1H, <i>J</i> = 8.0 Hz), 8.11 (d, 1H, <i>J</i> = 8.0 Hz) 7.45 (d, 1H, <i>J</i> = 2.0 Hz) 7.39 (t, 1H, <i>J</i> = 8.0 Hz), 7.36-7.37 (m, 1H), 7.25 (t, 1H, <i>J</i> = 8.0 Hz), 7.21 (d, 1H, <i>J</i> = 8.0 Hz),) 7.10 (dd, 1H, <i>J</i> = 2.0, 8.0 Hz), 4.78 (q, 2H, <i>J</i> = 8.8 Hz), 4.15 (s, 2H), 3.12-3.24 (m, 4H). MS (APCI): <i>m/e</i> 394.1 (M+1). CHN (for C ₁₉ H ₁₈ F ₃ N ₃ O ₃ • 1HCl • 0.80 H ₂ O) calcd: C 51.37, H 4.67, N 9.46; found: C 51.02, H 4.43, N 10.19.
890	7-nitro	2,2,3,3-tetra fluoropropyl	mp 175.0-176.8 °C; ¹ H NMR (400 MHz, dmso-d ₆): 11.81 (br s, 1H), 9.32 (br s, 2H), 8.13 (d, 1H, <i>J</i> = 8.0 Hz), 8.11 (d, 1H, <i>J</i> = 8.0 Hz) 7.45 (d, 1H, <i>J</i> = 2.0 Hz) 7.39 (t, 1H, <i>J</i> = 8.0 Hz), 7.31-7.32 (m, 1H), 7.25 (t, 1H, <i>J</i> = 8.0 Hz), 7.20 (d, 1H, <i>J</i> = 7.6 Hz),) 7.10 (dd, 1H, <i>J</i> = 2.4, 8.4 Hz), 6.69 (tt, 1H, <i>J</i> = 5.2, 52.0 Hz), 4.60 (t, 2H, <i>J</i> = 13.2 Hz), 4.16 (s, 2H), 3.18 (s, 4H). MS (APCI): <i>m/e</i> 426.1 (M+1). CHN (for C ₂₀ H ₁₉ F ₄ N ₃ O ₃ • 1HCl • 0.90 H ₂ O) calcd: C 50.25, H 4.60, N 8.79; found: C 49.98, H 4.38, N 9.47.

Example 8922-(7-Trifluoromethyl-1H-indol-3-yl)-ethylamine

- Combine in a 500 mL round bottom flask equipped with magnetic stirring, (2-trifluoromethyl-phenyl)-hydrazine (5.0 g, 28.4 mmol) and 4-aminobutyraldehyde dimethyl acetal (4.54 g, 34.1 mmol) and stir. After 5 minutes, slowly add 1N HCl (200 mL) and heat the reaction to 85° C for 2 hours forming an orange-red colored solution. Increase the temperature to 100° C for 10 minutes and cool to room temperature. Pour the reaction mixture over ice and stir for 10 minutes followed by adjustment to pH ~10 with ammonium hydroxide. Extract the mixture with methylene chloride, pool the organic phases, dry over sodium sulfate, and concentrate *in vacuo* to give a dark orange-brown oil. Purification on a pre-packed, HMDS treated silica column using a step gradient of 9% to 17 % methanol in methylene chloride gives the pure title compound as an orange oil: ¹H

-174-

NMR (400 MHz, dms -d_6): 11.18 (br s, 1H), 7.82 (d, 1H, $J = 7.6$ Hz), 7.40 (d, 1H, $J = 7.2$ Hz), 7.24 (d, 1H, $J = 2.0$ Hz), 7.13 (t, 1H, $J = 7.6$ Hz) 2.76-2.83 (m, 4H). MS (APCI): m/e 229.0 (M+1), 212.0 (M-NH $_2$).

5

Example 893(7-Nitro-1H-indol-3-yl)-acetonitrile

Dissolve in a 500 mL round bottom flask equipped with magnetic stirring, 7-nitro indole (4.55 g, 28.1 mmol) in 130 mL of glacial acetic acid and heat to 70° C. Add di-

- 10 After 45 minutes, cool the reaction mixture and remove the solvent *in vacuo* to give a crude yellow solid. Treat the crude material with 200 mL ammonium hydroxide and extract with ethyl acetate. Pool the organic phases and dry over magnesium sulphate and concentrate *in vacuo* to give the amine intermediate as a yellow crystalline solid. Immediately dissolve the intermediate in 200 mL of dimethyl sulphoxide, treat with
- 15 methyl iodide (4.55 mL, 56.2 mmole), and stir overnight at room temperature. Add potassium cyanide (18.30 g, 281 mmol), and 18-crown-6 (226 mg) and stir the mixture at 50°C for 25 minutes. Pour the resulting brown-yellow suspension over ice, stir for 10 minutes, saturate with sodium chloride, and extract with ethyl acetate. Wash the pooled extracts once with water, twice with brine, dry over sodium sulfate, and concentrate *in*
- 20 *vacuo* to give the title compound as a yellow-brown solid. No further purification was necessary. ^1H NMR (400 MHz, dms -d_6): 11.92 (br s, 1H) 8.14 (d, 1H, $J = 8.0$ Hz), 8.12 (d, 1H, $J = 8.0$ Hz) 7.53 (d, 1H, $J = 2.0$ Hz) 7.31 (t, 1H, $J = 8.0$ Hz), 4.16 (s, 2H), MS (ES-): m/e 200.0 (M-1).

25


Example 8942-(7-Nitro-1H-indol-3-yl)ethylamine

Dissolve in a 500 mL round bottom flask equipped with magnetic stirring, and a nitrogen inlet, (7-nitro-1H-indol-3-yl)-acetonitrile (5.27 g, 26 mmol) in dry tetrahydrofuran (150 mL). Treat the solution with 1M BH $_3$:THF (55mL, 55 mmol) and


30 stir at room temperature. After 20 hours, quench the reaction by the cautious dropwise addition of water(9 mL) and stir until foaming and gas evolution has stopped. Concentrate the mixture to dryness *in vacuo*, redissolve in 1 N HCl (300 ml) and extract with ethyl acetate. Basify the aqueous phase 5 N NaOH and extract with ethyl acetate. Pool the ethyl acetate extracts and dry over sodium sulfate and concentrate *in vacuo* to give the title

35 compound as an orange-brown solid: ^1H NMR (400 MHz, dms -d_6): 11.66 (br s, 1H) 8.07 (t, 2H, $J = 7.6$ Hz), 7.32 (s, 1H), 7.20 (t, 1H, $J = 8.0$ Hz) 2.79-2.83 (m, 4H), MS (APCI): m/e 189.0 (M-NH $_2$).

-175-

Example 895N-(2-(6-Fluoro-1H-indol-3-yl)ethyl)-4-fluoro-3-phenoxy-benzylamine

- 5 The method of Example 340 gives the hydrochloride of the title compound: mp 173 - 175 °C; MS(m/e): 379 (M+1), 377 (M-1); Calculated for $C_{23}H_{20}F_2N_2O \cdot HCl$: Calcd: C, 66.59; H, 5.10; N, 6.75. Found: C, 66.39; H, 5.05; N, 6.57.

Example 89610 N-(2-(6-Fluoro-1H-indol-3-yl)ethyl)-3-phenoxybenzylamine

- The method of Example 340 gives the hydrochloride of the title compound: mp 196 - 199 °C; MS(m/e): 361 (M+1), 359 (M-1); Calculated for $C_{23}H_{21}FN_2O \cdot HCl$: Calcd: C, 69.60; H, 5.59; N, 7.06. Found: C, 69.23; H, 5.58; N, 7.00.

15

Example 8974-Fluoro-1-methyl-3-phenoxybenzene

- Add triethylamine (28.6 mL, 205 mmol) dropwise to a mixture of 2-fluoro-5-methylphenol (5.18 g, 41.1 mmol), copper(II) acetate (7.46 g, 41.1 mmol), phenylboronic acid (10.0 g, 82.1 mmol), powdered 4Å sieves (7 g), and methylene chloride (400 mL).
 20 Stir at ambient temperature. After 22 h, filter and concentrate the filtrate. Purify the residue by silica gel chromatography (50% methylene chloride/hexanes), concentrate and purify again by silica
 gel chromatography (100% hexanes) to give 2.4 g (29%) of the title compound: MS(m/e):
 25 202 (M^+).

Example 8984-Fluoro-3-phenoxybenzaldehyde

- Combine 4-fluoro-1-methyl-3-phenoxybenzene (2.43 g, 12.0 mmol), *N*-bromosuccinimide (4.92 g, 27.6 mmol), benzoyl peroxide (408 mg, 1.68 mmol), and carbon tetrachloride (55 mL). Heat the mixture at reflux temperature for 6.5 h and cool to

30

-176-

- 0 °C for 64 h. Filter the solids and concentrate the filtrate. Dissolve the residue in chloroform and wash with ice cold sodium carbonate solution. Dry the chloroform solution over sodium sulfate, filter and concentrate under reduced pressure. Dissolve the residue in acetonitrile (50 mL) and add 4-methylmorpholine-4-oxide (4.6 g, 39.1 mmol) and powdered 4Å sieves (200 mg). Stir at ambient temperature for 20 h, filter and concentrate. Purify by silica gel chromatography (5%, 30% ethyl acetate/hexanes) to give 220 mg (8%) of the title compound: MS(m/e): 216 (M⁺).

Example 899

10 7-Fluorotryptamine

- Combine lithium aluminum hydride (12.8 g; 336.1 mmol) and 0 °C anhydrous tetrahydrofuran (160 mL). Cool resulting exotherm to 0 °C. Add 7-fluoro-3-(2-nitrovinyl)-1H-indole (11.55 g, 56.0 mmol) in anhydrous THF (200 mL) dropwise. After 30 min, warm to ambient temperature. After 4 hours, cool to 0 °C and add saturated sodium sulfate solution (35 mL) dropwise. Filter the solids and wash with THF and ethyl acetate. Concentrate the filtrate and dissolve the residue in methylene chloride. Filter the precipitate to give 1.26 g of product as brown crystals. Concentrate the filtrate and chromatograph on silica gel eluting with 5%, 7%, 10% 2N ammonia in methanol/methylene chloride to give product: MS(m/e): 179 (M+1), 177 (M-1);
- 20 Calculated for C₁₀H₁₁FN₂: Calcd: C, 67.40; H, 6.22; N, 15.72. Found: C, 67.06; H, 6.11; N, 15.48.

Example 900

3-(2-Nitrovinyl)-6-methanesulfonyl-1H-indole

- 25 Combine 1-dimethylamino-2-nitroethylene (892.1 mg, 7.68 mmol) and TFA (9.0 ml) and stir until dissolved. Add 6-methanesulfonyl-1H-indole (1.5 g, 7.68 mmol) and stir at ambient temperature. After 24 hours, pour the reaction mixture into ice/water, extract with ethyl acetate, then wash ethyl acetate with brine and saturated sodium bicarbonate. Filter, wash, and dry the precipitate to give the title compound as a yellow
- 30 powder: mp >250 °C. MS (ACPI): m/e 267.0 (M+1). Analysis for C₁₁H₁₀N₂O₄S: Calcd: C, 49.62; H, 3.79; N, 10.52; found: C, 49.86; H, 3.97; N, 10.25.

Example 901

3-(2-Nitrovinyl)-6-benzenesulfonyl-1H-indole

- 35 Combine 1-dimethylamino-2-nitroethylene (676.9 mg, 5.83 mmol) and TFA (9.0 ml) and stir until dissolved. Add 6-benzenesulfonyl-1H-indole (1.5 g, 5.83 mmol) and stir at

-177-

ambient temperature. After 24 hours, pour the reaction mixture into ice/water and adjust to pH 8. After stirring, filter the precipitate, wash with water, and dry to give the title compound as a yellow powder: mp 110 °C, dec. MS (ACPI): m/e 329.0 (M+1). Analysis for C₁₆H₁₂N₂O₄S: Calcd: C, 58.53; H, 3.68; N, 8.53; found: C, 58.54; H, 3.83; N, 7.85.

5

Example 902

(3-Phenoxybenzyl)-(2-pyridin-2-yl-ethyl)amine oxalic acid salt

Combine 2-pyridin-2-yl-ethylamine (Aldrich, 0.36 mL, 3.0 mmol), 3-phenoxybenzaldehyde (Aldrich, 0.58 mL, 3.66 mmol), 3A molecular sieves (0.5 g), and methanol (30 mL) and heat to reflux for 4 hours. Remove the molecular sieves by filtration. Add sodium borohydride (0.35 g, 9.0 mmol) slowly and stir the reaction at room temperature. After 1 hour, concentrate the reaction and dissolve the residue in a mixture of 1N NaOH solution and methylene chloride and extract the mixture with methylene chloride. Wash the organic extract with water, dry (Na₂SO₄) and concentrated to give a pale yellow oil. Form the salt with oxalic acid and crystallize from ethyl acetate to give a white solid: mp = 183 – 185°C; ms: ion at 305.2.

10

15

Example 903

(3-[1,3]Dioxolan-2-yl-phenyl)-pyridin-2-ylamine

Combine 2-aminopyridine (8.25g, 95mmol), 2-(3-bromo-phenyl)-[1,3]dioxolane (13.8 mL, 90mmol), sodium t-butoxide (12.2g, 126mmol), BINAP (210mg, 0.62mmol), Pd₂(dbu)₃ (630mg, 0.21mmol) and toluene (100mL) and heat to reflux for 48 hours. Cool the reaction to room temperature, dissolve in ether and filter and concentrate the resulting solution. Purification by flash chromatography (hexanes/EtOAc (8.5:1.5) and then hexanes/EtOAc (7:3)) provides the title compound as a yellow oil.

20

25

Example 904


3-(Pyridin-2-ylamino)-benzaldehyde

Dissolve (3-[1,3]dioxolan-2-yl-phenyl)-pyridin-2-yl-amine (10.32g, 42.6 mmol) in THF (150 mL). Add concentrated HCl solution (37.5 mL) and stir the solution at room temperature overnight. Concentrate the reaction, treat with water, and extract with CH₂Cl₂. Wash the organic extract with water, dry (Na₂SO₄) and concentrate to give the crude product. Purification by flash chromatography (hexanes/EtOAc (7:3)) provides the title compound as a yellow solid.

30


35

-178-

Example 905N-(3-(2-(6-Chloro-1H-indol-3-yl)ethyl)-3-(pyridin-2-ylamino)benzylamine

Combine 6-chlorotryptamine (0.22 g, 1.1 mmol), 3-(pyridin-2-ylamino)benzaldehyde (0.22 g, 1.1 mmol), 3A molecular sieves (0.5 g), and methanol (25 mL) and heat to reflux for 4 hours. Remove the molecular sieves by filtration. Add sodium borohydride (0.16g, 3.3 mmol) slowly and stir the reaction at room temperature. After 1 hour, concentrate the reaction and dissolve the residue in a mixture of 1N NaOH solution and methylene chloride and extract the mixture with methylene chloride. Wash the organic extract with water, dry (Na₂SO₄) and concentrate to give the crude product. Purification by flash chromatography (EtOAc/MeOH (9:1) with 2% concentrated NH₄OH solution) provides the desired product as a colorless oil. Form the dihydrochloride salt and crystallize from EtOAc to give the desired product: mp = 164 - 166°C; ms: ion at 377.1.


The following compounds were prepared following a procedure following Example 673:



No:	R ₁	R ₄	Data
906	pyrid-2-yl	phenyl	LC Method 3: R _f 2.83 min at 54/220nm; m/e 305.0 (M+1)
907	thien-2-yl	phenyl	LC Method 3: R _f 4.00 min at 254/220 nm; m/e 3309.9 (M+1)

20


The following compounds were prepared following a procedure following Example 673:



-179-

No.:	Z'	R ₄	Data
908	3-Br	propyl	LC Method 3: R _f 4.48 min at 254/220 nm; m/e 349.9 (M+1)
908a	3-COOCH ₃	phenyl	MS=362 (m+1), IR; 1718.51, 1584.26, 1489.84, 1445.78, 1285.67, 1253.07, 1199.51 cm ⁻¹


The following compounds were prepared following a similar procedure in Example 665:



No.:	Z'	R ₂	Data
909	H	isopropyl	LC Method 3: R _f 5.43 min at 254/220 nm; m/e 385.0(M+1)
910	methoxy	methyl	LC Method 2: R _f 4.86 min at 254/220 nm; m/e 385.0(M+1)

5

By the method of Example 221 the following compounds were prepared, isolated as the maleate except where noted:



No.	Z''	R ₄	Data
911	3-chloro	2-fluoro-benzyl	LC Method 3: R _f 4.61 min at 254/220 nm; m/e 369.9(M+1)
912	3-chloro	4-fluoro-benzyl	LC Method 3: R _f 4.62 min at 254/220 nm; m/e 369.9(M+1)
913	3-chloro	2,3-difluoro-benzyl	LC Method 3: R _f 4.76 min at 254/220 nm; m/e 387.9(M+1)

10

Example 914

3-Propoxybenzonitrile

Combine 3-hydroxybenzonitrile (11.052 gm; 92.8 mmol), n-propyl bromide (24.4 gm; 198 mmol), and potassium carbonate (38.65 gm; 280 mmol) in 2-butanone (175 mL) and heat and reflux. After 17 h., cool the mixture to room temperature, and decant the solution and concentrate by rotary evaporation. Partition the residue between diethyl ether

15

-180-

(150 mL) and water (150 mL), separate the layers and extract the aqueous layer with diethyl ether (2 x 100 mL). Combine the organic layers and wash with water, 1 N NaOH, and water, dry over MgSO₄, and concentrate. Distill the residue to give the title compound.

5

Example 915

3-Propoxybenzylamine hydrochloride

Combine 100 mL of lithium aluminum hydride (1 M in THF) and 50 mL of THF and add sulfuric acid (100%) dropwise at 10 °C. Allow the mixture to warm to room temperature and stir. After a 1 h period, remove the solids by filtration through diatomaceous earth using nitrogen pressure, and to the clear solution add dropwise a solution of nitrile in 50 mL of THF at 0 °C. Allow the reaction to stir. After 1 h at 0 °C, allow to warm to ambient temperature and stir over a 2.5 h period. Cool the reaction to 0 °C and add dropwise 16 mL of a 1:1 solution of water/THF, and add dropwise addition of 2 M NaOH (60 mL). Filter the resulting mixture, wash the solids with THF (2 X 100 mL), combine the organic layers dry over sodium sulfate and concentrate. Dissolve the residue in dry ether (250 mL) and acidify with HCl/dioxane solution (20 mL of 4 M solution). Wash the resulting solid with ether to yield the title compound as a white solid.

20

Example 916

2-(3-Bromophenyl)-N-(3-propoxybenzyl)acetamide

Combine 3-propoxy-benzylamine in 50 mL of dichloromethane and add dropwise to a mixture of 3-bromophenylacetyl chloride (4.90 gm; 21.0 mmol) and triethylamine (3.60 gm; 35.9 mmol) in 250 mL of dichloromethane at 0 °C. Allow the reaction to warm to room temperature and stir for 18 h. Pour the reaction into 100 mL of saturated brine, separate the layers and extract the aqueous layer with 100 mL of dichloromethane. Combine organic layers, wash with brine, dry (MgSO₄), and concentrate. Purification by chromatography on silica gel with 40% EtOAc in hexanes gives the title compound.

30

Example 917


2-(4'-Fluorobiphenyl-3-yl)-N-(3-propoxybenzyl)acetamide

Combine bromoamide (0.365 gm; 1.008 mmol), 4-fluorophenylboronic acid (0.175 gm; 1.25 mmol), cesium fluoride (0.360 gm; 2.37 mmol), and dichloro(bis(triphenylphosphine)palladium(II) (0.062 gm; 0.088 mmol) in NMP (3 mL) and heat at 104 °C. After 13.3 h. cool to ambient temperature and dilute with 40 mL each of dichloromethane and water. Separate the layers and extract the aqueous layer with dichloromethane (2 x 20 mL). Combine the organic layers, wash four times with 10 mL


35

-181-

portions of saturated brine, dry (MgSO_4) and concentrate. Purification by chromatography on silica gel with 40% EtOAc in hexanes gives the title compound.


Example 9185 N-(2-(3-(4-Fluorophenyl)phenyl)ethyl)-3-propoxybenzylamine

Combine 2-(4'-fluorobiphenyl-3-yl)-N-(3-propoxybenzyl)acetamide in 15 mL of THF and add a solution of $\text{BH}_3\text{-SMe}_2$ (2 M in THF) dropwise at 0 °C. Allow the reaction to warm to ambient temperature and stir. After 5 h. Cautiously add ethanol (1 mL), and
 10 concentrate the mixture. Dissolve the residue in ethanol (2 mL), heat to reflux for 2 h, and concentrate. Purification using gives the title compound as a tan solid. Dissolve the amine in 10 mL of 1:1 dichloromethane/methanol and add 600 mg of polyvinylpyridine hydrochloride. Shake the mixture for 4 h, removed by filtration the polymer and concentrate the residue and wash with ether to give the title compound as the
 15 hydrochloride: MS (ES⁺): m/e 364 (M+1).

Example 919N-(2-(5-Benzyloxy-1H-indol-3-yl)-ethyl)-3-phenoxybenzylamine

Combine 5-benzyloxy tryptamine (1.23 g, 4.6 mmol), 3-phenoxybenzaldehyde (97%, 1.09 g, 5.53 mmol) and molecular sieves 4A (1.0 g) and stir in methanol (15 mL) for 4h. Filter the molecular sieves and wash several times with MeOH. To this MeOH solution, add portionwise NaBH_4 (174 mg, 4.60 mmol), stir the resulting mixture at room temperature for 1h. Remove MeOH under vacuum, dilute the residue with CH_2Cl_2 /
 25 water, extract with CH_2Cl_2 , combine organic layers, dry over Na_2SO_4 and concentrate the solvent *in vacuo*. Purification by silica gel chromatography (CH_2Cl_2 / MeOH) to give the free base. Combine the free base with oxalic acid to form the salt: (300 MHz, DMSO-d_6) 2.95-3.15 (m, 4H), 3.93 (s, 2H), 4.10 (br, 1H), 5.05 (s, 2H), 6.85-7.46 (m, 18H), 10.67 (br, 1H); ms (ELECTROSPRAY) m/e: 449.2 (M+1).

-182-

Example 921N-(2-(5-Benzyloxy-1*H*-indol-3-yl)ethyl)-N-methyl-3-phenoxy-benzylamine


5 Combine N-(2-(5-benzyloxy-1*H*-indol-3-yl)ethyl)-3-phenoxy-benzylamine (1.61 g, 3.59 mmol) and NaOH (143.6 mg, 3.591.75 mmol) and dissolve in THF (25 mL) and stir at room temperature. After 15 min., add di-*tert*-butyl dicarbonate (1.57 g, 7.18 mmol) in THF (20 mL) and heat to reflux for 4 h. Remove the solvent, dilute with water, extract with CH₂Cl₂, (3 x 15 mL), dry over Na₂SO₄ and concentrate *in vacuo* to give a brown oil.

10 The crude product was used directly in the next step without purification.

Combine 1.0 M solution of LiAlH₄-THF (13.4 mL, 13.4 mmol) and (3-phenoxy-benzyl)-(2-(5-benzyloxy-1*H*-indol-3-yl)-ethyl)-carbamic acid *tert*-butyl ester (1.83 g, 3.34 mmol) and slowly add 15 mL dry THF. After addition, heat the reaction mixture to reflux. After 4.5h, cool down to room temperature. Quench the reaction by adding water (1.5 mL) cautiously, followed by 10 % NaOH. Filter off the suspension and wash repeatedly with ether. Dry the organic solution over Na₂SO₄ and concentrate the solvent *in vacuo*. Purification by silica gel chromatography using CH₂Cl₂ / MeOH as eluent to give the free base: ¹H NMR (300 MHz, CDCl₃) 2.35 (s, 3H), 2.69-2.74 (m, 2H), 2.91-2.96 (m, 2H), 3.65 (s, 2H), 5.07 (s, 2H), 6.90-7.53 (m, 18H), 7.80 (s, 1H). This compound reacted

15

20 further with oxalic acid to form the salt.


Example 922N-(2-(6,7-Difluoro-1*H*-indol-3-yl)-ethyl)-3-(pyridin-4-yloxy)benzylamine

25 Combine 6,7-difluoro tryptamine (0.285 g, 1.450mmol), 3-pyridin-4-yloxybenzaldehyde (0.303 g, 1.52 mmol, 1.05 eq.) and molecular sieves 4A (0.30 g) and stir in methanol (12 mL). After 4h. filter the molecular sieves and wash several times with MeOH. To this MeOH solution, add portionwise NaBH₄ (55.0 mg, 1.45 mmol), and stir at room temperature for 1h. Remove MeOH *in vacuo*, dilute the residue with CH₂Cl₂ /

-183-

water, extract with CH_2Cl_2 , combine organic layers, dry over Na_2SO_4 and concentrate *in vacuo*. Purification by flash chromatography on silica gel (CH_2Cl_2 / MeOH) to give the free base which is converted to the hydrochloride salt: ^1H NMR (300 MHz, DMSO- d_6) 3.13 (s, 4H), 4.20 (s, 2H), 6.85-7.55 (m, 10H), 8.47-8.50 (m, 1 H), 9.58 (br, 1H), 11.57 (br, 1H); MS (electrospray) m/e: 380.2 (M+1-HCl), 378.3 (M-1-HCl).

The present invention also provides novel intermediates of the compounds of formula I. The present invention provides intermediates of formula III:



wherein

R_3 is selected from the group consisting of hydrogen, fluoro, and methyl;
 R_4 is fluorinated C_2 - C_4 alkyl.

The present invention also provides novel crystalline forms of the compounds of formula I. Thus, for example, N-(2-(6-fluoro-1H-indol-3-yl)ethyl)-3-(2,2,3,3-tetrafluoropropoxy)benzylamine hydrochloride may be prepared by crystallization under controlled conditions to give novel crystalline forms. Crystallization from a solution and slurring techniques are contemplated to be within the scope of the present process. In practice, a number of factors can influence the form of N-(2-(6-fluoro-1H-indol-3-yl)ethyl)-3-(2,2,3,3-tetrafluoropropoxy)benzylamine hydrochloride obtained, including temperature and solvent composition. While the precise conditions under which crystalline N-(2-(6-fluoro-1H-indol-3-yl)ethyl)-3-(2,2,3,3-tetrafluoropropoxy)benzylamine hydrochloride is formed may be empirically determined it is only possible to give a number of methods which have been found to be suitable in practice. A preferred polymorphic form of N-(2-(6-fluoro-1H-indol-3-yl)ethyl)-3-(2,2,3,3-tetrafluoropropoxy)benzylamine hydrochloride can be prepared by crystallization or slurry from diethyl ether. Another preferred polymorphic form of N-(2-(6-fluoro-1H-indol-3-yl)ethyl)-3-(2,2,3,3-tetrafluoropropoxy)benzylamine hydrochloride can be prepared by crystallization from aqueous dichloromethane, aqueous acetone, ethyl acetate, ethyl acetate/cyclohexane, ethyl acetate/hexane, ethyl acetate/heptane, acetone/cyclohexane, isopropanol/hexanes, acetonitrile, acetonitrile/toluene, n-propanol/isoamylacetate/hexane, isopropyl acetate/ diethyl ether, methyl t-butyl ether/acetone, water, water/acetone, water/diethyl ether.

Crystalline N-(2-(6-fluoro-1H-indol-3-yl)ethyl)-3-(2,2,3,3-tetrafluoropropoxy)benzylamine may be prepared by direct crystallization under controlled conditions. The novel crystalline forms of the present invention may also be

-184-

prepared by dissolving (N-(2-(6-fluoro-1H-indol-3-yl)ethyl)-3-(2,2,3,3-tetrafluoropropoxy)benzylamine in a solvent and then forming the hydrochloride salt by the addition of a solution containing hydrochloric acid and then allowing crystallization while controlling the temperature.

5 A number of methods are available to characterize crystalline forms of organic compounds. For example methods include differential scanning calorimetry, solid state NMR spectrometry, infra-red spectroscopy, and X-ray powder diffraction. Among these X-ray powder diffraction and solid state NMR spectroscopy are very useful for identifying and distinguishing between crystalline forms.

10 X-ray powder diffraction analysis are performed by a variety of methods known to the skilled person. These methods can be varied to increase sensitivity by sample preparation techniques and by using more intense radiation, smaller scan steps, and slower scan rates. One method is as follows. Either with or without lightly grinding the sample with an agate mortar and pestle, the sample is loaded into a sample holder for the X-ray
15 powder diffraction measurement. The X-ray powder diffraction patterns are measured using a Siemens D5000 X-ray powder diffractometer equipped with a CuK_α source ($\lambda = 1.54056\text{\AA}$) operated at 50 kV and 40 mA using divergence slit size of 1 mm, receiving slit of 1 mm, and detector slit of 0.1 mm. Samples can be scanned between 4° and 35° (2θ) with a step size of 0.02° and a maximum scan rate of 3 sec/step. Data is collected using a
20 Kevex solid-state silicon lithium detector. Optimally, a silicon standard is run routinely to check the instrument alignment.

 It is well known in the crystallography art that, for any given crystal form, the relative intensities and peak widths of the diffraction peaks may vary due to a number of factors, including the effects of preferred orientation and/or particle size. Where the
25 effects of preferred orientation and/or particle size are present, peak intensities may be altered, but the characteristic peak positions of the polymorph are unchanged. See, e.g., The United States Pharmacopoeia #24, National Formulary #19, pages 1843-1844, 2000.

 Grinding may be used to minimize peak intensity. However, if grinding significantly alters the diffractogram or alters the crystalline state of the sample, then the
30 diffractogram of the unground sample should be used. Grinding is done in a small agate mortar and pestle. The mortar is held during the grinding and light pressure was applied to the pestle.

 Thus, a properly prepared sample crystalline compound of formula I may be characterized by one or more 2θ values in an X-ray diffraction pattern obtained as
35 described above.

-185-

Crystalline compounds of formula I may also be characterized by solid state NMR spectroscopy. Solid state ^{13}C chemical shifts reflect not only the molecular structure of but also the electronic environment of the molecule in the crystal.

Solid state NMR (^{13}C) analysis can be carried out using ^{13}C Cross polarization/magic angle spinning (CP/MAS). NMR (solid-state NMR or SSNMR) spectra are obtained using a Varian Unity 400 MHz spectrometer operating at a carbon frequency of 100.580 MHz, equipped with a complete solids accessory and Varian 7 mm VT CP/MAS probe. Acquisition parameters are readily determined and typically are 90° proton r.f. pulse width 4.0 μs , contact time 1.0 ms, pulse repetition time 5 s, MAS frequency 7.0 kHz, spectral width 50 kHz, and acquisition time 50 ms. Chemical shifts are generally reported by referenced to the methyl group of external hexamethylbenzene, that is, by sample replacement with hexamethylbenzene.

Thus, crystalline compounds of formula I may be characterized one or more resonances in the solid state ^{13}C nuclear magnetic spectra obtained as described above.

The compounds of the present invention can be administered alone or in the form of a pharmaceutical composition, that is, combined with pharmaceutically acceptable carriers or excipients. The compounds of the present invention, while effective themselves, may be formulated and administered in the form of their pharmaceutically acceptable salts, for purposes of stability, convenience, solubility, and the like. In practice, the compounds of formula I and II are usually administered in the form of pharmaceutical compositions, that is, in admixture with pharmaceutically acceptable carriers or diluents.

Thus, the present invention provides pharmaceutical compositions comprising a compound of the formula I or II and a pharmaceutically acceptable diluent.

The compounds of formula I and II can be administered by a variety of routes. In effecting treatment of a patient afflicted with disorders described herein, a compound of formula I and II can be administered in any form or mode which makes the compound bioavailable in an effective amount, including oral and parenteral routes. For example, compounds of formula I and II can be administered orally, by inhalation, subcutaneously, intramuscularly, intravenously, transdermally, intranasally, rectally, ocularly, topically, sublingually, buccally, and the like. Oral administration is generally preferred for treatment of the disorders described herein.

One skilled in the art of preparing formulations can readily select the proper form and mode of administration depending upon the particular characteristics of the compound selected, the disorder or condition to be treated, the stage of the disorder or condition, the solubility and chemical properties of the compound selected, the chosen route of administration, and other relevant circumstances considered in standard

-186-

pharmaceutical practice. (*Remington's Pharmaceutical Sciences*, 18th Edition, Mack Publishing Co. (1990)).

5 The pharmaceutical compositions are prepared in a manner well known in the pharmaceutical art. The carrier or excipient may be a solid, semi-solid, or liquid material which can serve as a vehicle or medium for the active ingredient. Suitable carriers or excipients are well known in the art. The pharmaceutical composition may be adapted for oral, inhalation, parenteral, or topical use and may be administered to the patient in the form of tablets, capsules, aerosols, inhalants, suppositories, solutions, suspensions, or the like.

10 The compounds of the present invention may be administered orally, for example, with an inert diluent or capsules or compressed into tablets. For the purpose of oral therapeutic administration, the compounds may be incorporated with excipients and used in the form of tablets, troches, capsules, elixirs, suspensions, syrups, wafers, chewing gums and the like. These preparations should contain at least 4% of the compound of the
15 present invention, the active ingredient, but may be varied depending upon the particular form and may conveniently be between 4% to about 70% of the weight of the unit. The amount of the compound present in compositions is such that a suitable dosage will be obtained. Preferred compositions and preparations according to the present invention may be determined by a person skilled in the art.

20 The tablets, pills, capsules, troches, and the like may also contain one or more of the following adjuvants: binders such as microcrystalline cellulose, gum tragacanth or gelatin; excipients such as starch or lactose, disintegrating agents such as alginic acid, Primogel, corn starch and the like; lubricants such as magnesium stearate or Sterotex; glidants such as colloidal silicon dioxide; and sweetening agents such as sucrose or
25 saccharin may be added or a flavoring agent such as peppermint, methyl salicylate or orange flavoring. When the dosage unit form is a capsule, it may contain, in addition to materials of the above type, a liquid carrier such as polyethylene glycol or a fatty oil. Other dosage unit forms may contain other various materials which modify the physical form of the dosage unit, for example, as coatings. Thus, tablets or pills may be coated
30 with sugar, shellac, or other coating agents. A syrup may contain, in addition to the present compounds, sucrose as a sweetening agent and certain preservatives, dyes and colorings and flavors. Materials used in preparing these various compositions should be pharmaceutically pure and non-toxic in the amounts used.

35 For the purpose of parenteral therapeutic administration, the compounds of the present invention may be incorporated into a solution or suspension. These preparations typically contain at least 0.1% of a compound of the invention, but may be varied to be between 0.1 and about 90% of the weight thereof. The amount of the compound of

-187-

formula I and II present in such compositions is such that a suitable dosage will be obtained. The solutions or suspensions may also include one or more of the following adjuvants: sterile diluents such as water for injection, saline solution, fixed oils, polyethylene glycols, glycerine, propylene glycol or other synthetic solvents; antibacterial agents such as benzyl alcohol or methyl paraben; antioxidants such as ascorbic acid or sodium bisulfite; chelating agents such as ethylene diaminetetraacetic acid; buffers such as acetates, citrates or phosphates and agents for the adjustment of tonicity such as sodium chloride or dextrose. The parenteral preparation can be enclosed in ampoules, disposable syringes or multiple dose vials made of glass or plastic. Preferred compositions and preparations are able to be determined by one skilled in the art.

The compounds of the present invention may also be administered topically, and when done so the carrier may suitably comprise a solution, ointment, or gel base. The base, for example, may comprise one or more of the following: petrolatum, lanolin, polyethylene glycols, bees wax, mineral oil, diluents such as water and alcohol, and emulsifiers, and stabilizers. Topical formulations may contain a concentration of the formula I and II or its pharmaceutical salt from about 0.1 to about 10% w/v (weight per unit volume).

The compounds of formula I and II are antagonists of 5-HT₆ receptors. Such antagonism can be identified by the methods below.

20

Example A

Assay for 5HT₆ binding

The assay buffer used is 50 mM Tris-HCl pH 7.4, 120 mM NaCl, 5 mM KCl, 5 mM MgCl₂, 1 mM EDTA. The radioligand used is ³H – LSD from New England Nuclear Cat. # NET 638 -75.9 Ci/mmol. The membranes used are from Receptor Biology, Cat. No. RB-HS6. These are membranes from HEK-293 cells expressing the Human 5HT₆ receptor.

Test compounds are obtained as 10 mM stocks in 100% DMSO. They are diluted to 1 mM in 100% DMSO by adding 180 µl DMSO to 20 µl of stock in 96 well plates using a multidrop. The 1 mM stocks are then diluted to make an 11 point concentration range from 125 µM down to 1.25 nM in half log increments using 10% DMSO as diluent. This is done using a TECAN robot. The final DMSO at this stage is 21.25%.

Radioligand is diluted in assay buffer to make a 125 nM solution and each vial of membranes is diluted up to 92 mL in assay buffer. The final assay volume is 250 µl consisting of 210 µl of diluted membranes, 20 µl of compound or 21.25% DMSO for total binding, and 20 µl of diluted radioligand. The compounds are transferred from drug dilution plates into corning 96 well assay plates using a 96 well Multimek pipettor.

-188-

Radioligand and membranes are added to assay plates using multidrop pipettors. Non-specific binding is determined in wells containing a final serotonin concentration of 10 μ M. In the final assay volume the radioligand is 10 nM and the membrane protein is approximately 25 μ g/well. The final drug concentration range in half logs is from 10 μ M
5 down to 0.1 nM. The final DMSO in the assay is 1.7%.

After addition of drug, membrane, and ligand, the plates are incubated for one hour at room temperature. During this time 96 well Millipore filter plates (MAFBN0B50) are soaked for a least 30 minutes with 200 μ l per well of 0.5% polyethyleneimine.

The 0.5% PEI is removed from filterplate wells using a TiterTek MAP aspirator
10 and 200 μ l of the incubation mixture is transferred from the incubation plate to the filterplate after mixing. This transfer is done using the 96 tip Mutimek pipettor. After transfer to the filterplate filterplates are extracted and washed twice with 220 μ l per well of cold buffer on the MAP aspirator. The peel away bottoms are removed from the filterplates and 100 μ l per well of microscint 20 scintillation fluid is added per well using
15 a multidrop. Plates are placed into suitable holders and are left at room temperature for three hours and are counted for ^3H in either a Wallac Microbeta counter or on a Packard Topcount.

In one embodiment, the present invention provides methods of treating disorders associated with the 5-HT₆ receptor, comprising: administering to a patient in need thereof
20 an effective amount of a compound of formula I. Thus, the present invention contemplates the various disorders described to be treated herein and others which can be treated by such antagonists as are appreciated by those skilled in the art.

In particular, because of their ability to antagonize the 5-HT₆ receptor, it is recognized that the compounds of the present invention are useful for treating cognitive
25 disorders, that is, disorders involving cognitive deficits. A number of the disorders which can be treated by 5-HT₆ antagonists are known according to established and accepted classifications, while others are not.

Some of the disorders to be treated according to the present invention are not well categorized and classified because cognition is a complicated and sometimes poorly
30 defined phenomenon. It is, however, widely recognized that cognition includes various "domains." These domains include short term memory, long term memory, working memory, executive function, and attention.

While many of the disorders which can be treated according to the present invention are not uniformly described and classified in the art, it is understood that the
35 compounds of the present invention are useful for treatment of disorders characterized by a deficit in any of the cognitive domains listed above or in other aspects of cognition. Thus the term "cognitive disorders" is meant to encompass any disorder characterized by

-189-

a deficit in one or more cognitive domain, including but not limited to short term memory, long term memory, working memory, executive function, and attention.

One cognitive disorder to be treated by the present invention is age-related cognitive decline. This disorder is not well defined, but includes decline in the cognitive domains, particularly the memory and attention domains, which accompany aging. Another is mild cognitive impairment. Again, this disorder is not well defined in the art, but involves decline in the cognitive domains, and is believed to represent a group of patients the majority of which have incipient Alzheimer's disease. Also, a wide variety of insults, including stroke, ischemia, hypoxia, inflammation, and infectious processes can result in cognitive deficits as a sequella which can be treated according to the present invention.

Where the disorders which can be treated by 5-HT₆ antagonists are, at present, known according to established and accepted classifications, these classifications can be found in various sources. For example, at present, the fourth edition of the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV™) (1994, American Psychiatric Association, Washington, D.C.), provides a diagnostic tool for identifying many of the disorders described herein. Also, the International Classification of Diseases, Tenth Revision, (ICD-10) provides classifications for many of the disorders described herein. The skilled artisan will recognize that there are alternative nomenclatures, nosologies, and classification systems for disorders described herein, including those not well characterized by the art and those described in the DMS-IV and ICD-10, and that terminology and classification systems evolve with medical scientific progress.

In one embodiment, the present invention provides methods of treating disorders selected from the group consisting of: age-related cognitive disorder, mild cognitive impairment, mood disorders (including depression, mania, bipolar disorders), psychosis (in particular schizophrenia), anxiety (particularly including generalized anxiety disorder, panic disorder, and obsessive compulsive disorder), idiopathic and drug-induced Parkinson's disease, epilepsy, convulsions, migraine (including migraine headache), substance withdrawal (including, substances such as opiates, nicotine, tobacco products, alcohol, benzodiazepines, cocaine, sedatives, hypnotics, etc.), sleep disorders (including narcolepsy), attention deficit/hyperactivity disorder, conduct disorder, learning disorders, dementia (including Alzheimer's disease and AIDS-induced dementia), Huntington's Chorea, cognitive deficits subsequent to cardiac bypass surgery and grafting, stroke, cerebral ischemia, spinal cord trauma, head trauma, perinatal hypoxia, cardiac arrest, and hypoglycemic neuronal damage, vascular dementia, multi-infarct dementia, amyotrophic lateral sclerosis, and multiple sclerosis, comprising: administering to a patient in need thereof an effective amount of a compound of formula I or II. That is, the present

-190-

invention provides for the use of a compound of formula I and II or pharmaceutical composition thereof for the treatment disorders associated with the 5-HT₆ receptor.

It is recognized that the terms "treatment" and "treating" are intended to include improvement of the cognitive deficit associated with each of the disorders associated with the 5-HT₆ receptor described herein. Also, it is also recognized that one skilled in the art may affect the disorders by treating a patient presently afflicted with the disorders or by prophylactically treating a patient believed to be susceptible to such disorders with an effective amount of the compound of formula I. Thus, the terms "treatment" and "treating" are intended to refer to all processes wherein there may be a slowing, interrupting, arresting, controlling, or stopping of the progression of the disorders described herein, but does not necessarily indicate a total elimination of all symptoms, and is intended to include prophylactic treatment of such disorders. For example, the present invention specifically encompasses the treatment of the cognitive deficits associated with schizophrenia, stroke, Alzheimer's disease, and the other disorders described herein. Thus, it is understood that the present invention includes adjunctive treatment of the disorders described herein. More specifically, the compounds of formula I and II are useful to treat cognition disorders in combination with a wide variety of therapeutic agents, in particular, in combination with AMPA potentiators; with typical and atypical antipsychotics, including olanzapine; with a variety of agents such as mGluR agonists, with NMDA antagonists, with IL 1-6 inhibitors, and the like; with cholinergics, including cholinesterase inhibitors, such as tacrine and donepezil, and compounds that inhibit amyloid protein processing, including inhibitors of amyloid precursor protein processing and antibodies directed against amyloid proteins; with antidepressants, including SSRIs; and with anxiolytic agents; etc. It is believed that the combinations above are synergistically beneficial providing efficacy at doses that are a small fraction of those required to produce the same effect with the individual components.

As used herein, the term "patient" refers to a warm blooded animal such as a mammal which is afflicted with one or more disorders associated with the 5-HT₆ receptor. It is understood that guinea pigs, dogs, cats, rats, mice, horses, cattle, sheep, pigs, and humans are examples of animals within the scope of the meaning of the term.

As used herein, the term "effective amount" of a compound of formula I or II refers to an amount, that is, the dosage which is effective in treating the disorders described herein.

An effective amount can be readily determined by the attending diagnostician, as one skilled in the art, by the use of conventional techniques and by observing results obtained under analogous circumstances. In determining an effective amount, the dose of a compound of formula I, a number of factors are considered by the attending

-191-

diagnostician, including, but not limited to: the compound of formula I or II to be administered; the co-administration of other therapies, if used; the species of mammal; its size, age, and general health; the specific disorder involved; the degree of involvement or the severity of the disorder; the response of the individual patient; the mode of administration; the bioavailability characteristics of the preparation administered; the dose regimen selected; the use of other concomitant medication; and other relevant circumstances.

An effective amount of a compound of formula I and II is expected to vary from about 0.1 milligram per kilogram of body weight per day (mg/kg/day) to about 100 mg/kg/day. Preferred amounts are able to be determined by one skilled in the art.

Of the disorders to be treated according to the present invention a number are particularly preferred.

In a preferred embodiment the present invention provides a method of treating cognitive disorders, comprising: administering to a patient in need thereof an effective amount of a compound of Claim 1.

In another preferred embodiment the present invention provides a method for treating Alzheimer's disease, comprising: administering to a patient in need thereof an effective amount of a compound of formula I.

In a preferred embodiment the present invention provides a method for treating schizophrenia, comprising: administering to a patient in need thereof an effective amount of a compound of formula I.

The fourth edition of the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV™) (1994, American Psychiatric Association, Washington, D.C.), provides a diagnostic tool including schizophrenia and related disorders, all of which are understood to be specifically included in the scope of this invention.

In a preferred embodiment the present invention provides a method for treating migraine, comprising: administering to a patient in need thereof an effective amount of a compound of formula I or II or a pharmaceutical composition thereof.

In one of the available sources of diagnostic tools, *Dorland's Medical Dictionary* (23rd Ed., 1982, W. B. Saunders Company, Philadelphia, PA), migraine is defined as a symptom complex of periodic headaches, usually temporal and unilateral, often with irritability, nausea, vomiting, constipation or diarrhea, and photophobia. As used herein the term "migraine" includes to these periodic headaches, both temporal and unilateral, the associated irritability, nausea, vomiting, constipation or diarrhea, photophobia, and other associated symptoms. The skilled artisan will recognize that there are alternative nomenclatures, nosologies, and classification systems for neurological and psychiatric

-192-

disorders, including migraine, and that these systems evolve with medical scientific progress.

In a preferred embodiment the present invention provides a method for treating anxiety disorders, including generalized anxiety disorder, panic disorder, and obsessive compulsive disorder, comprising: administering to a patient in need thereof an effective amount of a compound of formula I.

At present, the fourth edition of the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV™) (1994, American Psychiatric Association, Washington, D.C.), provides a diagnostic tool including anxiety and related disorders. These include: panic disorder with or without agoraphobia, agoraphobia without history of panic disorder, specific phobia, social phobia, obsessive-compulsive disorder, post-traumatic stress disorder, acute stress disorder, generalized anxiety disorder, anxiety disorder due to a general medical condition, substance-induced anxiety disorder and anxiety disorder not otherwise specified. As used herein the term "anxiety" includes treatment of those anxiety disorders and related disorders as specifically described in the DSM-IV and the term "anxiety" is intended to include like disorders that are described in other diagnostic sources.

A number of preclinical laboratory animal models have been described for the disorders described herein.

Example B

Fear Potentiated Startle Paradigm

Male Sprague-Dawley rats weighing 325-400 g were purchased from Harlan Sprague-Dawley, Inc. (Cumberland, IN) and given a one week acclimation period before testing. Rats were individually housed with food and water ad libitum in an animal room on a 12-hour light/dark cycle with lights on between 6:00 A.M. and 6:00 P.M. The compound of Example 16 was prepared in a suspension of 5% ethanol, 0.5% CMC, 0.5% Tween 80 and 99% water. 2S-2-amino-2-(1S,2S-2-carboxycyclopropan-1-yl)-3-(xanth-9-yl) propionic acid was prepared in sterile water. Control rats were given the respective vehicle.

The fear potentiated startle paradigm is conducted over three consecutive days. All three days begin with a 5-minute adaptation period before the trial starts. On day one (baseline startle) after the adaptation period, the animal receives 30 trials of 120dB auditory noise. The mean startle amplitude (V_{max}) is used to assign animals to groups with similar means before conditioning begins. Day two consists of conditioning the animals. Each animal receives 0.5 mA of shock for 500 msec preceded by a 5 second presentation of light which remains on for the duration of the shock. Ten presentations of the light and

-193-

shock are administered. Day three is the testing trial where drug administration occurs prior to testing. Twenty-four hours after conditioning, startle testing sessions are conducted. Ten trials of acoustic startle (120 dB), non-light paired, are presented at the beginning of the session. This is followed by 20 random trials of the noise alone and 20 random trials of noise preceded by light. Excluding the first 10 trials, the startle response amplitudes for each trial type are averaged for each animal. Data is presented as the difference between light + noise and noise-alone. Differences in startle response amplitudes were analyzed by Jmp statistical software using a One-way Anova (analysis of variance, t-test). Group differences were considered to be significant at $p < 0.05$.

The radial arm maze model can be used as a model of cognition and can be used to evaluate the present compounds.

Example C

Radial Arm Maze

The delayed non-match to sample task has been used to study the effect of drugs on memory retention (Pussinen, R. and Sirvio, J. *J of Psychopharm* 13: 171-179(1999); Staubli, U., et al. *Proc Natl Acad Sci* 91: 777-781(1994)) in the eight arm radial maze.

Well-trained rats were allowed to retrieve food rewards from four randomly selected arms of the maze (sampling phase). Some time later, the rats were exposed to eight open arms and were tested for their ability to remember and avoid the arms they had previously entered to obtain food. Re-entry into an arm that was baited during the sampling session was counted as a reference error, whereas entry into the same arm more than once during the retention session was counted as working error. The total (reference + working) number of errors made during the retention test increases with increasing delay periods. For example, young male rats made 0.66 (+ 0.4) errors at a 1 minute delay, 2 (+ 0.5) errors at a one hour delay, and 3.95 (+ 0.2) errors at a seven hour delay (observations of this lab).

Male Sprague-Dawley rats were individually housed and maintained on a 12h light-dark cycle (lights on at 6 am). The rats were given free access to water and maintained at 85% of their free-feeding weight by supplemental feedings of Purina Lab Chow.

The rats were initially trained to search for food at the end of each of the eight arms. Once the rats had reached the criteria of no more than two errors (i.e. entering the same arm more than once during a session) on three consecutive days, a delay of one minute was imposed between the fourth and the fifth arm choices. This training ensured that the rats were thoroughly familiar with the procedural aspects of the task before any drugs were administered. Once stable performance had been obtained on the delay task

-194-

(i.e. no more than one error was made on three consecutive days), drug and vehicle tests commenced using a seven hour delay period. A novel set of arms was baited each day for each rat and the maze was thoroughly cleaned during the delay period.

During the sampling session, each rat was placed on the center platform with access to all eight arms of the maze blocked. Four of the eight arms were randomly selected and baited with food. The gates of the baited arms were raised and the rat was allowed five minutes to obtain the food at the end of each of the four arms. As soon as the rat had obtained the food, it was removed, administered vehicle or various doses of compounds, and placed back in its home cage. Seven hours later (retention session), the rat was placed back onto the center platform with access to all eight arms blocked. The four arms that were previously baited during the sampling session, were baited and the gates to all eight arms were raised. The rat was allowed five minutes to obtain the remaining four pieces of food. An entry into a non-baited arm or a re-entry into a previously visited arm was counted as an error. Significance ($p < 0.05$) was determined using a repeated measure ANOVA followed by a Dunnett's test for comparison with control.

In order to compare test compounds with standards, scopolamine and tacrine were administered s.c. immediately after the sampling phase. The effects of scopolamine, a known amnesic, were tested after a three-hour delay, whereas the effect of tacrine, a cholinesterase inhibitor used in the treatment of Alzheimer's disease was tested after a six-hour delay. Scopolamine disrupted retention after a three-hour delay in a dose-related fashion. Tacrine significantly improved retention after a six-hour delay at 10, but not at 3 mg/kg.

25 Example D

Acquisition in the Radial Maze 8-arm radial maze acquisition

A prominent early feature of Alzheimer's disease (AD) symptomology is a pronounced deficit in declarative memory (R.W. Parks, R.F. Zec & R.S. Wilson (Eds.), *Neuropsychology of Alzheimer's disease and other dementias*. NY: Oxford University Press pp. 3-80 (1993).

As the disease progresses, other domains of cognition become severely affected as well. Among the brain regions affected early in the progression of AD is the hippocampus, which is a critical neural substrate for declarative memory (West M.J., Coleman P.D., Flood D.G. & Troncoso J.C.. Differences in the pattern of hippocampal neuronal loss in normal aging and Alzheimer's disease. *Lancet*, 344: 769-772(1994). One behavioral test that is often used to assess hippocampal function in animal models is the

-195-

8-arm radial maze (Olton D.S. The radial arm maze as a tool in behavioral pharmacology. *Physiology & Behavior*, 40: 793-797 (1986)).

Lesions or pharmacological blockade of the hippocampus disrupt performance of this task. Moreover, aged animals generally show deficits in this task (Porsolt R.D., Roux S. & Wettstein J.G. Animal models of dementia. *Drug Development Research*, 35: 214-229(1995)).

In this test of spatial learning and memory, a hungry rat is placed in the center of the maze and allowed to traverse the maze in search of food located at the end of each runway arm. In this version of the maze, the rat learns a win-shift strategy in which a visited arm is not replaced. Therefore, the most efficient foraging strategy is to visit each arm once. The version of the maze also taps into general learning processes as the rat is naïve to the maze on day one of the four day experiment.

Upon arrival, male Sprague Dawley®, rats were individually housed in a regular light-cycle colony room and allowed to acclimate for at least 4 days prior to testing. Each rat was reduced to and maintained at 85% of their target body weight throughout the experiment. Proper body weight was maintained by adjusting the allotment of lab chow based on a combination of age and the rat's daily bodyweight reading.

A session began with an individual rat being placed into the hub of the maze and then all guillotine doors were raised, allowing free access to all areas of the maze. A food hopper was located at the end of each of the 8 runway arms and a single food pellet was placed in each food hopper. Each daily session terminated when either all 8 food-hoppers had been visited or when the rat timed out (15 min on Day 1: 5 min on Days 2-4). The number of arm entries was recorded. Errors were counted as repeat arm entries or failures to visit an arm in the session period. An animal was excluded from the study if it failed to visit at least one arm on Day 1, 2 arms on Day 2, and at least 4 arms on Days 3 & 4.

Each rat was pseudo-randomly assigned to either a vehicle or drug group and received the same treatment throughout the experimental period. Vehicle consisted of 5% acacia within sterile water. Injections were administered subcutaneously 20-30 minutes prior to each daily session.


In this acquisition task, vehicle-treated animals do not consistently show significant acquisition of maze learning as compared to the number of errors committed on Day 1. We have found that in compounds that facilitate acquisition of maze learning, the effects are often not observed until the fourth day of training. Therefore, results consisted of total Day 4 errors across treatment groups.

35

-196-

WE CLAIM:

1. A compound of the formula



formula I

- 5 wherein
- X is selected from the group consisting of -O-, -NH-,
 -S-, -SO₂-, -CH₂-, -CH(F)-, -CH(OH)-, and -C(O)-;
- R₁ is selected from the group consisting of optionally substituted phenyl, optionally
 substituted naphthyl, optionally substituted 5 to 6 membered monocyclic aromatic
10 heterocycle having one heteroatom selected from the group consisting of nitrogen,
 oxygen, and sulfur and which 5 to 6 membered monocyclic aromatic heterocycle is
 optionally benzofused;
- R₂ is selected from the group consisting of hydrogen and C₁-C₃ alkyl;
- R₃ is selected from the group consisting of hydrogen, fluoro, and methyl;
- 15 R₄ is selected from the group consisting of hydrogen, allyl, C₂-C₄ alkyl, fluorinated C₂-C₄
 alkyl, optionally substituted phenyl, optionally substituted phenylsulfonyl, optionally
 substituted benzyl, and optionally substituted 5 to 6 membered monocyclic aromatic
 heterocycle having one or two heteroatoms selected from the group consisting of nitrogen,
 oxygen, and sulfur, provided that R₄ is not optionally substituted phenylsulfonyl when X
20 is -SO₂-, -CH₂-, CH(F)-,
 -CH(OH)-, or -C(O)-; and
- pharmaceutically acceptable salts thereof.
2. A compound according to Claim 1 wherein X is selected from the group consisting of
25 -O- and -NH-.
3. A compound according to Claim 2 wherein X is -O-.
4. A compound according to Claim 3 wherein R₃ is hydrogen.

30

-197-

5. A compound according to Claim 4 wherein R_4 is selected from the group consisting of C_2 - C_4 fluorinated alkyl and optionally substituted phenyl.
6. A compound according to Claim 5 wherein R_1 is selected from the group consisting of optionally substituted phenyl and optionally substituted 5 to 6 membered monocyclic aromatic heterocycle having one heteroatom selected from the group consisting of nitrogen, oxygen, and sulfur and which 5 to 6 membered monocyclic aromatic heterocycle is optionally benzofused.
7. A compound according to Claim 6 wherein R_1 is optionally substituted phenyl.
8. A compound according to Claim 6 wherein R_1 is optionally substituted 5 to 6 membered monocyclic aromatic heterocycle having one heteroatom selected from the group consisting of nitrogen, oxygen, and sulfur and which 5 to 6 membered monocyclic aromatic heterocycle is benzofused.
9. A compound according to Claim 8 wherein the benzofused 5 to 6 membered monocyclic aromatic heterocycle is optionally substituted indol-3-yl.
10. A compound according to any one of Claims 7 to 9 wherein R_4 is optionally substituted phenyl.
11. A compound according to any one of Claims 7 to 9 wherein R_4 is C_2 - C_4 fluorinated alkyl.
12. A compound of Claim 1 wherein the compound is N-(2-(6-Fluoro-1H-indol-3-yl)ethyl)-3-(2,2,3,3-tetrafluoropropoxy)benzylamine.
13. A compound of Claim 1 wherein the compound is N-(2-(6-Fluoro-1H-indol-3-yl)ethyl)-3-(2,2,3,3-tetrafluoropropoxy)benzylamine hydrochloride.
14. A compound of Claim 1 selected from the group consisting of N-(2-(5-Methoxy-6-fluoro-1H-indol-3-yl)ethyl)-3-(2,2,2-trifluoroethylamino)benzylamine, N-(2-(5-Methoxy-6-fluoro-1H-indol-3-yl)ethyl)-3-(2,2,3,3,3-pentafluoropropylamino)benzylamine, N-(2-(5-Methoxy-6-fluoro-1H-indol-3-yl)ethyl)-3-(2,2,3,3-tetrafluoropropoxy)benzylamine, N-(2-(5-Methoxy-6-fluoro-1H-indol-3-yl)ethyl)-3-(2,2,-trifluoroethoxy) benzylamine, N-(2-(5-Chloro-1H-indol-3-yl)ethyl)-3-(2,2,2-trifluoroethoxy)benzylamine, N-(2-(5-Chloro-1H-

-198-

- indol-3-yl)ethyl)-3-(3-fluoropropoxy)benzylamine, N-(2-(5-Chloro-1H-indol-3-yl)ethyl)-3-(2,2,3,3,3-pentafluoropropoxy) benzylamine, N-(2-(5-Chloro-1H-indol-3-yl)ethyl)-3-(2,2,3,3-tetrafluoropropoxy)benzylamine, N-(2-(5-Methoxy-1H-indol-3-yl)ethyl)-3-(2,2,2-trifluoroethoxy)benzylamine, N-(2-(5-Cyano-1H-indol-3-yl)ethyl)-3-(2,2,2-
- 5 trifluoroethoxy benzylamine, N-(2-(5-Methylsulfonyl-1H-indol-3-yl)ethyl)-3-(2,2,2-trifluoroethoxy)benzylamine, N-(2-(5-Cyano-1H-indol-3-yl)ethyl)-3-(3,3,3-trifluoropropoxy)benzylamine, N-(2-(5-Methylsulfonyl-1H-indol-3-yl)ethyl)-3-(3,3,3-trifluoropropoxy)benzylamine, N-(2-(4-Fluoro-1H-indol-3-yl)ethyl)-3-(2,2,2-trifluoroethoxy)benzylamine, N-(2-(4-Fluoro-1H-indol-3-yl)ethyl)-3-(2,2,3,3,3-
- 10 pentafluoropropoxy)benzylamine, N-(2-(4-Fluoro-1H-indol-3-yl)ethyl)-3-(2,2,3,3-tetrafluoropropoxy)benzylamine, N-(2-(7-Fluoro-1H-indol-3-yl)ethyl)-3-(2,2,2-trifluoroethoxy) benzylamine , N-(2-(7-Fluoro-1H-indol-3-yl)ethyl)-3-(2,2,3,3,3-pentafluoropropoxy)benzylamine, N-(2-(7-Fluoro-1H-indol-3-yl)ethyl)-3-(2,2,3,3-tetrafluoropropoxy)benzylamine, N-(2-(5-Amido-1H-indol-3-yl)ethyl)-3-(3,3,3-
- 15 trifluoropropoxy)benzylamine, N-(2-(5-Amido-1H-indol-3-yl)ethyl)-3-(2,2,2-trifluoroethoxy)benzylamine, N-(2-(6-Phenyl-1H-indol-3-yl)ethyl)-3-(2,2,2-trifluoroethoxy)benzylamine, N-(2-(6-Methyl-1H-indol-3-yl)ethyl)-3-(2,2,3,3,3-pentafluoropropoxy)benzylamine, N-(2-(6-Phenyl-1H-indol-3-yl)ethyl)-3-(2,2,3,3,3-
- 20 pentafluoropropoxy)benzylamine, N-(2-(6-Phenyl-1H-indol-3-yl)ethyl)-3-(2,2,3,3-tetrafluoropropoxy)benzylamine, N-(2-(6-Methyl-1H-indol-3-yl)ethyl)-3-(2,2,2-trifluoroethoxy)benzylamine, N-(2-(6-Methyl-1H-indol-3-yl)ethyl)-3-(2,2,3,3-tetrafluoropropoxy)benzylamine, N-(2-(6-Ethoxycarbonyl-1H-indol-3-yl)ethyl)-3-(2,2,3,3-
- 25 tetrafluoropropoxy)benzylamine, N-(2-(6-Ethoxycarbonyl-1H-indol-3-yl)ethyl)-3-(2,2,2-trifluoroethoxy)benzylamine, N-(2-(6-Cyano-1H-indol-3-yl)ethyl)-3-(2,2,2-trifluoroethoxy)benzylamine, N-(2-(6-Cyano-1H-indol-3-yl)ethyl)-3-(2,2,3,3-
- 30 tetrafluoropropoxy)benzylamine, N-(2-(6-Amido-1H-indol-3-yl)ethyl)-3-(2,2,2-trifluoroethoxy)benzylamine , N-(2-(6-Amido-1H-indol-3-yl)ethyl)-3-(2,2,3,3-tetrafluoropropoxy)benzylamine, N-(2-(6-Trifluoromethoxy-1H-indol-3-yl)ethyl)-3-(2,2,3,3-tetrafluoropropoxy) benzylamine, N-(2-(6-Trifluoromethoxy-1H-indol-3-
- 35 yl)ethyl)-3-(2,2,2-trifluoroethoxy)benzylamine, N-(2-(7-Chloro-1H-indol-3-yl)ethyl)-3-(2,2,3,3-tetrafluoropropoxy) benzylamine, N-(2-(7-Chloro-1H-indol-3-yl)ethyl)-3-(2,2,2-trifluoroethoxy)benzylamine, N-(2-(5-Trifluoromethyl-1H-indol-3-yl)ethyl)-3-(2,2,2-trifluoroethoxy)benzylamine , N-(2-(5-Trifluoromethyl-1H-indol-3-yl)ethyl)-3-(3,3,3-trifluoropropoxy)benzylamine, N-(2-(4-Methoxy-1H-indol-3-yl)ethyl)-3-(2,2,3,3-
- 35 tetrafluoropropoxy)benzylamine, N-(2-(5-Cyano-1H-indol-3-yl)ethyl)-3-(2,2,3,3-tetrafluoropropoxy)benzylamine, N-(2-(5-Cyano-1H-indol-3-yl)ethyl)-3-(2,2,3,3,3-pentafluoropropoxy)benzylamine, N-(2-(5-(4-Fluorophenyl)-1H-indol-3-yl)ethyl)-3-

-199-

- (2,2,3,3-tetrafluoropropoxy)benzylamine, N-(2-(5-(4-Fluorophenyl)-1H-indol-3-yl)ethyl)-3-(2,2,3,3,3-pentafluoropropoxy) benzylamine, N-(2-(5-Phenyl-1H-indol-3-yl)ethyl)-3-(2,2,3,3-tetrafluoropropoxy)benzylamine, N-(2-(5-Phenyl-1H-indol-3-yl)ethyl)-3-(2,2,3,3,3-pentafluoropropoxy)benzylamine, N-(2-(5-(4-Fluorophenyl)-1H-indol-3-yl)ethyl)-3-(2,2,2-trifluoroethoxy)benzylamine, N-(2-(5-Phenyl-1H-indol-3-yl)ethyl)-3-(2,2,2-trifluoroethoxy)benzylamine, N-(2-(4-Phenyl-1H-indol-3-yl)ethyl)-3-(2,2,3,3,3-pentafluoropropoxy)benzylamine, N-(2-(4-Phenyl-1H-indol-3-yl)ethyl)-3-(2,2,2-trifluoroethoxy)benzylamine, N-(2-(4-Phenyl-1H-indol-3-yl)ethyl)-3-(2,2,3,3-tetrafluoropropoxy)benzylamine, N-(2-(7-Phenyl-1H-indol-3-yl)ethyl)-3-(2,2,2-trifluoroethoxy)benzylamine, N-(2-(7-Phenyl-1H-indol-3-yl)ethyl)-3-(2,2,3,3-tetrafluoropropoxy)benzylamine, N-(2-(6-Chloro-1H-indol-3-yl)ethyl)-N-methyl-3-(2,2,3,3-tetrafluoropropoxy) benzylamine, N-(2-(4-Methoxy-1H-indol-3-yl)ethyl)-N-methyl-3-(2,2,3,3-tetrafluoropropoxy)benzylamine, N-(2-(4-Fluoro-1H-indol-3-yl)ethyl)-N-methyl-3-(2,2,2-trifluoroethoxy) benzylamine, N-(2-(6-Phenyl-1H-indol-3-yl)ethyl)-N-methyl-3-(2,2,3,3-tetrafluoropropoxy) benzylamine, N-(2-(6-Carboxy-1H-indol-3-yl)ethyl)-3-(2,2,3,3-tetrafluoropropoxy) benzylamine, N-(2-(6-Carboxy-1H-indol-3-yl)ethyl)-3-(2,2,2-trifluoroethoxy)benzylamine, N-(2-(6-Chloro-7-fluoro-1H-indol-3-yl)ethyl)-3-(2,2,2-trifluoroethoxy)benzylamine, N-(2-(6-Chloro-7-fluoro-1H-indol-3-yl)ethyl)-3-(2,2,3,3-tetrafluoropropoxy) benzylamine, N-(2-(5,7-Difluoro-1H-indol-3-yl)ethyl)-3-(2,2,2-trifluoroethoxy)benzylamine, N-(2-(5,7-Difluoro-1H-indol-3-yl)ethyl)-3-(2,2,3,3-tetrafluoropropoxy) benzylamine, N-(2-(6,7-Difluoro-1H-indol-3-yl)ethyl)-3-(2,2,2-trifluoroethoxy)benzylamine, N-(2-(6,7-Difluoro-1H-indol-3-yl)ethyl)-3-(2,2,3,3-tetrafluoropropoxy) benzylamine, N-(2-(5,6,7-Trifluoro-1H-indol-3-yl)ethyl)-3-(2,2,2-trifluoroethoxy)benzylamine, N-(2-(5,6,7-Trifluoro-1H-indol-3-yl)ethyl)-3-(2,2,3,3-tetrafluoropropoxy)benzylamine, N-(2-(4,5,7-Trifluoro-1H-indol-3-yl)ethyl)-3-(2,2,2-trifluoroethoxy)benzylamine, N-(2-(4,5,7-Trifluoro-1H-indol-3-yl)ethyl)-3-(2,2,3,3-tetrafluoropropoxy)benzylamine, N-(2-(7-Cyano-1H-indol-3-yl)ethyl)-3-(2,2,2-trifluoroethoxy)benzylamine, N-(2-(7-Cyano-1H-indol-3-yl)ethyl)-3-(2,2,3,3-tetrafluoropropoxy) benzylamine, N-(2-(5-Fluoro-1H-indol-3-yl)ethyl)-3-(2,2,2-trifluoroethoxy)benzylamine, N-(2-(5-Fluoro-1H-indol-3-yl)ethyl)-3-(2,2,3,3,3-pentafluoropropoxy)benzylamine, N-(2-(5-Methoxy-1H-indol-3-yl)ethyl)-3-(2,2,3,3,3-pentafluoropropoxy) benzylamine, N-(2-(5-Methoxy-1H-indol-3-yl)ethyl)-3-(2,2,3,3-tetrafluoropropoxy)benzylamine, N-(2-(5-Methoxy-1H-indol-3-yl)ethyl)-3-(3,3,3-trifluoropropoxy) benzylamine, N-(2-(5-Fluoro-1H-indol-3-yl)ethyl)-3-(2,2,3,3-tetrafluoropropoxy)benzylamine, N-(2-(4-Chloro-5-methoxy-1H-indol-3-yl)ethyl)-3-(2,2,2-trifluoroethoxy) benzylamine, N-(2-(4-Chloro-5-methoxy-1H-indol-3-yl)ethyl)-3-(2,2,3,3-tetrafluoropropoxy)benzylamine, N-(2-(4-Chloro-5-methoxy-1H-indol-3-

-200-

- yl)ethyl)-3-(3,3,3-trifluoropropoxy)benzylamine, N-(2-(6-Fluoro-1H-indol-3-yl)ethyl)-3-(2,2,2-trifluoroethoxy)benzylamine, N-(2-(6-Fluoro-1H-indol-3-yl)ethyl)-3-(2,2,3,3,3-pentafluoropropoxy)benzylamine, N-(2-(6-Chloro-5-methoxy-1H-indol-3-yl)ethyl)-3-(2,2,2-trifluoroethoxy) benzylamine, N-(2-(6-Chloro-5-methoxy-1H-indol-3-yl)ethyl)-3-
- 5 (2,2,3,3-tetrafluoropropoxy) benzylamine, N-(2-(6-Fluoro-1H-indol-3-yl)ethyl)-3-(3,3,3-trifluoropropoxy)benzylamine, N-(2-(5-Trifluoromethoxy-1H-indol-3-yl)ethyl)-3-(2,2,3,3,3-pentafluoropropoxy) benzylamine, N-(2-(5-Trifluoromethoxy-1H-indol-3-yl)ethyl)-3-(2,2,3,3-tetrafluoropropoxy)benzylamine, N-(2-(5-Trifluoromethoxy-1H-indol-3-yl)ethyl)-3-(2,2,2-trifluoroethoxy)benzylamine, N-(2-(5-Nitro-1H-indol-3-
- 10 yl)ethyl)-3-(2,2,2-trifluoroethoxy)benzylamine, N-(2-(5-Nitro-1H-indol-3-yl)ethyl)-3-(2,2,3,3-tetrafluoropropoxy)benzylamine, N-(2-(6-Nitro-1H-indol-3-yl)ethyl)-3-(2,2,2-trifluoroethoxy)benzylamine, N-(2-(6-Nitro-1H-indol-3-yl)ethyl)-3-(2,2,3,3-tetrafluoropropoxy)benzylamine, N-(2-(6-Nitro-1H-indol-3-yl)ethyl)-3-(2,2,3,3,3-pentafluoropropoxy)benzylamine, N-(2-(5-Amino-1H-indol-3-yl)ethyl)-3-(2,2,2-
- 15 trifluoroethoxy)benzylamine, N-(2-(5-Amino-1H-indol-3-yl)ethyl)-3-(2,2,3,3-tetrafluoropropoxy)benzylamine, N-(2-(6-Amino-1H-indol-3-yl)ethyl)-3-(2,2,2-trifluoroethoxy) benzylamine, N-(2-(6-Amino-1H-indol-3-yl)ethyl)-3-(2,2,3,3-tetrafluoropropoxy)benzylamine, N-(2-(5-Propoxy-1H-indol-3-yl)ethyl)-3-(2,2,3,3-tetrafluoropropoxy)benzylamine, N-(2-(5-n-Propyl amido-1H-indol-3-yl)ethyl)-3-(2,2,2-
- 20 trifluoroethoxy)benzylamine, N-(2-(5-Ethoxycarbonyl-1H-indol-3-yl)ethyl)-3-(2,2,3,3-tetrafluoropropoxy) benzylamine, N-(2-(5-Phenoxy-1H-indol-3-yl)ethyl)-3-(2,2,3,3-tetrafluoropropoxy)benzylamine, N-(2-(1H-Indol-3-yl)ethyl)-3-(2,2,2-trifluoroethoxy)benzylamine, N-(2-(1H-Indol-3-yl)ethyl)-3-(2,2,3,3,3-
- 25 pentafluoropropoxy) benzylamine, N-(2-(5-n-Butylamido-1H-indol-3-yl)ethyl)-3-(2,2,2-trifluoroethoxy)benzylamine, N-(2-(5-Hydroxy-1H-indol-3-yl)ethyl)-3-(2,2,2-trifluoroethoxy)benzylamine, N-(2-(5-Benzyloxy-1H-indol-3-yl)ethyl)-3-(2,2,3,3,3-pentafluoropropoxy)benzylamine, N-(2-(6-Benzyloxy-1H-indol-3-yl)ethyl)-3-(2,2,2-
- 30 trifluoroethoxy)benzylamine, N-(2-(6-Benzyloxy-1H-indol-3-yl)ethyl)-3-(2,2,3,3-tetrafluoropropoxy)benzylamine, N-(2-(6-Butyloxy-1H-indol-3-yl)ethyl)-3-(2,2,2-trifluoroethoxy)benzylamine, N-(2-(5-Butyloxy-1H-indol-3-yl)ethyl)-3-(2,2,3,3-tetrafluoropropoxy)benzylamine, N-(2-(6-Ethoxy-1H-indol-3-yl)ethyl)-3-(2,2,2-
- 35 trifluoroethoxy)benzylamine, N-(2-(6-Phenylsulfonyloxy-1H-indol-3-yl)ethyl)-3-(2,2,2-trifluoroethoxy)benzylamine, N-(2-(6-Phenylsulfonyloxy-1H-indol-3-yl)ethyl)-3-(2,2,3,3-tetrafluoropropoxy) benzylamine, N-(2-(6-Butyloxy-1H-indol-3-yl)ethyl)-3-(2,2,3,3-tetrafluoropropoxy)benzylamine, N-(2-(5-Hydroxy-1H-indol-3-yl)ethyl)-3-(2,2,2-
- trifluoroethoxy) benzylamine, N-(2-(5-Hydroxy-1H-indol-3-yl)ethyl)-3-(2,2,3,3-tetrafluoropropoxy)benzylamine, N-(2-(6-Hydroxy-1H-indol-3-yl)ethyl)-3-(2,2,2-

-201-

- trifluoroethoxy) benzylamine, N-(2-(6-Hydroxy-1H-indol-3-yl)ethyl)-3-(2,2,3,3-tetrafluoropropoxy) benzylamine, N-(2-(5-Carboxy-1H-indol-3-yl)ethyl)-3-(2,2,2-trifluoroethoxy) benzylamine, N-(2-(6-Chloro-1H-indol-3-yl)ethyl)-3-(3-fluoropropoxy) benzylamine, N-(2-(6-Fluoro-1H-indol-3-yl)ethyl)-3-(3-fluoropropoxy) benzylamine, N-(2-(6-Fluoro-1H-indol-3-yl)ethyl)-3-(2,2-difluoroethoxy) benzylamine, N-(2-(6-Chloro-1H-indol-3-yl)ethyl)-3-(2,2-difluoroethoxy) benzylamine, N-(2-(6-Chloro-1H-indol-3-yl)ethyl)-3-(2,2,3,3,3-pentafluoropropoxy) benzylamine, N-(2-(5-Isopropyl-1H-indol-3-yl)ethyl)-3-(2,2,3,3,3-pentafluoropropoxy) benzylamine, N-(2-(5-Isopropyl-1H-indol-3-yl)ethyl)-3-(2,2,3,3-tetrafluoropropoxy) benzylamine, N-(2-(6-Chloro-1H-indol-3-yl)ethyl)-N-methyl-3-(2,2-difluoroethoxy) benzylamine, N-(2-(6-Fluoro-1H-indol-3-yl)ethyl)-N-methyl-3-(3-fluoropropoxy) benzylamine, N-(2-(6-Chloro-1H-indol-3-yl)ethyl)-N-methyl-3-(3-fluoropropoxy) benzylamine, N-(2-(6-Chloro-1H-indol-3-yl)ethyl)-N-isopropyl-3-(2,2,3,3,3-pentafluoropropoxy) benzylamine, N-(2-(6-Chloro-1H-indol-3-yl)ethyl)-N-propyl-3-(2,2,3,3,3-pentafluoropropoxy) benzylamine, N-(2-(6-Chloro-1H-indol-3-yl)ethyl)-N-ethyl-3-(2,2,3,3,3-pentafluoropropoxy) benzylamine, N-(2-(6-Chloro-5-methoxy-1H-indol-3-yl)ethyl)-3-(2,2,3,3-tetrafluoropropoxy) benzylamine, N-(2-(6-Chloro-5-methoxy-1H-indol-3-yl)ethyl)-3-(2,2,2-trifluoroethoxy) benzylamine, N-(2-(4-Chloro-5-methoxy-1H-indol-3-yl)ethyl)-3-(2,2,2-trifluoroethoxy) benzylamine, N-(2-(4-Chloro-5-methoxy-1H-indol-3-yl)ethyl)-3-(2,2,3,3-tetrafluoropropoxy) benzylamine, N-(2-(5-Methoxy-1H-indol-3-yl)ethyl)-3-(2-fluoroethoxy) benzylamine, N-(2-(6-Methoxy-1H-indol-3-yl)ethyl)-3-(2,2,3,3,3-pentafluoropropoxy) benzylamine, N-(2-(4-Chloro-1H-indol-3-yl)ethyl)-3-(2,2,3,3,3-pentafluoropropoxy) benzylamine, N-(2-(4-Methoxy-1H-indol-3-yl)ethyl)-3-(2,2,3,3,3-pentafluoropropoxy) benzylamine, N-(2-(5-Methoxy-2-methyl-1H-indol-3-yl)ethyl)-3-(2,2,3,3,3-pentafluoropropoxy) benzylamine, N-(2-(7-Methoxy-1H-indol-3-yl)ethyl)-3-(2,2,3,3,3-pentafluoropropoxy) benzylamine, N-(2-(4-Methoxy-1H-indol-3-yl)ethyl)-3-(2,2,3,3-tetrafluoropropoxy) benzylamine, N-(2-(5-Methoxy-2-methyl-1H-indol-3-yl)ethyl)-3-(2,2,3,3-tetrafluoropropoxy) benzylamine, N-(2-(7-Methoxy-1H-indol-3-yl)ethyl)-3-(2,2,3,3-tetrafluoropropoxy) benzylamine, N-(2-(6-Methoxy-1H-indol-3-yl)ethyl)-3-(2-fluoroethoxy) benzylamine, N-(2-(4-Chloro-1H-indol-3-yl)ethyl)-3-(2-fluoroethoxy) benzylamine, N-(2-(4-Methoxy-1H-indol-3-yl)ethyl)-3-(2-fluoroethoxy) benzylamine, N-(2-(5-Methoxy-2-methyl-1H-indol-3-yl)ethyl)-3-(2-fluoroethoxy) benzylamine, N-(2-(7-Methoxy-1H-indol-3-yl)ethyl)-3-(2-fluoroethoxy) benzylamine, N-(2-(6-Chloro-1H-indol-3-yl)ethyl)-3-(2-fluoroethoxy) benzylamine, N-(2-(6-Methoxy-1H-indol-3-yl)ethyl)-3-(2,2-difluoroethoxy) benzylamine, N-(2-(6-Fluoro-1H-indol-3-yl)ethyl)-3-(2-fluoroethoxy) benzylamine, N-(2-(4,6-Dichloro-5-methoxy-1H-indol-3-yl)ethyl)-3-(2,2-

-202-

- difluoroethoxy) benzylamine, N-(2-(4-Chloro-1H-indol-3-yl)ethyl)-3-(2,2-difluoroethoxy)benzylamine, N-(2-(4-Methoxy-1H-indol-3-yl)ethyl)-3-(2,2-difluoroethoxy) benzylamine, N-(2-(5-Methoxy-2-methyl-1H-indol-3-yl)ethyl)-3-(2,2-difluoroethoxy)benzylamine, N-(2-(7-Methoxy-1H-indol-3-yl)ethyl)-3-(2,2-difluoroethoxy)benzylamine, N-(2-(6-Methoxy-1H-indol-3-yl)ethyl)-3-(3-fluoropropoxy) benzylamine, N-(2-(4-Chloro-1H-indol-3-yl)ethyl)-3-(3-fluoropropoxy)benzylamine, N-(2-(4-Methoxy-1H-indol-3-yl)ethyl)-3-(3-fluoropropoxy)benzylamine, N-(2-(5-Methoxy-2-methyl-1H-indol-3-yl)ethyl)-3-(3-fluoropropoxy) benzylamine, N-(2-(7-Methoxy-1H-indol-3-yl)ethyl)-3-(3-fluoropropoxy)benzylamine, N-(2-(6-Chloro-1H-indol-3-yl)ethyl)-3-(3-fluoropropoxy)benzylamine, N-(2-(6-Methoxy-1H-indol-3-yl)ethyl)-3-(2,2,2-trifluoroethoxy)benzylamine, N-(2-(4-Chloro-1H-indol-3-yl)ethyl)-3-(2,2,2-trifluoroethoxy)benzylamine, N-(2-(4-Methoxy-1H-indol-3-yl)ethyl)-3-(2,2,2-trifluoroethoxy) benzylamine, N-(2-(5-Methoxy-2-methyl-1H-indol-3-yl)ethyl)-3-(2,2,2-trifluoroethoxy)benzylamine, N-(2-(7-Methoxy-1H-indol-3-yl)ethyl)-3-(2,2,2-trifluoroethoxy)benzylamine, N-(2-(6-Chloro-1H-indol-3-yl)ethyl)-3-(2,2,2-trifluoroethoxy) benzylamine, N-(2-(6-Methoxy-1-methyl-1H-indol-3-yl)ethyl)-3-(2,2-difluoroethoxy)benzylamine, N-(2-(5-Methoxy-4,6-difluoro-1-methyl-1H-indol-3-yl)ethyl)-3-(3-fluoropropoxy)benzylamine, N-(2-(5-Methoxy-4,6-difluoro-1-methyl-1H-indol-3-yl)ethyl)-3-(2,2,2-trifluoroethoxy) benzylamine, N-(2-(5-Methoxy-1-methyl-1H-indol-3-yl)ethyl)-3-(2,2,3,3-tetrafluoropropoxy) benzylamine, N-(2-(5-Methoxy-1-methyl-1H-indol-3-yl)ethyl)-3-(2,2,2-trifluoroethoxy) benzylamine, N-(2-(6-Trifluoromethyl-1H-indol-3-yl)ethyl)-3-(3,3,3-trifluoropropoxy)benzylamine, N-(2-(5-Fluoro-6-chloro-1H-indol-3-yl)ethyl)-3-(3,3,3-trifluoropropoxy) benzylamine, N-(2-(5,6-Difluoro-1H-indol-3-yl)ethyl)-3-(3,3,3-trifluoropropoxy)benzylamine, N-(2-(5-Fluoro-6-chloro-1H-indol-3-yl)ethyl)-3-(2,2-difluoroethoxy) benzylamine, N-(2-(5,6-Difluoro-1H-indol-3-yl)ethyl)-3-(2,2-difluoroethoxy)benzylamine, N-(2-(6-Trifluoromethyl-1H-indol-3-yl)ethyl)-3-(2,2,2-trifluoroethoxy)benzylamine, N-(2-(5-Fluoro-6-chloro-1H-indol-3-yl)ethyl)-3-(2,2,3,3,3-pentafluoropropoxy) benzylamine, N-(2-(5,6-Difluoro-1H-indol-3-yl)ethyl)-3-(2,2,3,3,3-pentafluoropropoxy)benzylamine, N-(2-(6-Trifluoromethyl-1H-indol-3-yl)ethyl)-3-(2,2,3,3,3-pentafluoropropoxy)benzylamine, N-(2-(5,6-Difluoro-1H-indol-3-yl)ethyl)-3-(2,2,2-trifluoroethoxy)benzylamine, N-(2-(5,6-Difluoro-1H-indol-3-yl)ethyl)-3-(2,2,2-trifluoroethoxy)benzylamine, N-(2-(5,6-Difluoro-1H-indol-3-yl)ethyl)-3-(3-fluoropropoxy)benzylamine, N-(2-(6-Trifluoromethyl-1H-indol-3-yl)ethyl)-3-(3-fluoropropoxy)benzylamine, N-(2-(6-Fluoro-1-methyl-1H-indol-3-yl)ethyl)-3-(3,3,3-trifluoropropoxy)benzylamine, N-(2-(6-Fluoro-1-methyl-1H-indol-3-yl)ethyl)-3-(2,2,3,3,3-pentafluoropropoxy)benzylamine, N-(2-(6-Fluoro-1-methyl-1H-indol-3-

- yl)ethyl)-3-(2,2,2-trifluoroethoxy)benzylamine, N-(2-(5-Methyl-1H-indol-3-yl)ethyl)-3-(2-fluoroethoxy)benzylamine, N-(2-(5-Fluoro-1H-indol-3-yl)ethyl)-3-(2-fluoroethoxy)benzylamine, N-(2-(5-Methyl-1H-indol-3-yl)ethyl)-3-(2,2-difluoroethoxy)benzylamine, N-(2-(5-Fluoro-1H-indol-3-yl)ethyl)-3-(2,2-difluoroethoxy)benzylamine, N-(2-(5-Chloro-1H-indol-3-yl)ethyl)-3-(2,2-difluoroethoxy)benzylamine, N-(2-(5-Fluoro-1H-indol-3-yl)ethyl)-3-(4,4,4-trifluorobutoxy) benzylamine, N-(2-(5-Fluoro-1H-indol-3-yl)ethyl)-3-(2,2,2-trifluoroethoxy)benzylamine, N-(2-(5-Chloro-1H-indol-3-yl)ethyl)-3-(4,4,4-trifluorobutoxy)benzylamine, N-(2-(5-Fluoro-1H-indol-3-yl)ethyl)-3-(3-fluoropropoxy) benzylamine, N-(2-(5-Chloro-1H-indol-3-yl)ethyl)-3-(3-fluoropropoxy)benzylamine, N-(2-(5-Methyl-1H-indol-3-yl)ethyl)-3-(2,2,3,3-tetrafluoropropoxy)benzylamine, N-(2-(5-Methyl-1H-indol-3-yl)ethyl)-3-(2,2,3,3,3-pentafluoropropoxy)benzylamine, N-(2-(5-Methyl-1H-indol-3-yl)ethyl)-3-(3-fluoropropoxy)benzylamine, N-(2-(5-Methoxy-1H-indol-3-yl)ethyl)-3-(4,4,4-trifluorobutoxy) benzylamine, N-(2-(5-Chloro-1H-indol-3-yl)ethyl)-3-(2-fluoroethoxy)benzylamine, N-(2-(4,7-Difluoro-1H-indol-3-yl)ethyl)-3-(2,2,2-trifluoroethoxy)benzylamine, N-(2-(4,5,6,7-Tetrafluoro-1H-indol-3-yl)ethyl)-3-(2,2,2-trifluoroethoxy)benzylamine, N-(2-(6-Bromo-1H-indol-3-yl)ethyl)-3-(2,2,3,3-tetrafluoropropoxy) benzylamine, N-(2-(6-Bromo-1H-indol-3-yl)ethyl)-3-(2,2,2-trifluoroethyl)benzylamine, N-(2-(6-Methanesulfonyl-1H-indol-3-yl)ethyl)-3-(2,2,3,3-tetrafluoropropoxy) benzylamine, N-(2-(6-Metanesulfonyl-1H-indol-3-yl)ethyl)-3-(2,2,2-trifluoroethoxy)benzylamine, N-(2-(6-Benzenesulfonyl-1H-indol-3-yl)ethyl)-3-(2,2,3,3-tetrafluoropropoxy)benzylamine, N-(2-(6-Benzenesulfonyl-1H-indol-3-yl)ethyl)-3-(2,2,2-trifluoroethoxy) benzylamine, N-(2-(6-Methoxy-1H-indol-3-yl)ethyl)-3-(2,2,3,3-tetrafluoropropoxy)benzylamine, N-(2-(7-Fluoro-1H-indol-3-yl)ethyl)-3-(3,3,3-trifluoropropoxy) benzylamine, N-(2-(6-Bromo-1H-indol-3-yl)ethyl)-3-(2,2,3,3-tetrafluoropropoxy)benzylamine, N-(2-(6-Bromo-1H-indol-3-yl)ethyl)-3-(2,2,3,3,3-pentafluoropropoxy)benzylamine, N-(2-(6-Methoxycarbonyl-1H-indol-3-yl)ethyl)-3-(2,2,2-trifluoroethoxy) benzylamine, N-2-(5-Benzamido-1H-indol-3-yl)-ethyl)-3-(2,2,2-trifluoroethoxy)benzylamine, N-2-(5-Benzamido-1H-indol-3-yl)-ethyl)-3-(2,2,3,3-tetrafluoropropoxy)benzylamine, N-2-(5-Methylsulfonylamino-1H-indol-3-yl)-ethyl)-3-(2,2,2-trifluoroethoxy)benzylamine, N-2-(5-Methylsulfonylamino-1H-indol-3-yl)-ethyl)-3-(2,2,3,3-tetrafluoropropoxy) benzylamine, N-2-(5-Isopropyl-1H-indol-3-yl)-ethyl)-3-(2,2,2-trifluoroethoxy)benzylamine, N-2-(5-Isopropyl-1H-indol-3-yl)-ethyl)-3-(2,2,3,3-tetrafluoropropoxy) benzylamine, N-2-(5-Ethoxy-1H-indol-3-yl)-ethyl)-3-(2,2,2-trifluoroethoxy)benzylamine, N-2-(5-Ethoxy-1H-indol-3-yl)-ethyl)-3-(2,2,3,3-tetrafluoropropoxy)benzylamine, N-2-(5-(2,2,2-Trifluoroethoxy)-1H-indol-3-yl)-ethyl)-3-(2,2,2-trifluoroethoxy)benzylamine, N-2-(5-(2,2,2-Trifluoroethoxy)-1H-indol-3-yl)-

-204-

ethyl)-3-(2,2,3,3-tetrafluoropropoxy) benzylamine, N-2-(5-Butoxy-1H-indol-3-yl)-ethyl)-3-(pyrid-2-yloxy)benzylamine, N-2-(5-Benzenesulfonyl-1H-indol-3-yl)-ethyl)-3-(2,2,3,3-tetrafluoropropoxy)benzylamine, N-2-(5-Benzenesulfonyl-1H-indol-3-yl)-ethyl)-3-(2,2,2-trifluoroethoxy)benzylamine, N-2-(5-Ethoxycarbonyl-1H-indol-3-yl)-ethyl)-3-(2,2,2-trifluoroethoxy)benzylamine, N-2-(5-(N'-Propylamido)-1H-indol-3-yl)-ethyl)-3-(2,2,3,3-tetrafluoropropoxy)benzylamine, N-2-(5-(N'-Propylamido)-1H-indol-3-yl)-ethyl)-3-(2,2,2-trifluoroethoxy)benzylamine, N-2-(5-(N'-Butylamido)-1H-indol-3-yl)-ethyl)-3-(2,2,3,3-tetrafluoropropoxy)benzylamine, N-2-(1H-Indol-3-yl)-ethyl)-3-(2,2,3,3-tetrafluoropropoxy)benzylamine, N-2-(5-Benzyloxy-1H-indol-3-yl)-ethyl)-3-(2,2,2-trifluoroethoxy) benzylamine, N-2-(5-Benzyloxy-1H-indol-3-yl)-ethyl)-3-(2,2,3,3-tetrafluoropropoxy)benzylamine, N-2-(5-Phenoxy-1H-indol-3-yl)-ethyl)-3-(2,2,3,3-tetrafluoropropoxy) benzylamine, N-2-(5-Phenoxy-1H-indol-3-yl)-ethyl)-3-(2,2,2-trifluoroethoxy)benzylamine, N-2-(5-(Pyrid-3-yloxy)-1H-indol-3-yl)-ethyl)-3-(2,2,3,3-tetrafluoropropoxy) benzylamine, N-2-(5-(Pyrid-3-yloxy)-1H-indol-3-yl)-ethyl)-3-(2,2,2-trifluoroethoxy)benzylamine, N-Methyl-N-(2-(6-Chloro-1H-indol-3-yl)ethyl)-3-(2,2-difluoroethoxy)benzylamine, N-2-(4,7-Difluoro-1H-indol-3-yl)-ethyl)-3-(2,2,2-trifluoroethoxy)benzylamine, N-2-(4,5,6,7-Tetrafluoro-1H-indol-3-yl)-ethyl)-3-(2,2,2-trifluoroethoxy)benzylamine, N-2-(4,7-Difluoro-1H-indol-3-yl)-ethyl)-3-(2,2,3,3-tetrafluoropropoxy)benzylamine, N-2-(4,5,6,7-Tetrafluoro-1H-indol-3-yl)-ethyl)-3-(2,2,3,3-tetrafluoropropoxy) benzylamine, N-2-(7-Trifluoromethyl-1H-indol-3-yl)-ethyl)-3-(2,2,2-trifluoroethoxy)benzylamine, N-2-(7-Trifluoromethyl-1H-indol-3-yl)-ethyl)-3-(2,2,3,3-tetrafluoropropoxy) benzylamine, N-2-(7-Nitro-1H-indol-3-yl)-ethyl)-3-(2,2,2-trifluoroethoxy)benzylamine, and N-2-(7-Nitro-1H-indol-3-yl)-ethyl)-3-(2,2,3,3-tetrafluoropropoxy)benzylamine.

25

15. A compound of Claim 1 selected from the group consisting of N-(2-Phenylphenylethyl)-3-(2,2,2-trifluoroethoxy)benzylamine, N-(4-Phenylphenylethyl)-3-(2,2,2-trifluoroethoxy)benzylamine, N-(2-(4-Chlorophenyl)ethyl)-3-(2,2,2-trifluoroethoxy)benzylamine, N-(2-(3-Trifluoromethylphenyl)ethyl)-3-(2,2,2-trifluoroethoxy)benzylamine, N-(2-(3-Chlorophenyl)ethyl)-3-(2,2,3,3,3-pentafluoropropoxy)benzylamine, N-(2-(3-Trifluoromethylphenyl)ethyl)-3-(2,2,3,3,3-pentafluoropropoxy)benzylamine, N-(2-(3-Chlorophenyl)ethyl)-3-(2,2,3,3-tetrafluoropropoxy) benzylamine, N-(2-(3-Trifluoromethylphenyl)ethyl)-3-(2,2,3,3-tetrafluoropropoxy)benzylamine, N-(2-(3-Trifluoromethylphenyl)ethyl)-3-(3,3,3-trifluoropropoxy)benzylamine, N-(2-(3-Chlorophenyl)ethyl)-3-(3,3,3-trifluoropropoxy)benzylamine, N-(2-(3,5-Dimethoxyphenyl)ethyl)-3-(2,2,2-trifluoroethoxy) benzylamine, N-(2-(3-Chlorophenyl)ethyl)-3-(2-

35

- fluoroethoxy)benzylamine, N-(2-(3-Trifluoromethylphenyl)ethyl)-3-(2-fluoroethoxy) benzylamine, N-(2-(3,5-Dimethoxyphenyl)ethyl)-3-(2-fluoroethoxy)benzylamine, N-(2-(3-Trifluoromethyl-4-fluorophenyl)ethyl)-3-(3,3,3-trifluoropropoxy) benzylamine, N-(2-(3-Trifluoromethyl-4-fluorophenyl)ethyl)-3-(2,2-difluoroethoxy)benzylamine, N-(2-(3-Trifluoromethyl-4-fluorophenyl)ethyl)-3-(2,2,3,3,3-pentafluoropropoxy) benzylamine, N-(2-(2-(3-Trifluoromethyl-4-fluorophenyl)ethyl)-3-(2,2,2-trifluoroethoxy)benzylamine, and N-(2-(3-Trifluoromethyl-4-fluorophenyl)ethyl)-3-(3-fluoropropoxy)benzylamine.
16. A compound of Claim 1 selected from the group consisting of N-(2-(5-Methoxy-1H-indol-3-yl)ethyl)-3-phenoxybenzylamine, N-(2-(1H-Indol-3-yl)ethyl)-3-phenoxybenzylamine, N-(2-(5-Methoxy-1H-indol-3-yl)ethyl)-3-(3-fluorophenoxy)benzylamine, N-(2-(1H-Indol-3-yl)ethyl)-3-(3-fluorophenoxy)benzylamine, N-(2-(5-Methoxy-1H-indol-3-yl)ethyl)-3-(2-fluorophenoxy)benzylamine, N-(2-(1H-Indol-3-yl)ethyl)-3-(2-fluorophenoxy)benzylamine, N-(2-(5-Methoxy-1H-indol-3-yl)ethyl)-3-(4-fluorophenoxy)benzylamine, N-(2-(1H-Indol-3-yl)ethyl)-3-(4-fluorophenoxy)benzylamine, N-(2-(5-Hydroxy-1H-indol-3-yl)ethyl)-3-phenoxybenzylamine, N-(2-(5-Phenoxy-1H-indol-3-yl)ethyl)-3-phenoxybenzylamine, N-(2-(5-p-Tolyloxy-1H-indol-3-yl)ethyl)-3-phenoxybenzylamine, N-(2-(5-o-Tolyloxy-1H-indol-3-yl)ethyl)-3-phenoxy benzylamine, N-(2-(5-m-Tolyloxy-1H-indol-3-yl)ethyl)-3-phenoxybenzylamine, N-(2-(6-Fluoro-1H-indol-3-yl)ethyl)-2-fluoro-3-phenoxybenzylamine, N-(2-(6-Fluoro-1H-indol-3-yl)ethyl)-6-fluoro-3-phenoxybenzylamine, N-Methyl-N-(2-(5-m-tolyloxy-1H-indol-3-yl)ethyl)-3-phenoxybenzylamine, N-(2-(5-Nitro-1H-indol-3-yl)ethyl)-3-phenoxybenzylamine, N-(2-(5-Amino-1H-indol-3-yl)ethyl)-3-phenoxybenzylamine, N-(2-(5-Propoxy-1H-indol-3-yl)ethyl)-3-phenoxybenzylamine, N-(2-(5-Ethoxycarbonyl-1H-indol-3-yl)ethyl)-3-phenoxy benzylamine, N-(2-(5-Phenyl-1H-indol-3-yl)ethyl)-3-phenoxybenzylamine, N-(2-(5-(4-Fluorophenyl)-1H-indol-3-yl)ethyl)-3-phenoxybenzylamine, N-(2-(6-Phenyl-1H-indol-3-yl)ethyl)-3-phenoxybenzylamine, N-(2-(6-Trifluoromethyl-1H-indol-3-yl)ethyl)-3-phenoxybenzylamine, N-(2-(6-Fluoro-1H-indol-3-yl)ethyl)-3-phenoxybenzylamine N-(2-(6-Chloro-5-methoxy-1H-indol-3-yl)ethyl)-3-phenoxybenzylamine, N-(2-(4-Chloro-5-methoxy-1H-indol-3-yl)ethyl)-3-phenoxy benzylamine, N-(2-(6-Chloro-1H-indol-3-yl)ethyl)-3-phenoxybenzylamine, N-(2-(4,6-Dichloro-5-methoxy-1H-indol-3-yl)ethyl)-3-phenoxybenzylamine, N-(2-(6-Methoxy-1H-indol-3-yl)ethyl)-3-phenoxybenzylamine, N-(2-(4-Chloro-1H-indol-3-yl)ethyl)-3-phenoxybenzylamine, N-(2-(4-Methoxy-1H-indol-3-yl)ethyl)-3-phenoxybenzylamine, N-(2-(5-Methoxy-2-methyl-1H-indol-3-yl)ethyl)-3-phenoxybenzylamine, N-(2-(7-Methoxy-1H-indol-3-yl)ethyl)-3-phenoxybenzylamin, N-(2-(5-Methoxy-4,6-difluoro-1-methyl-1H-indol-3-yl)ethyl)-3-(2,2,2-trifluoroethoxy)benzylamine, N-(2-(5-Methyl-1H-indol-3-

-206-

yl)ethyl)-3-phenoxybenzylamine, N-(2-(5-Chloro-1H-indol-3-yl)ethyl)-3-phenoxybenzylamine, N-(2-(5-(4-Fluorophenyl)-1H-indol-3-yl)ethyl)-3-phenoxybenzylamine, N-(2-(5-Methanesulfonyl-1H-indol-3-yl)-ethyl)-3-phenoxybenzylamine, N-(2-(5-Cyano-1H-indol-3-yl)-ethyl)-3-phenoxybenzylamine, N-(2-(5-Methoxycarbonyl-1H-indol-3-yl)-ethyl)-3-phenoxybenzylamine, N-(2-(5-Amido-1H-indol-3-yl)-ethyl)-3-phenoxybenzylamine, N-2-(5-Nitro-1H-indol-3-yl)-ethyl)-3-phenoxybenzylamine, N-2-(5-Butoxy-1H-indol-3-yl)-ethyl)-3-phenoxybenzylamine, N-2-(5-Benzamido-1H-indol-3-yl)-ethyl)-3-phenoxybenzylamine, N-2-(5-Methylsulfonylamino-1H-indol-3-yl)-ethyl)-3-phenoxybenzylamine, N-2-(5-Isopropyl-1H-indol-3-yl)-ethyl)-3-phenoxybenzylamine, N-2-(5-Ethoxy-1H-indol-3-yl)-ethyl)-3-phenoxybenzylamine, N-2-(5-(2,2,2-Trifluoroethoxy)-1H-indol-3-yl)-ethyl)-3-phenoxybenzylamine, N-2-(5-Benzenesulfonyl-1H-indol-3-yl)-ethyl)-3-phenoxybenzylamine, N-2-(5-(N'-Propylamido)-1H-indol-3-yl)-ethyl)-3-phenoxybenzylamine, N-2-(5-(N'-Butylamido)-1H-indol-3-yl)-ethyl)-3-phenoxybenzylamine, N-2-(5-Phenoxy-1H-indol-3-yl)-ethyl)-3-phenoxybenzylamine, N-Methyl-N-2-(5-Phenoxy-1H-indol-3-yl)-ethyl)-3-phenoxybenzylamine, N-(2-(6-Fluoro-1H-indol-3-yl)ethyl)-4-fluoro-3-phenoxybenzylamine, N-(2-(6-Fluoro-1H-indol-3-yl)ethyl)-3-phenoxybenzylamine, N-(2-(5-Benzoyloxy-1H-indol-3-yl)-ethyl)-3-phenoxybenzylamine, and N-(2-(5-Benzoyloxy-1H-indol-3-yl)-ethyl)-N-methyl-3-phenoxybenzylamine.

17. A compound of Claim 1 selected from the group consisting of N-(2-(3-Chlorophenyl)ethyl)-3-phenoxybenzylamine, N-(2-(3-Trifluoromethylphenyl)ethyl)-3-phenoxybenzylamine, N-(2-(4-Methoxyphenyl)ethyl)-3-phenoxybenzylamine, N-(2-(3,4-Dimethoxyphenyl)ethyl)-3-phenoxybenzylamine, N-(2-(3-Methoxyphenyl)ethyl)-3-phenoxybenzylamine, N-(2-(3,4-Dichlorophenyl)ethyl)-3-phenoxybenzylamine, N-(2-(3-Chlorophenyl)ethyl)-3-(3-trifluoromethylphenoxy)benzylamine, N-(2-(3-Chlorophenyl)ethyl)-3-(4-t-butylphenoxy)benzylamine, N-(2-(3-Chlorophenyl)ethyl)-3-(4-chlorophenoxy)benzylamine, N-(2-(3-Chlorophenyl)ethyl)-3-(4-methoxyphenoxy)benzylamine, N-(2-(3-Chlorophenyl)ethyl)-3-(4-methylphenoxy)benzylamine, N-(2-(3-Chlorophenyl)ethyl)-3-(3,5-dichlorophenoxy)benzylamine, N-(2-(3-Chlorophenyl)ethyl)-3-(3,4-dichlorophenoxy)benzylamine, N-(2-Phenylethyl)-3-phenoxybenzylamine, N-(2-(4-Chlorophenyl)ethyl)-3-phenoxybenzylamine, N-(2-(3-Chlorophenyl)ethyl)-3-(4-fluorophenoxy)benzylamine, N-(2-(3-Trifluoromethylphenyl)ethyl)-3-(4-fluorophenoxy)benzylamine, N-(2-(3-Chlorophenyl)ethyl)-3-(2-fluorophenoxy)benzylamine, N-(2-(3-Trifluoromethylphenyl)ethyl)-3-(2-fluorophenoxy)benzylamine, N-(2-(3-

-207-

- Chlorophenyl)ethyl)-3-(3-fluorophenoxy)benzylamine, N-(2-(3-Trifluoromethylphenyl)ethyl)-3-(3-fluorophenoxy)benzylamine, N-(2-(2-Chlorophenyl)ethyl)-3-phenyloxybenzylamine, N-(2-(3,4-Dimethoxyphenyl)ethyl)-3-phenoxybenzylamine, N-(2-(3-Chlorophenyl)ethyl)-N-methyl-3-phenoxybenzylamine, N-(2-(3-Chlorophenyl)ethyl)-N-ethyl-3-phenoxybenzylamine, N-(2-(2-Fluorophenyl)ethyl)-3-phenoxybenzylamine, N-(2-(3-Fluorophenyl)ethyl)-3-phenoxybenzylamine, N-(2-(4-Chlorophenyl)ethyl)-3-phenoxybenzylamine, N-(2-(4-Hydroxyphenyl)ethyl)-3-phenoxybenzylamine, N-(2-(2-Methoxyphenyl)ethyl)-3-phenoxybenzylamine, N-(2-(3-Bromo-3-methoxyphenyl)ethyl)-3-phenoxybenzylamine, N-(2-(4-Fluorophenyl)ethyl)-3-phenoxybenzylamine, N-(2-(2-Chlorophenyl)ethyl)-3-phenoxybenzylamine, N-(2-(4-Bromophenyl)ethyl)-3-phenoxybenzylamine, N-(2-(4-Methylphenyl)ethyl)-3-phenoxybenzylamine, N-(2-(3-Methoxyphenyl)ethyl)-3-phenoxybenzylamine, N-(2-(4-Methoxyphenyl)ethyl)-3-phenoxybenzylamine, N-(2-(2-Ethoxyphenyl)ethyl)-3-phenoxybenzylamine, N-(2-(4-Ethoxyphenyl)ethyl)-3-phenoxybenzylamine, N-(2-(4-Phenoxyphenyl)ethyl)-3-phenoxybenzylamine, N-(2-(4-Sulfonamidophenyl)ethyl)-3-phenoxybenzylamine, N-(2-(3,4-Dichlorophenyl)ethyl)-3-phenoxybenzylamine, N-(2-(2,5-Dichlorophenyl)ethyl)-3-phenoxybenzylamine, N-(2-(2,6-Dichlorophenyl)ethyl)-3-phenoxybenzylamine, N-(2-(2,5-Dimethoxyphenyl)ethyl)-3-phenoxybenzylamine, N-(2-(2,3-Dimethoxyphenyl)ethyl)-3-phenoxybenzylamine, N-(2-(3,5-Dimethoxyphenyl)ethyl)-3-phenoxybenzylamine, and N-(2-(3-Ethoxy-4-methoxyphenyl)ethyl)-3-phenoxybenzylamine.
18. A compound of Claim 1 selected from the group consisting of N-(2-Pyrid-4-ylethyl)-3-phenoxybenzylamine,
- N-(2-Pyrid-3-ylethyl)-3-phenoxybenzylamine, N-(2-(Pyrid-2-ylethyl)-3-phenoxybenzylamine, N-(2-Imidazol-4-ylethyl)-3-phenoxybenzylamine, N-(2-Naphth-2-ylethyl)-3-phenoxybenzylamine, N-(2-Naphth-1-ylethyl)-3-phenoxybenzylamine, and N-(2-(Thien-2-ylethyl)-3-phenoxybenzylamine.
19. A compound of Claim 1 selected from the group consisting of N-(2-(1H-Indol-3-yl)ethyl)-3-thiophenoxybenzylamine, N-(2-(1H-Indol-3-yl)ethyl)-3-phenylsulfonylbenzylamine, N-(2-(5-Methoxy-1H-indol-3-yl)ethyl)-3-thiophenoxybenzylamine, N-(2-(5-Methoxy-1H-indol-3-yl)ethyl)-3-phenylsulfonylbenzylamine, N-(2-(1H-Indol-3-yl)ethyl)-3-(4-methylthiophenoxy)benzylamine, N-(2-(1H-Indol-3-yl)ethyl)-3-(4-methylphenylsulfonyl)benzylamine, N-(2-(5-Methoxy-1H-indol-3-yl)ethyl)-3-(α -fluorobenzyl)benzylamine, N-(2-(1H-Indol-3-yl)ethyl)-3-(α -fluorobenzyl)benzylamine, N-(2-(5-Methoxy-1H-indol-3-

-208-

- yl)ethyl)-3-(benzyl)benzylamine, N-(2-(1H-Indol-3-yl)ethyl)-3-(benzyl)benzylamine, N-(2-(5-Methoxy-1H-indol-3-yl)ethyl)-3-(α -hydroxybenzyl)benzylamine, N-(2-(5-Methoxy-1H-indol-3-yl)ethyl)-3-(N-phenylamino)benzylamine, N-(2-(1H-Indol-3-yl)ethyl)-3-(N-phenylamino)benzylamine, N-(2-(5-Methoxy-1H-indol-3-yl)ethyl)-3-(N-
- 5 benzylamino)benzylamine, N-(2-(1H-Indol-3-yl)ethyl)-3-(N-benzylamino)benzylamine, N-(2-(5-Methoxy-1H-indol-3-yl)ethyl)-3-hydroxybenzylamine, N-(2-(1H-Indol-3-yl)ethyl)-3-hydroxybenzylamine, N-(2-(5-Methoxy-1H-indol-3-yl)ethyl)-3-naphth-1-yloxybenzylamine, N-(2-(1H-Indol-3-yl)ethyl)-3-naphth-1-yloxybenzylamine, N-(2-(5-Methoxy-1H-indol-3-yl)ethyl)-3-naphth-2-
- 10 yloxybenzylamine, N-(2-(1H-Indol-3-yl)ethyl)-3-naphth-2-yloxybenzylamine, N-(2-(5-Methoxy-1H-indol-3-yl)ethyl)-3-benzyloxybenzylamine, N-(2-(1H-Indol-3-yl)ethyl)-3-benzyloxybenzylamine, N-(2-(5-Methoxy-1H-indol-3-yl)ethyl)-3-pyrimid-5-yloxybenzylamine, N-(2-(1H-Indol-3-yl)ethyl)-3-pyrimid-5-yloxybenzylamine, N-(2-(5-Methoxy-1H-indol-3-yl)ethyl)-3-pyrid-4-yloxybenzylamine, N-(2-(1H-Indol-3-yl)ethyl)-3-pyrid-4-yloxybenzylamine, N-(2-(6-Chloro-1H-indol-3-yl)ethyl)-3-pyrid-4-
- 15 yloxybenzylamine, N-(2-(5-Methoxy-1H-indol-3-yl)ethyl)-3-pyrid-3-yloxybenzylamine, N-(2-(1H-Indol-3-yl)ethyl)-3-pyrid-3-yloxybenzylamine, N-(2-(5-Fluoro-1H-indol-3-yl)ethyl)-3-pyrid-3-yloxybenzylamine, N-(2-(6-Chloro-1H-indol-3-yl)ethyl)-3-pyrid-3-yloxybenzylamine, N-(2-(5-Methoxy-1H-indol-3-yl)ethyl)-3-pyrid-2-yloxybenzylamine, N-(2-(1H-Indol-3-yl)ethyl)-3-pyrid-2-yloxybenzylamine, N-(2-(6-Chloro-1H-indol-3-yl)ethyl)-3-pyrid-2-yloxybenzylamine, N-(2-(5-Methoxy-1H-indol-3-yl)ethyl)-3-thiazol-2-yloxybenzylamine, N-(2-(1H-Indol-3-yl)ethyl)-3-thiazol-2-yloxybenzylamine, N-(2-(5-Methoxy-1H-indol-3-yl)ethyl)-3-(2,6-difluororsulfonyloxy)benzylamine, N-(2-(1H-Indol-3-yl)ethyl)-3-(2,6-difluororsulfonyloxy)benzylamine, N-(2-
- 25 (5-Methoxy-1H-indol-3-yl)ethyl)-3-(pyrid-2-ylamino)benzylamine, N-(2-(1H-Indol-3-yl)ethyl)-3-(pyrid-2-ylamino)benzylamine, N-(2-(6-Chloro-1H-indol-3-yl)ethyl)-3-(pyrid-2-ylamino)benzylamine, N-(2-(5-Methoxy-1H-indol-3-yl)ethyl)-3-(pyrid-3-ylamino)benzylamine, N-(2-(1H-Indol-3-yl)ethyl)-3-(pyrid-3-ylamino)benzylamine, N-(2-(6-Chloro-1H-indol-3-yl)ethyl)-3-(pyrid-3-ylamino)benzylamine, N-(2-(5-Methoxy-1H-indol-3-yl)ethyl)-3-(pyrid-4-ylamino)benzylamine, N-(2-(1H-Indol-3-yl)ethyl)-3-(pyrid-4-ylamino)benzylamine, N-(2-(6-Chloro-1H-indol-3-yl)ethyl)-3-(pyrid-4-ylamino)benzylamine, N-(2-(5-Methoxy-1H-indol-3-yl)ethyl)-3-benzoylbenzylamine, N-(2-(1H-Indol-3-yl)ethyl)-3-benzoylbenzylamine, N-(2-(6-Fluoro-1H-indol-3-yl)ethyl)-3-(pyrid-4-yloxy)benzylamine, N-(2-(6-Fluoro-1H-indol-3-yl)ethyl)-3-(pyrid-3-yloxy)
- 30 benzylamine, N-(2-(3-Chlorophenyl)ethyl)-3-thiophenoxybenzylamine, N-(2-(3-Chlorophenyl)ethyl)-3-sulfonylphenylbenzylamine, N-(2-Phenylethyl)-3-thiophenoxybenzylamine, N-(2-Phenylethyl)-3-sulfonylphenylbenzylamine, N-(2-(3-

-209-

Chlorophenyl)ethyl)-3-(4-methylthiophenoxy)benzylamine, N-(2-(3-Chlorophenyl)ethyl)-3-sulfonyl-4-methylphenylbenzylamine, N-(2-(3-Chlorophenyl)ethyl)-3-(N-benzylamino)benzylamine, N-(2-(3-Chlorophenyl)ethyl)-3-(N-phenylamino)benzylamine, N-(2-(3-Chlorophenyl)ethyl)-3-(α -hydroxybenzyl)benzylamine, N-(2-(3-Chlorophenyl)ethyl)-3-benzylbenzylamine, N-(2-(3-Chlorophenyl)ethyl)-3-(α -fluorobenzyl)benzylamine, N-(2-(3-Chlorophenyl)ethyl)-3-naphth-2-yloxybenzylamine, N-(2-(3-Trifluoromethylphenyl)ethyl)-3-naphth-2-yloxybenzylamine, N-(2-(3-Chlorophenyl)ethyl)-3-naphth-1-yloxybenzylamine, N-(2-(3-Trifluoromethylphenyl)ethyl)-3-naphth-1-yloxybenzylamine, N-(2-(3-Chlorophenyl)ethyl)-3-hydroxybenzylamine, N-(2-(3-Trifluoromethylphenyl)ethyl)-3-hydroxybenzylamine, N-(2-(3-Trifluoromethylphenyl)ethyl)-3-benzyloxybenzylamine, N-(2-(3-Chlorophenyl)ethyl)-3-(2,4-difluoro phenylsulfonyloxy)benzylamine, N-(2-(3-Trifluoromethylphenyl)ethyl)-3-(2,4-difluoro phenylsulfonyloxy)benzylamine, N-(2-(3-Chlorophenyl)ethyl)-3-thiazol-2-yloxybenzylamine, N-(2-(3-Trifluoromethylphenyl)ethyl)-3-thiazol-2-yloxy benzylamine, N-(2-(3-Chlorophenyl)ethyl)-3-pyrid-3-yloxybenzylamine, N-(2-(3-Trifluoromethylphenyl)ethyl)-3-pyrid-3-yloxybenzylamine, N-(2-(3-Methoxyphenyl)ethyl)-3-pyrid-2-yloxybenzylamine, N-(2-(3-Trifluoromethylphenyl)ethyl)-3-pyrid-4-yloxybenzylamine, N-(2-(3-Chlorophenyl)ethyl)-3-pyrimid-5-yloxybenzylamine, N-(2-(3-Trifluoromethylphenyl)ethyl)-3-pyrimid-5-yloxybenzylamine, N-(2-(3-Trifluoromethylphenyl)ethyl)-3-pyrid-2-yloxybenzylamine, N-(2-(3-Chlorophenyl)ethyl)-3-pyrid-3-ylaminobenzylamine, N-(2-(3-Trifluoromethylphenyl)ethyl)-3-pyrid-3-ylaminobenzylamine, N-(2-(3-Chlorophenyl)ethyl)-3-pyrid-4-ylaminobenzylamine, N-(2-(3-Trifluoromethylphenyl)ethyl)-3-pyrid-4-ylaminobenzylamine, N-(2-(3-Chlorophenyl)ethyl)-3-pyrid-2-ylaminobenzylamine, N-(2-(3-Trifluoromethylphenyl)ethyl)-3-pyrid-2-ylaminobenzylamine, N-(2-(3-Chlorophenyl)ethyl)-3-benzyloxybenzylamine, N-(2-(3-Trifluoromethylphenyl)ethyl)-3-(phenylamino)benzylamine, N-(2-(7-Fluoro-1H-indol-3-yl)ethyl)-3-(pyrid-4-yloxy)benzylamine, N-(2-(7-Fluoro-1H-indol-3-yl)ethyl)-3-(pyrid-3-yloxy)benzylamine, N-(2-(3-Chlorophenyl)ethyl)-3-(pyrid-2-ylmethoxy)benzylamine, N-(2-(3-Chlorophenyl)ethyl)-3-(pyrid-3-ylmethoxy)benzylamine, N-(2-(3-Chlorophenyl)ethyl)-3-(pyrid-4-ylmethoxy) benzylamine, N-2-(5-Butoxy-1H-indol-3-yl)-ethyl)-3-(pyrid-2-yloxy)benzylamine, and N-(2-(6,7-Difluoro-1H-indol-3-yl)-ethyl)-3-(pyridin-4-yloxy)benzylamine.

35

20. A compound of Claim 1 selected from the group consisting of N-(2-(3-Chlorophenyl)ethyl)-3-ethoxybenzylamine, N-(2-(3-Chlorophenyl)ethyl)-3-

- propoxybenzylamine, N-(2-(3-Trifluoromethylphenyl)ethyl)-3-propoxybenzylamine, N-(2-(3-Trifluoromethylphenyl)ethyl)-3-ethoxybenzylamine, N-(2-(5-Chloro-1H-indol-3-yl)ethyl)-3-ethoxybenzylamine, N-(2-(5-Chloro-1H-indol-3-yl)ethyl)-3-propoxybenzylamine, N-(2-(5-Fluoro-1H-indol-3-yl)ethyl)-3-propoxybenzylamine, N-(2-(5-Fluoro-1H-indol-3-yl)ethyl)-3-ethoxybenzylamine, N-(2-(5-Trifluoromethyl-1H-indol-3-yl)ethyl)-3-ethoxybenzylamine, N-(2-(5-Trifluoromethyl-1H-indol-3-yl)ethyl)-3-propoxybenzylamine, N-(2-(5-Methoxy-1H-indol-3-yl)ethyl)-3-ethoxybenzylamine, N-(2-(5-Methoxy-1H-indol-3-yl)ethyl)-3-propoxybenzylamine, N-(2-(4-Chloro-1H-indol-3-yl)ethyl)-3-propoxybenzylamine, N-(2-(4-Methoxy-1H-indol-3-yl)ethyl)-3-propoxybenzylamine, N-(2-(5-Methoxy-2-methyl-1H-indol-3-yl)ethyl)-3-propoxybenzylamine, N-(2-(7-Methoxy-1H-indol-3-yl)ethyl)-3-propoxybenzylamine, N-(2-(6-Chloro-1H-indol-3-yl)ethyl)-3-propoxybenzylamine, N-(2-(6-Fluoro-1-methyl-1H-indol-3-yl)ethyl)-N-methyl-3-propoxybenzylamine, N-(2-(3-Trifluoromethyl-4-fluorophenyl)ethyl)-3-propoxy-4-methylbenzylamine, N-(2-(5-Fluoro-6-chloro-1H-indol-3-yl)ethyl)-3-propoxy-4-methylbenzylamine, N-(2-(6-Fluoro-1-methyl-1H-indol-3-yl)ethyl)-3-propoxy-4-methylbenzylamine, N-(2-(3-Chlorophenyl)ethyl)-3-propoxybenzylamine, N-(2-(3-Chlorophenyl)ethyl)-3-butoxybenzylamine, N-(2-(3-Chlorophenyl)ethyl)-3-hexoxybenzylamine, N-(2-(2-Fluorophenyl)ethyl)-3-propoxybenzylamine, N-(2-(3-Fluorophenyl)ethyl)-3-propoxybenzylamine, N-(2-(4-Fluorophenyl)ethyl)-3-propoxybenzylamine, N-(2-(2-Chlorophenyl)ethyl)-3-propoxybenzylamine, N-(2-(4-Chlorophenyl)ethyl)-3-propoxybenzylamine, N-(2-(4-Bromophenyl)ethyl)-3-propoxybenzylamine, N-(2-(4-Methylphenyl)ethyl)-3-propoxybenzylamine, N-(2-(4-Hydroxyphenyl)ethyl)-3-propoxybenzylamine, N-(2-(2-Methoxyphenyl)ethyl)-3-propoxybenzylamine, N-(2-(3-Methoxyphenyl)ethyl)-3-propoxybenzylamine, N-(2-(4-Methoxyphenyl)ethyl)-3-propoxybenzylamine, N-(2-(3-Ethoxyphenyl)ethyl)-3-propoxybenzylamine, N-(2-(4-Ethoxyphenyl)ethyl)-3-propoxybenzylamine, N-(2-(4-Phenoxyphenyl)ethyl)-3-propoxybenzylamine, N-(2-(4-Sulfonamidophenyl)ethyl)-3-propoxybenzylamine, N-(2-(3,4-Dichlorophenyl)ethyl)-3-propoxybenzylamine, N-(2-(2,5-Dichlorophenyl)ethyl)-3-propoxybenzylamine, N-(2-(2,6-Dichlorophenyl)ethyl)-3-propoxybenzylamine, N-(2-(3,4-Dimethoxyphenyl)ethyl)-3-propoxybenzylamine, N-(2-(2,5-Dimethoxyphenyl)ethyl)-3-propoxybenzylamine, N-(2-(2,3-Dimethoxyphenyl)ethyl)-3-propoxybenzylamine, N-(2-(3,5-Dimethoxyphenyl)ethyl)-3-propoxybenzylamine, N-(2-(3-Bromo-4-methoxyphenyl)ethyl)-3-propoxybenzylamine, N-(2-(3-Methoxy-4-ethoxyphenyl)ethyl)-3-propoxybenzylamine, N-(2-(3-Ethoxy-4-methoxyphenyl)ethyl)-3-propoxybenzylamine, N-(2-(Pyrid-2-yl)ethyl)-3-propoxybenzylamine, N-(2-(Pyrid-3-yl)ethyl)-3-propoxybenzylamine, N-(2-(Pyrid-4-yl)ethyl)-3-propoxybenzylamine, N-(2-(7-

-211-

Methyl-1H-indol-3-yl)ethyl)-3-propoxybenzylamine, N-(2-(6-Methoxy-1H-indol-3-yl)ethyl)-3-propoxybenzylamine, N-(2-(Thien-3-yl)ethyl)-3-propoxybenzylamine, N-(2-(5-Methyl-1H-indol-3-yl)ethyl)-3-propoxybenzylamine, N-(2-(5-Methoxy-1H-indol-3-yl)ethyl)-3-propoxybenzylamine, N-(2-(3-Bromophenyl)ethyl)-3-propoxybenzylamine, N-
5 (2-(3-Methoxycarbonylphenyl)ethyl)-3-propoxybenzylamine, and N-(2-(3-(4-Fluorophenyl)phenyl)-ethyl)-3-propoxybenzylamine.

21. A pharmaceutical composition comprising a compound of Claim 1 and a pharmaceutically acceptable diluent.
10
22. A method of treating disorders associated with the 5-HT₆ receptor, comprising: administering to a patient in need thereof an effective amount of a compound of Claim 1.
23. A method according to Claim 22 wherein the disorder is selected from the group
15 consisting of cognitive disorders, schizophrenia, anxiety, and Alzheimer's disease, comprising: administering to a patient in need thereof an effective amount of a compound of Claim 1.
24. A method for treating cognitive disorders comprising: administering to a patient in
20 need thereof an effective amount of a compound of Claim 1.
25. A method of treating memory disorders, comprising: administering to a patient in need thereof an effective amount of a compound of Claim 1.
- 25 26. A method of treating psychosis, comprising: administering to a patient in need thereof an effective amount of a compound of Claim 1.
27. A method of treating schizophrenia, comprising: administering to a patient in need thereof an effective amount of a compound of Claim 1.
30
28. A method of treating anxiety, comprising: administering to a patient in need thereof an effective amount of a compound of Claim 1.
28. The use of a compound of Claim 1 as a pharmaceutical.
35
29. The use of a compound of Claim 1 for the manufacture of a medicament for treating disorders associated with the 5-HT₆ receptor.

-212-

30. The use of a compound of Claim 1 for the manufacture of a medicament for treating cognitive disorders.

5 31. The use of a compound of Claim 1 for the manufacture of a medicament for treating memory disorders.


32. The use of a compound of Claim 1 for the manufacture of a medicament for treating psychosis.

10

33. The use of a compound of Claim 1 for the manufacture of a medicament for treating schizophrenia.

15 34. The use of a compound of Claim 1 for the manufacture of a medicament for treating anxiety.

35. A compound of the formula



formula II

20 wherein

Y is selected from the group consisting of O, NH, and NR₉, wherein R₉ is selected from the group consisting of C₁-C₄ alkyl, and optionally substituted phenyl;


R₅ and R₆ are hydrogen or taken together with the atoms to which they are attached form a benzo ring, provided that R₅ and R₆ are hydrogen when Y is NR₉;

25 R₇ is selected from the group consisting of optionally substituted phenyl, optionally substituted naphthyl, optionally substituted 5 to 6 membered monocyclic aromatic heterocycle having one heteroatom selected from the group consisting of nitrogen, oxygen, and sulfur and which 5 to 6 membered monocyclic aromatic heterocycle is optionally benzofused;

30 R₈ is selected from the group consisting of hydrogen and C₁-C₃ alkyl; and pharmaceutically acceptable salts thereof.

36. A compound according to Claim 31 wherein the moiety

-213-



is attached by either the 4-position or the 6-position.

37. A compound according to Claim 36 wherein Y is O.
- 5 38. A compound according to Claim 36 wherein Y is NH.
39. A compound according to Claim 36 wherein Y is NR₉.
- 10 40. A compound according to Claim 36 wherein R₅ and R₆ are taken together with the atoms to which they are attached for a phenyl ring.
41. A compound according to Claim 36 wherein R₅ and R₆ are hydrogen.
- 15 42. A compound according to any one of Claims 36 to 41 wherein R₇ is selected from the group consisting of optionally substituted phenyl and optionally substituted 5 to 6 membered monocyclic aromatic heterocycle having one heteroatom selected from the group consisting of nitrogen, oxygen, and sulfur and which 5 to 6 membered monocyclic aromatic heterocycle is optionally benzofused.
- 20 43. A compound according to Claim 42 wherein the optionally benzofused 5 to 6 membered monocyclic aromatic heterocycle is optionally substituted indol-3-yl.
44. A pharmaceutical composition comprising a compound of Claim 35 and a
- 25 pharmaceutically acceptable diluent.
45. A method of treating disorders associated with the 5-HT₆ receptor, comprising: administering to a patient in need thereof an effective amount of a compound of Claim 35.
- 30 46. A method according to Claim 45 wherein the disorder is selected from the group consisting of cognitive disorders, schizophrenia, anxiety, and Alzheimer's disease.
47. The use of a compound of Claim 35 for the manufacture of a medicament.

-214-

48. The use of a compound of Claim 35 for the manufacture of a medicament for treating disorders associated with the 5-HT₆ receptor.

49. A use according to Claim 48 wherein the disorder is selected from the group
5 consisting of cognitive disorders, schizophrenia, anxiety, and Alzheimer's disease.

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
10 October 2002 (10.10.2002)

PCT

(10) International Publication Number
WO 02/078693 A3

(51) International Patent Classification⁷: A61K 31/4045, C07D 209/16, A61P 25/18, 25/28, C07D 403/12, 401/12, A61K 31/4406, 31/4409, 31/506, 31/4402, C07C 217/54, 323/01, 317/14, C07D 213/64, 277/34, 239/34, 233/24, 333/20, 307/91, 209/14

(21) International Application Number: PCT/US02/05115

(22) International Filing Date: 15 March 2002 (15.03.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/279,928 29 March 2001 (29.03.2001) US
60/329,449 15 October 2001 (15.10.2001) US

(71) Applicant (for all designated States except US): ELI LILLY AND COMPANY [US/US]; Lilly Corporate Center, Indianapolis, IN 46285 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): CHEN, Zhaogen [CN/US]; 9972 Parkshore Drive, Noblesville, IN 46060 (US). COHEN, Michael, Philip [US/US]; 8141 Bowline Court, Indianapolis, IN 46236 (US). FISHER, Matthew, Joseph [US/US]; 10650 Quail Ridge Court, Mooresville, IN 46158 (US). GIETHLEN, Bruno [FR/FR], Prestwick Chemical, Inc., Bat. Tycho Brahe, Rue Tobias Stimmer, F-67400 Illkirch-Graffenstaden (FR). GILLIG, James, Ronald [US/US]; 3634 Toronto Court, Indianapolis, IN 46268 (US). MCCOWAN, Jefferson, Ray [US/US], 2653 Crescent Hill Lane, Indianapolis, IN 46228 (US). MILLER, Shawn, Christopher [US/US]; 7359 Hardin Oak Drive, Noblesville, IN 46060 (US). SCHAUS, John, Mehnert [US/US]; 135 Raintree Drive, Zionsville, IN 46077 (US).

(74) Agents: DAWALT, Elizabeth, A. et al.; Eli Lilly and Company, Lilly Corporate Center, Indianapolis, IN 46285 (US).

(81) Designated States (national): AE, AG, AL, AM, AT (utility model), AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA,

CH, CN, CO, CR, CU, CZ (utility model), CZ, DE (utility model), DE, DK (utility model), DK, DM, DZ, EC, EE (utility model), EE, ES, FI (utility model), FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK (utility model), SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.



(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

- as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii)) for the following designations AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZM, ZW, ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG)
- as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii)) for the following designations AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZM, ZW, ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),

[Continued on next page]

(54) Title: N-(2-ARYLETHYL)BENZYLAMINES AS ANTAGONISTS OF THE 5-HT₆ RECEPTOR



(57) Abstract: The present invention provides compounds of formula (I), which are antagonists of the 5-HT₆ receptor.



European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG)

(88) Date of publication of the international search report:
5 December 2002

Published:

— *with international search report*

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 02/05115

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A61K31/4045 C07D209/16 A61P25/18 A61P25/28 C07D403/12
 C07D401/12 A61K31/4406 A61K31/4409 A61K31/506 A61K31/4402
 C07C217/54 C07C323/01 C07C317/14 C07D213/64 C07D277/34

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D C07C

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, BEILSTEIN Data, PAJ, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 00 34242 A (GLENNON RICHARD A ;ROTH BRYAN L (US); UNIV VIRGINIA COMMONWEALTH () 15 June 2000 (2000-06-15) claim 1	1-49
A	US 6 187 805 B1 (MC ALLISTER GEORGE ET AL) 13 February 2001 (2001-02-13) claim 1	1-49

	---/---	



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents :

A document defining the general state of the art which is not considered to be of particular relevance

E earlier document but published on or after the international filing date

L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

O document referring to an oral disclosure, use, exhibition or other means

P document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

& document member of the same patent family

Date of the actual completion of the international search

20 August 2002

Date of mailing of the international search report

03/09/2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax: (+31-70) 340-3016

Authorized officer

Gettins, M

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 02/05115

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D239/34 C07D233/24 C07D333/20 C07D307/91 C07D209/14

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>DATABASE CROSSFIRE BEILSTEIN 'Online! Beilstein Institut zur Förderung der Chemischen Wissenschaften, Frankfurt am Main, DE; (3-ethoxy-benzyl)-(4-ethoxy-phenethyl)-ami ne), IDE, BUCK: Database accession no. 3365864 XP002209139 abstract & J.AM.CHEM.SOC., vol. 59, 1937, page 726ff</p> <p style="text-align: center;">--- -/-</p>	1



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- * & * document member of the same patent family

Date of the actual completion of the international search

20 August 2002

Date of mailing of the international search report

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Gettins, M

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 02/05115

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DATABASE CROSSFIRE BEILSTEIN 'Online! Beilstein Institut zur Förderung der Chemischen Wissenschaften, Frankfurt am Main, DE; (3-ethoxy-benzyl)-(3-ethoxy-phenethyl)-ami ne), Database accession no. 3365863 XP002209140 abstract & IDE, BUCK: J.AM.CHEM.SOC, vol. 59, 1937, page 726ff -----	1
X	DATABASE CROSSFIRE BEILSTEIN 'Online! Beilstein Institut zur Förderung der Chemischen Wissenschaften, Frankfurt am Main, DE; (3-ethoxy-benzyl)-(2-ethoxy-phenethyl)-ami ne), IDE, BUCK: Database accession no. 3380345 XP002209141 abstract & J.AM.CHEM.SOC., vol. 59, 1937, page 726ff -----	1

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US 02/05115

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

Although claims 22-28, 45 and 46 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☒ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:

see FURTHER INFORMATION sheet PCT/ISA/210
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Present claims 1-11, and 21-49 relate to an extremely large number of possible compounds due to the references to "optionally substituted". Support within the meaning of Article 6 PCT and/or disclosure within the meaning of Article 5 PCT is to be found, however, for only a very small proportion of the compounds claimed. In the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search over the whole of the claimed scope is impossible. Consequently, the search has been carried out for those parts of the claims which appear to be supported and disclosed, namely those parts relating to the compounds where optionally substituted is as defined on pages 6-8.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 02/05115

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 0034242	A	15-06-2000	AU 2356200 A	26-06-2000
			EP 1149078 A1	31-10-2001
			US 2002103382 A1	01-08-2002
			US 2002103383 A1	01-08-2002
			WO 0034242 A1	15-06-2000
US 6187805	B1	13-02-2001	GB 2341549 A	22-03-2000